



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07K 14/47, C12N 15/12, C07K 16/18, G01N 33/50, 33/53, C12Q 1/68, A61K 38/17	A2	(11) International Publication Number: WO 00/52047 (43) International Publication Date: 8 September 2000 (08.09.00)
(21) International Application Number: PCT/US00/05452 (22) International Filing Date: 3 March 2000 (03.03.00) (30) Priority Data: 09/263,022 5 March 1999 (05.03.99) US (71) Applicant: MILLENNIUM PHARMACEUTICALS, INC. [US/US]; 75 Sidney Street, Cambridge, MA 02139 (US). (72) Inventor: MCCARTHY, Sean; 62 Commonwealth Avenue, Apartment #4, Boston, MA 02116 (US). (74) Agents: MANDRAGOURAS, Amy, E.; Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 (US) et al.		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: HUMAN DICKKOPF-RELATED PROTEIN AND NUCLEIC ACID MOLECULES AND USES THEREFOR (57) Abstract Novel Dkk and Dkk-related polypeptides, proteins, and nucleic acid molecules are disclosed. In addition to isolated, full-length Dkk and Dkk-related proteins, the invention further provides isolated fusion proteins, antigenic peptides and antibodies. The invention also provides Dkk and Dkk-related nucleic acid molecules, recombinant expression vectors containing a nucleic acid molecule of the invention, host cells into which the expression vectors have been introduced and non-human transgenic animals in which a Dkk and Dkk-related gene has been introduced or disrupted. Diagnostic, screening and therapeutic methods utilizing compositions of the invention are also provided.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

HUMAN DICKKOPF-RELATED PROTEIN AND NUCLEIC ACID MOLECULES AND USES THEREFOR

Background of the Invention

5 Secreted proteins play an integral role in the formation, differentiation, and maintenance of cells in multicellular organisms. For instance, secretory proteins are known in the art to be involved in signaling between cells which are not in direct contact. Such secreted signaling molecules are particularly important in the development of vertebrate tissue during embryogenesis as well as in the maintenance of
10 the differentiated state of adult tissues. For example, inductive interactions that occur between neighboring cell layers and tissues in the developing embryo are largely dependent on the existence and regulation of secreted signaling molecules. In inductive interactions, biochemical signals secreted by one cell population influence the developmental fate of a second cell population, typically by altering the fate of the
15 second cell population. For example, the Wnt proteins are now recognized as one of the major families of developmentally important signaling molecules in organisms ranging from *Drosophila* to mice.

 The Wnt gene family encode a large class of secreted proteins related to the *Int1/Wnt1* proto-oncogene and *Drosophila* wingless ("Wg"), a *Drosophila* Wnt1
20 homologue, (Cadigan *et al.* (1997) *Genes & Development* 11:3286-3305). Wnts are expressed in a variety of tissues and organs and are required for many developmental processes, including segmentation in *Drosophila*, endoderm development in *Caenorhabditis elegans*, establishment of limb polarity, neural crest differentiation, kidney morphogenesis, sex determination, and brain development in mammals
25 (reviewed in Parr and McMahon (1994) *Curr. Opinion Genetics & Devel.* 4:523-528; Cadigan and Nusse, *supra*).

 Recent studies in diverse organisms have led to identification of several components of the Wnt signal transduction pathway in responding cells (Cadigan and Nusse, *supra*). Wnt signals are transduced by the Frizzled ("Fz") family of seven
30 transmembrane domain receptors (Bhanot *et al.* (1996) *Nature* 382:225-230). The resulting signal leads to the activation of the cytoplasmic protein Dishevelled (Dsh) and

stabilization of Armadillo/ β -catenin (Perrimon (1994) *Cell* 76:781-784). Negative regulators of the Wnt pathway include glycogen synthase kinase 3 (GSK3)/shaggy (Perrimon, *supra*), the tumor suppressor gene product adenomatous polyposis coli (APC) (Gumbiner (1997) *Curr. Biol.* 7:R443-436) and a novel protein, called Axin
5 (Zeng *et al.* (1997) *Cell* 90:181-192). In the absence of a Wnt ligand, these proteins promote phosphorylation and then degradation of β -catenin, whereas Wnt signaling inactivates GSK3, thus preventing β -catenin degradation. As a result, β -catenin is translocated to the nucleus, where it forms a complex with TCF transcription factors and activates target gene expression (Cadigan and Nusse, *supra*). Deregulation of this
10 pathway can lead to carcinogenesis (reviewed by Gumbiner, *supra*), emphasizing the long-recognized connection between Wnts, normal development and cancer. This connection has been further established recently with the identification the c-Myc proto-oncogene as a target of Wnt signaling (He *et al.* (1998) *Science* 281:1509-3512).

While the outcome of Wnt signaling may be influenced by multiple intracellular
15 regulatory mechanisms, recent studies have identified several classes of secreted factors which can modulate Wnt action outside of the cell. These include Cerberus, a secreted Wnt inhibitor implicated in head development (Bouwmeester *et al.* (1996) *Nature* 382:595-601), and a family of proteins related to the extracellular domain of Frizzled. These Frizzled-related proteins ("FRPs") (Rattner *et al.* (1997) *Proc. Natl. Acad. Sci.*
20 *USA* 94:2859-2863), also known as secreted apoptosis-related proteins ("SARPs"), are encoded by several independently discovered genes including FrzA/FRP1, SDF5/FRP2, FrzB/FRP3, FRP4 and Sizzled (Melkonyan *et al.* (1997) *Proc. Natl. Acad. Sci. USA* 94:13636-13641; Finch *et al.* (1997) *Proc. Natl. Acad. Sci. USA* 94:6770-6775; Wang *et al.* (1997) *Cell* 88:747-766; Leyns *et al.* (1997) *Cell* 88:747-756; Mayr *et al.* (1997)
25 *Mech. Dev.* 63:109-325; and Salic *et al.* (1997) *Development* 124:4739-4748). These proteins inhibit the ability of Xwnt8 to induce a secondary axis in frog embryos (for review see Zorn (1997) *Curr. Biol.* 7:R501-504), and are thought to compete for binding of Wnt ligands to the Frizzled receptors. Data on binding of certain FRPs to Xwnt8 (Wang *et al.*, (1997) *Biochem. Biophys. Res. Comm.* 236:502-504; and Leyns *et al.*,
30 *supra*) and Wg corroborate this notion (Rattner *et al.*, *supra*).

It is now recognized that many of these families of signaling molecules have a dual role to play in both the development of an organism as well as in promoting or maintaining the differentiated state of tissues in the adult animal. Furthermore, major families of signaling molecules have been implicated in controlling proliferation of cells in mature adult tissue, for example, during normal cell turnover in the adult organism as well as in tissue regeneration activated as a result of damage to the adult tissue. Given the important role of these signalling molecules such as the Wnts and FRPs in both developing and adult tissues, there exists a need for identifying novel modulators of such molecules for use in regulating a variety of cellular processes.

Summary of the Invention

The present invention is based, at least in part, on the discovery of nucleic acid molecules which encode a novel family of secreted human proteins, referred to herein as the human Dickkopf proteins or "hDkks" (formerly referred to as the "Cysteine-Rich Secreted Proteins", "CRSPs", "CRISPYs", or "CRSP proteins"). The Dkk molecules of the present invention are useful as modulating agents in regulating a variety of cellular processes. Accordingly, in one aspect, this invention provides isolated nucleic acid molecules encoding Dkk proteins or biologically active portions thereof, as well as nucleic acid fragments suitable as primers or hybridization probes for the detection of Dkk-encoding nucleic acids. In another aspect, this invention provides isolated nucleic acid molecules encoding Dkk-related proteins (*e.g.*, Soggy proteins) or biologically active portions thereof, as well as nucleic acid fragments suitable as primers or hybridization probes for the detection of Dkk- or Soggy-encoding nucleic acids.

In one embodiment, a Dkk nucleic acid molecule is 60% homologous to the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:3, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452 or complement thereof. In yet another embodiment, a Dkk nucleic acid molecule is 80% homologous to the nucleotide sequence shown in SEQ ID NO:4, SEQ ID NO:6, or a complement thereof. In yet another embodiment, a Dkk nucleic acid molecule is 60% homologous to the nucleotide sequence shown in SEQ ID NO:7, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as

Accession Number 98633, or a complement thereof. In yet another embodiment, a Dkk nucleic acid molecule is 85% homologous to the nucleotide sequence shown in SEQ ID NO:7, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or a complement thereof. In yet

5 another embodiment, a Dkk nucleic acid molecule is 70% homologous to the nucleotide sequence shown in SEQ ID NO:20, SEQ ID NO:22, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a complement thereof. In yet another embodiment, a nucleic acid molecule of the present invention (*e.g.*, a Dkk-related nucleic acid molecule) is 90% homologous to the
10 nucleotide sequence shown in SEQ ID NO:13, SEQ ID NO:15, or a complement thereof.

In a preferred embodiment, an isolated Dkk nucleic acid molecule has the nucleotide sequence shown SEQ ID NO:3, or a complement thereof. In another embodiment, a Dkk nucleic acid molecule further comprises nucleotides 1-37 of SEQ ID NO:1. In yet another preferred embodiment, a Dkk nucleic acid molecule further
15 comprises nucleotides 1088-2479 of SEQ ID NO:1. In another preferred embodiment, an isolated Dkk nucleic acid molecule has the nucleotide sequence shown in SEQ ID NO:1.

In another preferred embodiment, an isolated Dkk nucleic acid molecule has the nucleotide sequence shown SEQ ID NO:6, or a complement thereof. In another
20 embodiment, a Dkk nucleic acid molecule further comprises nucleotides 1-124 of SEQ ID NO:4. In yet another preferred embodiment, a Dkk nucleic acid molecule further comprises nucleotides 797-848 of SEQ ID NO:4. In another preferred embodiment, an isolated Dkk nucleic acid molecule has the nucleotide sequence shown in SEQ ID NO:4.

In another preferred embodiment, an isolated Dkk nucleic acid molecule has the
25 nucleotide sequence shown SEQ ID NO:9, or a complement thereof. In another embodiment, a Dkk nucleic acid molecule further comprises nucleotides 1-108 of SEQ ID NO:7. In yet another preferred embodiment, a Dkk nucleic acid molecule further comprises nucleotides 907-1536 of SEQ ID NO:7. In another preferred embodiment, an isolated Dkk nucleic acid molecule has the nucleotide sequence shown in SEQ ID NO:7.

In another preferred embodiment, an isolated Dkk nucleic acid molecule has the nucleotide sequence shown SEQ ID NO:22, or a complement thereof. In another embodiment, a Dkk nucleic acid molecule further comprises nucleotides 1-723 of SEQ ID NO:20. In yet another preferred embodiment, a Dkk nucleic acid molecule further
5 comprises nucleotides 1501-3687 of SEQ ID NO:20. In yet another preferred embodiment, an isolated Dkk nucleic acid molecule has the nucleotide sequence shown in SEQ ID NO:20.

In another preferred embodiment, an isolated nucleic acid molecule of the present invention (*e.g.*, a Dkk-related nucleic acid molecule) has the nucleotide sequence
10 shown SEQ ID NO:15, or a complement thereof. In another embodiment, a nucleic acid molecule further comprises nucleotides 1-74 of SEQ ID NO:13. In yet another preferred embodiment, a nucleic acid molecule further comprises nucleotides 801-928 of SEQ ID NO:13. In yet another preferred embodiment, an isolated nucleic acid molecule has the nucleotide sequence shown in SEQ ID NO:13.

15 In another embodiment, a Dkk nucleic acid molecule includes a nucleotide sequence encoding a protein having an amino acid sequence sufficiently homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:21. In another preferred embodiment, a Dkk nucleic acid molecule includes a nucleotide sequence encoding a protein having an amino acid sequence at least 60%
20 homologous to the amino acid sequence of SEQ ID NO:2. In yet another preferred embodiment, a Dkk nucleic acid molecule includes a nucleotide sequence encoding a protein having an amino acid sequence at least 60% homologous to the amino acid sequence of SEQ ID NO:5. In yet another preferred embodiment, a Dkk nucleic acid molecule includes a nucleotide sequence encoding a protein having an amino acid
25 sequence at least 60% homologous to the amino acid sequence of SEQ ID NO:8. In yet another preferred embodiment, a Dkk nucleic acid molecule includes a nucleotide sequence encoding a protein having an amino acid sequence at least 75% homologous to the amino acid sequence of SEQ ID NO:8. In yet another preferred embodiment, a Dkk nucleic acid molecule includes a nucleotide sequence encoding a protein having an
30 amino acid sequence at least 65% homologous to the amino acid sequence of SEQ ID NO:21. In another embodiment, a nucleic acid molecule of the present invention (*e.g.*, a

Dkk-related nucleic acid molecule) includes a nucleotide sequence encoding a protein having an amino acid sequence sufficiently homologous to the amino acid sequence of SEQ ID NO:14 (*e.g.*, encodes a protein having an amino acid sequence which is 60% homologous to the amino acid sequence of SEQ ID NO:14).

5 In another embodiment, an isolated nucleic acid molecule of the present invention encodes a Dkk protein which includes a signal sequence and at least one cysteine-rich region, and is secreted. In another embodiment, an isolated nucleic acid molecule of the present invention encodes a Dkk protein which includes a signal sequence and a cysteine-rich region, wherein the cysteine-rich region comprises at least
10 one cysteine-rich domain, and is secreted. In yet another embodiment, a Dkk nucleic acid molecule encodes a Dkk protein and is a naturally occurring nucleotide sequence.

In another embodiment, an isolated nucleic acid molecule of the present invention encodes a Dkk-related protein (*e.g.*, a Soggy protein) which includes a signal sequence, lacks cysteine-rich domains, and is secreted. In another embodiment, an
15 isolated nucleic acid molecule of the present invention encodes a Dkk-related protein (*e.g.*, a Soggy protein) which includes a signal sequence and a Soggy domain, and is secreted. In yet another embodiment, a nucleic acid molecule of the present invention encodes a Dkk-related protein and is a naturally occurring nucleotide sequence.

Another embodiment of the invention features nucleic acid molecules which
20 specifically detect Dkk nucleic acid molecules relative to nucleic acid molecules encoding non-Dkk proteins (or specifically detect Dkk-related nucleic acid molecules). For example, in one embodiment, a nucleic acid molecule hybridizes under stringent conditions to a nucleic acid molecule consisting of nucleotides 470-2479 of nucleotide sequence shown in SEQ ID NO:1, to nucleotides 1-475 of nucleotide sequence shown in
25 SEQ ID NO:4, or to nucleotides 1-600 of nucleotide sequence shown in SEQ ID NO:7, or hybridizes under stringent conditions to the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, to the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or to the nucleotide sequence of the DNA insert of the plasmid deposited with
30 ATCC as Accession Number _____. In another embodiment, the nucleic acid molecule is at least 500 nucleotides in length and hybridizes under stringent conditions to a

nucleic acid molecule comprising the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20 or a complement thereof.

Another embodiment of the invention provides an isolated nucleic acid molecule which is antisense to the coding strand of a Dkk nucleic acid or Dkk-related nucleic acid. Another embodiment of the invention provides an isolated nucleic acid molecules in a form suitable for expression of mRNA. In another embodiment, the isolated nucleic acid molecules are in a form suitable for expression of protein. In yet another embodiment, the isolated nucleic acid molecules are free from vector sequences.

Another aspect of the invention provides a vector comprising a Dkk nucleic acid molecule or Dkk-related nucleic acid molecule. In certain embodiments, the vector is a recombinant expression vector. In another embodiment, the invention provides a host cell containing a vector of the invention. The invention also provides a method for producing a Dkk protein or Dkk-related protein by culturing in a suitable medium, a host cell of the invention containing a recombinant expression vector such that a Dkk protein or Dkk-related protein is produced.

Another aspect of this invention features isolated or recombinant Dkk proteins and polypeptides or Dkk-related proteins and polypeptides. In one embodiment, an isolated Dkk protein has a signal sequence and a cysteine-rich region which comprises two cysteine-rich domains, and is secreted. In another embodiment, an isolated Dkk protein has an amino acid sequence sufficiently homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:21. In a preferred embodiment, a Dkk protein has an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ ID NO:2. In another preferred embodiment, a Dkk protein has an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ ID NO:5. In another preferred embodiment, a Dkk protein has an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ ID NO:8. In another preferred embodiment, a Dkk protein has an amino acid sequence at least about 75% homologous to the amino acid sequence of SEQ ID NO:8. In another preferred embodiment, a Dkk protein has an amino acid sequence at least about 65% homologous to the amino acid sequence of SEQ ID NO:21. In another embodiment, a Dkk protein has the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID

NO:8, or SEQ ID NO:21. In another preferred embodiment, a protein of the present invention has an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ ID NO:14. In another embodiment, a protein has the amino acid sequence of SEQ ID NO:14.

5 Another embodiment of the invention features an isolated Dkk protein which is encoded by a nucleic acid molecule having a nucleotide sequence at least about 60% homologous to a nucleotide sequence of SEQ ID NO:1, or a complement thereof. Another embodiment of the invention features an isolated Dkk protein which is encoded by a nucleic acid molecule having a nucleotide sequence at least about 80% homologous
10 to a nucleotide sequence of SEQ ID NO:4, or a complement thereof. Another embodiment of the invention features an isolated Dkk protein which is encoded by a nucleic acid molecule having a nucleotide sequence at least about 60% homologous to a nucleotide sequence of SEQ ID NO:7, or a complement thereof. Another embodiment of the invention features an isolated Dkk protein which is encoded by a nucleic acid
15 molecule having a nucleotide sequence at least about 85% homologous to a nucleotide sequence of SEQ ID NO:7, or a complement thereof. Another embodiment of the invention features an isolated Dkk protein which is encoded by a nucleic acid molecule having a nucleotide sequence at least about 70% homologous to a nucleotide sequence of SEQ ID NO:20, or a complement thereof. Another embodiment of the invention
20 features an isolated protein which is encoded by a nucleic acid molecule having a nucleotide sequence at least about 90% homologous to a nucleotide sequence of SEQ ID NO:13, or a complement thereof. This invention further features an isolated protein which is encoded by a nucleic acid molecule having a nucleotide sequence which hybridizes under stringent hybridization conditions to a nucleic acid molecule
25 comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, or a complement thereof.

 The proteins of the present invention, or biologically active portions thereof, can be operatively linked to a non-Dkk polypeptide or non-Dkk-related polypeptide to form fusion proteins. The invention further features antibodies that specifically bind Dkk or
30 Dkk-related proteins, such as monoclonal or polyclonal antibodies. In addition, the

proteins or biologically active portions thereof can be incorporated into pharmaceutical compositions, which optionally include pharmaceutically acceptable carriers.

In another aspect, the present invention provides a method for detecting Dkk expression (or the expression of a Dkk-related molecule) in a biological sample by
5 contacting the biological sample with an agent capable of detecting a nucleic acid molecule, protein or polypeptide of the present invention such that the presence of a Dkk (of Dkk-related) nucleic acid molecule, protein or polypeptide is detected in the biological sample.

In another aspect, the present invention provides a method for detecting the
10 presence of a Dkk activity (or Dkk-related activity) in a biological sample by contacting the biological sample with an agent capable of detecting an indicator of Dkk activity (or Dkk-related activity) such that the presence of the activity is detected in the biological sample.

In another aspect, the invention provides a method for modulating Dkk activity
15 (or Dkk-related activity) comprising contacting the cell with an agent that modulates the activity such that the activity in the cell is modulated. In one embodiment, the agent inhibits Dkk activity (or Dkk-related activity). In another embodiment, the agent stimulates Dkk activity (or Dkk-related activity). In one embodiment, the agent is an antibody that specifically binds to a Dkk (or Dkk-related) protein. In another
20 embodiment, the agent modulates expression of a protein (*e.g.*, a Dkk or a Dkk-related protein) by modulating transcription of a gene or translation of a mRNA of the present invention. In yet another embodiment, the agent is a nucleic acid molecule having a nucleotide sequence that is antisense to the coding strand of a mRNA or gene of the present invention.

25 In one embodiment, the methods of the present invention are used to treat a subject having a disorder characterized by aberrant expression or activity of a protein or nucleic acid of the invention by administering to the subject an agent which is a modulator of Dkk or a Dkk-related molecule. In one embodiment, the modulator is a Dkk or Dkk-related protein. In another embodiment the modulator is a Dkk or Dkk-
30 related nucleic acid molecule. In yet another embodiment, the modulator is an antibody peptide, peptidomimetic, or other small molecule. In a preferred embodiment, the

disorder characterized by aberrant protein or nucleic acid expression is a developmental, differentiative, or proliferative disorder.

The present invention also provides a diagnostic assay for identifying the presence or absence of a genetic alteration characterized by at least one of (i) aberrant
5 modification or mutation of a gene encoding a Dkk or Dkk-related protein; (ii) mis-regulation of said gene; and (iii) aberrant post-translational modification of a Dkk or Dkk-related protein, wherein a wild-type form of said gene encodes an protein with a Dkk or Dkk-related activity.

In another aspect the invention provides a method for identifying a compound
10 that binds to or modulates the activity of a Dkk or Dkk-related protein, by providing a indicator composition comprising a Dkk or Dkk-related protein having a biological activity, contacting the indicator composition with a test compound, and determining the effect of the test compound on the activity in the indicator composition to identify a compound that modulates the activity of a Dkk or Dkk-related protein.

15 Other features and advantages of the invention will be apparent from the following detailed description and claims.

Brief Description of the Drawings

Figure 1A-B depicts the cDNA sequence and predicted amino acid sequence of
20 human Dkk-3. The nucleotide sequence corresponds to nucleic acids 1 to 2479 of SEQ ID NO:1. The amino acid sequence corresponds to amino acids 1 to 350 of SEQ ID NO:2.

Figure 2 depicts the cDNA sequence and predicted amino acid sequence of
human Dkk-4. The nucleotide sequence corresponds to nucleic acids 1 to 848 of SEQ
25 ID NO:4. The amino acid sequence corresponds to amino acids 1 to 224 of SEQ ID NO:5.

Figure 3 depicts the cDNA sequence and predicted amino acid sequence of
human Dkk-1. The nucleotide sequence corresponds to nucleic acids 1 to 1536 of SEQ
ID NO:7. The amino acid sequence corresponds to amino acids 1 to 266 of SEQ ID
30 NO:8.

Figure 4A-B depicts the cDNA sequence and predicted amino acid sequence of full-length human Dkk-2. The nucleotide sequence corresponds to nucleic acids 1 to 3687 of SEQ ID NO:20. The amino acid sequence corresponds to amino acids 1 to 259 of SEQ ID NO:21.

5 *Figure 5A-B* depicts the cDNA sequence and predicted amino acid sequence of murine Dkk-3. The nucleotide sequence corresponds to nucleic acids 1 to 2380 of SEQ ID NO:16. The amino acid sequence corresponds to amino acids 1 to 349 of SEQ ID NO:17.

10 *Figure 6* depicts a multiple sequence alignment of the amino acid sequences of hDkk-1 (corresponding the SEQ ID NO:8), mDkk-1 (having Accession No. AF030433), *Xenopus* Dkk-1 ("xDkk-1") (having Accession No. AF030434), hDkk-2 (corresponding to SEQ ID NO:21), hDkk-3 (corresponding to SEQ ID NO:2), mDkk-3 (corresponding to SEQ ID NO:17), chicken Dkk-3 ("cDkk-3") (having Accession No. D26311), and hDkk-4 (corresponding to SEQ ID NO:5). The alignment was performed using the
15 ClustalW algorithm as implemented in the GCG program PILEUP. The alignment provides information regarding the relationship between the Dkk proteins of the instant invention. Predicted signal peptides are underlined, N-glycosylation sites are indicated by a thick bar, CRD-1 by an open box, CRD-2 by a shaded box. The proteolytic cleavage site within hDkk4 is indicated by an arrow.

20 *Figure 7* depicts the cDNA sequence and predicted amino acid sequence of human Soggy. The nucleotide sequence corresponds to nucleic acids 1 to 928 of SEQ ID NO:13. The amino acid sequence corresponds to amino acids 1 to 242 of SEQ ID NO:14.

25 *Figure 8* depicts the cDNA sequence and predicted amino acid sequence of murine Soggy-1. The nucleotide sequence corresponds to nucleic acids 1 to 835 of SEQ ID NO:26. The amino acid sequence corresponds to amino acids 1 to 230 of SEQ ID NO:27.

30 *Figure 9* is a schematic diagram illustrating the Dkk and Dkk-related proteins of the instant invention. The figure depicts the structural domains of the human Dkks and Soggy. Signal peptides are indicated by darkened boxes. The cysteine-rich domains of

a Dkk cysteine-rich region are depicted as CRD-1 and CRD-2. Branches indicate sites of N-glycosylation.

Figure 10 depicts a multiple sequence alignment of hSoggy-1 (corresponding to SEQ ID NO:14), murine Soggy-1 (corresponding to SEQ ID NO:27), hDkk-3 (corresponding to SEQ ID NO:2), and mDkk-3 (corresponding to SEQ ID NO:17). The alignment was generated as described in the legend to Figure 6. The alignment provides details regarding the relationship between the Dkk-3 and Soggy-1 proteins of the instant invention. Predicted signal peptides are underlined, N-glycosylation sites are indicated by a thick bar. CRD-1 and CRD-2 within Dkk-3 are indicated for reference by open and shaded boxes.

Figure 11 depicts a multiple sequence alignment of the carboxy-terminal cysteine-rich domains of hDkk-1, hDkk-2, hDkk-3, hDkk-4 with human colipase (having accession No. J02883). The carboxy-terminal cysteine-rich domains of the Dkk proteins are indicated by an open box. The alignment was generated using PILEUP (gap penalties of 12 for opening and 12 for extending). A minor adjustment was necessary since PILEUP inserts a single gap in hDkk-1 and hDkk-2 between Gly56 and Ser57, even with a gap opening penalty of 15. The conserved residues are indicated. The disulfide-bonding pattern typical for the colipase family and predicted for the Dkk family is indicated below the alignment.

Figure 12 is a schematic diagram depicting the relationship between the hDkk-3 nucleotide sequence (corresponding to SEQ ID NO:1) and those of RIG and RIG-like 7-1 (Accession Nos. U32331 and AF034208, respectively). Thick bars indicate regions of sequence identity between hDkk-3 and RIG or RIG-like 7-1 mRNAs. As between RIG and hDkk-3, there exists a short region of identity within the 3' untranslated regions of the mRNAs when the mRNAs are aligned in reverse orientation. As between hDkk-3 and RIG-like 7-1, there exists a longer region of identity, however, RIG-like 7-1 lacks a signal sequence and, accordingly, is not predicted to be secreted.

Detailed Description of the Invention

The present invention is based on the discovery of novel molecules, referred to herein as Dkk protein and nucleic acid molecules, which comprise a family of molecules

having certain conserved structural and functional features. The term "family" when referring to the protein and nucleic acid molecules of the invention is intended to mean two or more proteins or nucleic acid molecules having a common structural domain and having sufficient amino acid or nucleotide sequence homology as defined herein. Such
5 family members can be naturally-occurring and can be from either the same or different species. For example, a family can contain a first protein of human origin, as well as other, distinct proteins of human origin or alternatively, can contain homologues of non-human origin. Members of a family may also have common functional characteristics.

In one embodiment, a Dkk family member is identified based on the presence of
10 at least one "cysteine-rich domain" in the protein molecule or corresponding amino acid sequence. As defined herein, a "cysteine-rich domain" refers to a portion of a Dkk protein (*e.g.*, hDkk-3) which is rich in cysteine residues. In a preferred embodiment, a "cysteine-rich domain" is a protein domain having an amino acid sequence of about 45-85 amino acids of which preferably 10 amino acids are cysteine residues located at the
15 same relative amino acid position as the cysteine residues in human Dkk-3 having SEQ ID NO:2 (*e.g.*, amino acid residues 147-195 of SEQ ID NO:2). In another embodiment, a "cysteine-rich domain" has 30-100 amino acids, preferably about 35-95 amino acids, more preferably about 40-90 amino acids, more preferably about 50-80 amino acids, even more preferably about 55-75, 60-70, or 65 amino acids, of which at least about 3-
20 20, preferably about 5-15, or more preferably about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 amino acids are cysteine residues.

A preferred Dkk protein of the present invention has a first cysteine-rich domain ("CRD-1") referred to herein as an "amino-terminal cysteine-rich domain" or "N-terminal cysteine-rich domain" and a second cysteine-rich domain ("CRD-2"), referred
25 to herein as a "carboxy-terminal cysteine-rich domain" or "C-terminal cysteine-rich domain". As defined herein, an "amino-terminal cysteine-rich domain" is a protein domain having an amino acid sequence of about 45-55 amino acids of which preferably 10 amino acids are cysteine residues located at the same relative position as the cysteine residues in an amino-terminal cysteine-rich domain of human Dkk-3 having SEQ ID
30 NO:2 (*e.g.*, amino acid residues 147-195 of SEQ ID NO:2). In another embodiment, an "amino-terminal cysteine-rich domain" has 30-70, preferably 35-65, more preferably

about 40-60, and even more preferably about 46, 47, 48, 49, 50, 51, 52, 53, or 54 amino acids, of which at least about 3-20, preferably about 5-15, or more preferably about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 amino acids are cysteine residues. In a preferred embodiment, an amino-terminal cysteine-rich domain has the following

5 consensus sequence: C - X(2) - D - X(2) - C - X(5) - C - X(8-13) - C - X(2) - C - X(6) - C - X(5) - C - C - X(4) - C - X(4) - C (SEQ ID NO:23). The consensus sequences described herein are described according to standard Prosite Signature designation (*e.g.*, all amino acids are indicated according to their universal single letter designation; X designates any amino acid; X(n) designates any n amino acids, *e.g.*, X (2) designates any

10 2 amino acids; and [LIVM] indicates any one of the amino acids appearing within the brackets, *e.g.*, any one of L, I, V, or M, in the alternative, any one of Leu, Ile, Val, or Met.)

As defined herein, a "carboxy-terminal cysteine-rich domain" is a protein domain having an amino acid sequence of about 80-85 amino acids of which preferably 10

15 amino acids are cysteine residues located at the same relative position as the cysteine residues in a carboxy-terminal cysteine-rich domain of human Dkk-3 having SEQ ID NO:2 (*e.g.*, amino acid residues 201-284 of SEQ ID NO:2). In another embodiment, a "carboxy-terminal cysteine-rich domain" has 65-100, preferably 70-95, more preferably about 75-90, and even more preferably about 81, 82, 83, or 84 amino acids, of which at

20 least about 3-20, preferably about 5-15, or more preferably about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 amino acids are cysteine residues. In a preferred embodiment, a carboxy-terminal cysteine-rich domain has the following consensus sequence: C - X(4) - D - C - X(2) - G - X - C - C - X(8-10) - C - X - P - X(4) - G - X(2) - C - X(16-24) - C - X - C - X(2) - P - X(4) - G - X(2) - C - X(16-24) - C - X - C - X(2) -

25 G - L - X - C - X(10-17) - C (SEQ ID NO:24).

A preferred protein of the present invention is a hDkk-3 protein (human Dkk-3) containing an amino-terminal cysteine-rich domain including about amino acids 147-195 of SEQ ID NO:2, having 10 cysteine residues, and a carboxy-terminal cysteine-rich domain including about amino acids 201-284 of SEQ ID NO:2, having 10 cysteine

30 residues (the positions of the cysteine residues are depicted in Figure 6). In another embodiment, a hDkk-4 (human Dkk-4) protein contains an amino-terminal cysteine-rich

domain including about amino acids 41-90 of SEQ ID NO:5, having 10 cysteine residues, and a carboxy-terminal cysteine-rich domain including about amino acids 138-218 of SEQ ID NO:5, having 10 cysteine residues (the positions of the cysteine residues are depicted in Figure 6). In another embodiment, a hDkk-1 protein (human Dkk-1) contains an amino-terminal cysteine-rich domain including about amino acids 85-138 of SEQ ID NO:8, having 10 cysteine residues, and a carboxy-terminal cysteine-rich domain including about amino acids 182-263 of SEQ ID NO:8, having 10 cysteine residues (the positions of the cysteine residues are depicted in Figure 6). In another embodiment, a hDkk-2 protein (human Dkk-2) contains an amino-terminal cysteine-rich domain including about amino acids 78-127 of SEQ ID NO:21, having 10 cysteine residues, and a carboxy-terminal cysteine-rich domain including about amino acids 176-256 of SEQ ID NO:21, having 10 cysteine residues (the positions of the cysteine residues are depicted in Figure 6).

Alignment of the human Dkk proteins with human colipase (having Accession No. J02883) indicates that the carboxy-terminal cysteine-rich domains of the human Dkk proteins have a pattern of cysteines typical of colipase (Figure 11 and Avarind and Koonin, *supra*). Within colipase, these cysteine residues are involved in disulfide bonding which gives rise to a structure termed the "colipase fold". The "colipase fold" is typical of a range of small proteins which are involved in protein-protein interactions including, but not limited to the colipases, snake and scorpion toxins and protease inhibitors (Hubbard *et al.* (1997) *Nucleic Acids Res.* 25:236-239. These proteins have a series of short β strands with large connecting loops, which are held together by disulfide bonds. The disulfide-bonding pattern typical for colipase and predicted for the Dkk family is indicated below the alignment of Figure 11. Conserved hydrophobic residues between the Dkks and human colipase suggest that the Dkks, like the colipases, interact with lipids (*e.g.*, Leu51 of human colipase, SEQ ID NO:25 which corresponds to Leu271 of hDkk-3 (SEQ ID NO:2); Leu200 of hDkk-4 (SEQ ID NO:5); Leu243 of hDkk-1 (SEQ ID NO:8); and Leu237 of hDkk-2 (SEQ ID NO:21). The carboxy-terminal cysteine-rich domain of the Dkk family, may function in the membrane association of Dkk, which in turn may be required for the inhibition of Wnt secretion or Wnt:7 transmembrane receptor interaction. In addition, inhibition of Wnt function by

the Dkk family may be closely associated with the cell membrane and the carboxy-terminal cysteine-rich domain of the Dkk family may mediate this association.

Furthermore, the amino-terminal cysteine-rich domain of the Dkk family may directly interact with Wnt or its receptor. Accordingly, a preferred Dkk protein of the present invention comprises a carboxy-terminal cysteine-rich domain. In one embodiment, a Dkk protein comprising a carboxy-terminal cysteine-rich domain lacks the amino-terminal cysteine-rich domain.

In a preferred embodiment, the cysteine residues of a cysteine-rich domain are located at the same relative amino acid position as the cysteine residues in human Dkk-3 having SEQ ID NO:2. In another preferred embodiment, the cysteine residues of a cysteine-rich domain are located at the same relative position as the cysteine residues in a cysteine-rich domain of human Dkk-3 having SEQ ID NO:2. For example, as shown in Figure 6, human Dkk-4 has at least about 10 cysteine residues located at the same relative amino acid position as the cysteine residues in human Dkk-3 having SEQ ID NO:2 (*e.g.*, cys151 in Dkk-4, SEQ ID NO:5, is located at the same relative amino acid position as cys214 in Dkk-3, SEQ ID NO:2; cys156 in Dkk-4, SEQ ID NO:5, is located at the same relative amino acid position as cys219 in Dkk-3, SEQ ID NO:2; and cys157 in Dkk-4, SEQ ID NO:5, is located at the same relative amino acid position as cys220 in Dkk-3, SEQ ID NO:2). Similarly, as shown in Figure 6, Dkk-1 has at least about 10 cysteine residues located at the same relative amino acid position as the cysteine residues in human Dkk-3 having SEQ ID NO:2. As also shown in Figure 6, Dkk-2 has at least about 10 cysteine residues located at the same relative amino acid position as the cysteine residues in human Dkk-3 having SEQ ID NO:2. Table I sets forth at least 20 cysteine residues in each of hDkk-4, hDkk-1, and hDkk-2 which are located in the same relative position as 20 cysteine residues in hDkk-3.

Table I:

cysteine	aa position in hDkk-3	aa position in hDkk-4	aa position in hDkk-1	aa position in hDkk-2
1	147	41	85	78
2	153	47	91	84
3	159	53	97	90
4	168	63	111	100
5	171	66	114	103
6	178	73	121	110
7	184	79	127	116
8	185	80	128	117
9	190	85	133	122
10	195	90	138	127
11	208	145	189	183
12	214	151	195	189
13	219	156	200	194
14	220	157	201	195
15	231	166	210	204
16	241	176	220	214
17	265	194	237	231
18	267	196	239	233
19	273	202	245	239
20	284	218	263	256

The first 10 rows of Table I contain 10 cysteine residues that are included within
5 the first, or amino-terminal, cysteine-rich domain of each of hDkks-3, -4, -1, and -2.
The last 10 rows of Table I contain 10 cysteine residues that are included within the
second, or carboxy-terminal, cysteine-rich domain of each of hDkks-3, -4, -1, and -2.

Preferred Dkk proteins have more than one cysteine-rich domain, more preferably have at least two cysteine-rich domains and, thus, have a cysteine-rich region. As used herein, the term "cysteine-rich region" refers to a protein domain which includes at least two cysteine-rich domains and has an amino acid sequence of about 120-200 amino acid residues of which at least about 20 of the amino acids are cysteine residues. In another embodiment, a "cysteine-rich region" has preferably about 140-180 amino acid residues, and even more preferably at least about 135-175 amino acids of which at least about 10-30, preferably about 15-20, and more preferably about 16, 17, 18, or 19 of the amino acids are cysteine residues. In a preferred embodiment, a cysteine-rich region is located in the C-terminal region of a Dkk protein. For example, in one embodiment, a hDkk-3 protein contains a cysteine rich region containing about amino acids 147-284 of SEQ ID NO:2, having 20 cysteine residues at the positions indicated in Figure 6. In another embodiment, a hDkk-4 protein contains a cysteine rich region containing about amino acids 41-218 of SEQ ID NO:5, having 20 cysteine residues at the positions indicated in Figure 6. In another embodiment, a hDkk-1 protein contains a cysteine rich region containing about amino acids 85-263 of SEQ ID NO:8, having 20 cysteine residues at the positions indicated in Figure 6. In another embodiment, a hDkk-2 protein contains a cysteine rich region containing about amino acids 78-256 of SEQ ID NO:21, having 20 cysteine residues at the positions indicated in Figure 6.

In another embodiment, in addition to cysteine-rich domains, the cysteine-rich region contains a spacer region which separates the first and second cysteine-rich domains. As used herein, the "spacer region" refers to amino acid residues which are located between the first and second cysteine-rich domains of a cysteine-rich region and includes amino acid residues located C-terminal to the first cysteine-rich domain and N-terminal to the second cysteine-rich domain. As defined herein, a "spacer region" refers to a protein domain of about 5-70 amino acids, preferably about 10-65 amino acids, more preferably about 15-60 amino acids, even more preferably about 20-55 amino acids, and even more preferably about 25-50, 30-45 or 35-40 amino acids. For example, hDkk-3 protein contains a spacer region of about amino acids 196-200 of SEQ ID NO:2; hDkk-4 protein contains a spacer region of about amino acids 91-137 of SEQ ID NO:5; hDkk-1 protein contains a spacer region of about amino acids 139-181 of SEQ ID NO:8;

and hDkk-2 protein contains a spacer region of about amino acids 128-175 of SEQ ID NO:21. The spacer regions of hDkk-1, hDkk-2 and hDkk-4 are remarkably conserved in length (*e.g.*, the spacer region of hDkk-1 consists of 43 amino acid residues, the spacer region of hDkk-2 consists of 48 amino acid residues and the spacer region of hDkk-4 consists of 47 amino acid residues, suggesting that the close proximity of CRD-1 and CRD-2 is important in Dkk function. Accordingly, in one embodiment, the spacer region functions to spatially restrict the separation of CRD-1 from CRD-2.

In another embodiment of the invention, the Dkk protein has at least one cysteine-rich domain, preferably a cysteine-rich region, and a signal sequence. As used herein, a "signal sequence" refers to a peptide containing about 18-24 amino acids which occurs at the N-terminus of secretory and integral membrane proteins and which contains at least about 40-70% hydrophobic amino acid residues (*e.g.*, alanine, valine, leucine, isoleucine, phenylalanine, tyrosine, tryptophan, or proline). In another embodiment, a signal sequence contains at least about 8-34, 9-33, 10-32, 11-31, 12-30, 13-29, 14-28 amino acid residues, preferably about 15-27 amino acid residues, more preferably about 16-26 amino acid residues, more preferably about 17-25 amino acid residues, and more preferably about 18-24, 19-23, 20-22, or 21 amino acid residues, and has at least about about 50-65%, and more preferably about 55-60% hydrophobic amino acid residues (*e.g.*, alanine, valine, leucine, isoleucine, phenylalanine, tyrosine, tryptophan, or proline). Such a "signal sequence", also referred to in the art as a "signal peptide", serves to direct a protein containing such a sequence to a lipid bilayer. For example, in one embodiment, a hDkk-3 protein contains a signal sequence of about amino acids 1-23 of SEQ ID NO:2. In another embodiment, a hDkk-4 protein contains a signal sequence of about amino acids 1-19 of SEQ ID NO:5. In another embodiment, a hDkk-1 protein contains a signal sequence of about amino acids 1-20 of SEQ ID NO:8. In another embodiment, a hDkk-2 protein contains a signal sequence of about amino acids 1-33 of SEQ ID NO:21. A preferred Dkk protein of the present invention is a human protein (*e.g.*, encoded by a nucleotide sequence corresponding to a naturally-occurring human gene).

Accordingly, one embodiment of the invention features a Dkk protein having at least one cysteine-rich domain, preferably at least one cysteine-rich region. Another embodiment features a Dkk protein having at least one cysteine-rich region, wherein the cysteine-rich region includes at least one cysteine-rich domain. Another embodiment
5 features a Dkk protein having at least one cysteine-rich region, wherein the cysteine-rich region includes at least two cysteine-rich domains. Another embodiment features a protein or domain within a protein having 20, 30, 40, 50, 60, 70, 80, 90, 95, or 99% homology to a cysteine-rich domain of a Dkk protein of the invention (*e.g.*, hDkk-3, hDkk-4, hDkk-1, or hDkk-2).

10 Yet another embodiment of the invention features a Dkk protein having at least one cysteine-rich domain, preferably at least one cysteine-rich region and a signal peptide. Another embodiment features a Dkk protein having at least one cysteine-rich domain, preferably at least one cysteine-rich region and a signal peptide, wherein the cysteine-rich region includes at least two cysteine-rich domains. Another embodiment
15 features a Dkk protein having at least one cysteine-rich domain, preferably at least one cysteine-rich region and a signal peptide, wherein the cysteine-rich region includes at least two cysteine-rich domains and a spacer.

Yet another aspect of the invention features Dkk proteins having domains and/or regions which are conserved among a subset of Dkk proteins but are not necessarily
20 conserved among all Dkk family members. In one embodiment, a Dkk protein (*e.g.*, Dkk-3) has an "extended N-terminal region" which is extended in length as compared to, for example, the "N-terminal regions" of other Dkk family members (*e.g.*, Dkk-4, Dkk-1, and Dkk-2). As defined herein, an "N-terminal region" of a Dkk proteins consists of amino acid residues found between the signal peptide and CRD-1 of a Dkk protein.
25 Preferably, the first amino acid residue of an N-terminal region of Dkk is the first residue of a mature Dkk protein and the last residue of an N-terminal region of Dkk is the residue preceeding the first cysteine residue of CRD-1. In a preferred embodiment, an N-terminal region is about 1-20 amino acid residues in length, preferably about 21-30, 31-40, 41-50, 51-60, 61-70, 71-80, 81-90, 91-100, 101-110, 111-120, 121-130, 131-
30 140, 141-150, 151-160 or more amino acid residues in length. In contrast, an "extended N-terminal region" is at least about 71-80, 81-90, 91-100, 101-110, 111-120, 121-130,

131-140, 141-150, 151-160 or more amino acid residues in length. For example, in one embodiment, a hDkk-4 protein includes an "N-terminal region" of about amino acids 20-40 of SEQ ID NO:5 (21 amino acid residues in length). In another embodiment, a hDkk-1 protein includes an N-terminal region of about amino acids 21-84 of SEQ ID NO:8 (64 amino acid residues in length). In another embodiment, a hDkk-2 protein includes an "N-terminal region" of about amino acids 34-77 of SEQ ID NO:21 (44 amino acid residues in length). In another embodiment, a hDkk-3 protein has an "extended N-terminal region" of about amino acids 23-146 of SEQ ID NO:2 (124 amino acid residues in length).

10 In another embodiment, a Dkk protein (*e.g.*, Dkk-3) has an "acidic C-terminal region" which includes amino acid residues found C-terminal to CRD-2 of a Dkk protein. Preferably, the first amino acid residue of an acidic C-terminal region is the residue following the last cysteine of CRD-2 and the last residue of an acidic C-terminal region is the last residue of a Dkk protein. In a preferred embodiment, an acidic C-terminal
15 region is about 65-66 amino acid residues in length and has about 27-25% acidic amino acid residues (*e.g.*, glutamic acid or aspartic acid). In another preferred embodiment, an acidic C-terminal region is about 55-80 amino acid residues in length, preferably about 60-75 amino acid residues in length, and more preferably about 64-70 amino acid residues in length and has about 21-35% acidic amino acid residues, preferably about
20 23-33% acidic amino acid residues, and more preferably about 25-31% acidic amino acid residues. Preferably, an acidic C-terminal region is involved in protein-protein interactions. For example, in one embodiment, a hDkk-3 protein has an acidic C-terminal region from about amino acids 285-350 of SEQ ID NO:2.

Preferred Dkk molecules of the present invention have an amino acid sequence
25 sufficiently homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:21. As used herein, the term "sufficiently homologous" refers to a first amino acid or nucleotide sequence which contains a sufficient or minimum number of identical or equivalent (*e.g.*, an amino acid residue which has a similar side chain) amino acid residues or nucleotides to a second amino acid or
30 nucleotide sequence such that the first and second amino acid or nucleotide sequences share common structural domains and/or a common functional activity. For example,

amino acid or nucleotide sequences which share common structural domains have at least about 40% homology, preferably 50% homology, more preferably 60%-70% homology across the amino acid sequences of the domains and contain at least one, preferably two, more preferably three, and even more preferably four, five or six structural domains, are defined herein as sufficiently homologous. Furthermore, amino acid or nucleotide sequences which share at least 40%, preferably 50%, more preferably 60, 70, or 80% homology and share a common functional activity are defined herein as sufficiently homologous.

As used interchangeably herein, a "Dkk activity", "biological activity of Dkk" or "functional activity of Dkk", refers to an activity exerted by a Dkk protein, polypeptide or nucleic acid molecule (*e.g.*, an activity on a Dkk responsive cell) as determined *in vivo*, or *in vitro*, according to standard techniques. In one embodiment, a Dkk activity is a direct activity, such as an association with a Dkk-target molecule. As used herein, a "target molecule" is a molecule with which a Dkk protein binds or interacts in nature, such that Dkk-mediated function is achieved. A Dkk target molecule can be a non-Dkk molecule or a Dkk protein or polypeptide of the present invention. In an exemplary embodiment, a Dkk target molecule is a membrane-bound protein (*e.g.*, a cell-surface receptor or "Dkk receptor") or a modified form of such a protein which has been altered such that the protein is soluble (*e.g.*, recombinantly produced such that the protein does not express a membrane-binding domain). In another embodiment, a Dkk target is a second soluble protein molecule (*e.g.*, a "Dkk binding partner" or "Dkk substrate"). In such an exemplary embodiment, a Dkk binding partner can be a second soluble non-Dkk protein or a second Dkk protein molecule of the present invention. Alternatively, a Dkk activity is an indirect activity, such as a cellular signaling activity mediated by interaction of the Dkk protein with a second protein (*e.g.*, a Dkk receptor). As used herein, the term "Dkk receptor" refers to a protein or protein complex, to which a Dkk protein, *e.g.*, human Dkk, can bind. A receptor can be a cell surface receptor, *e.g.*, a peptide, growth factor, or nuclear hormone receptor. Dkk receptors can be isolated by methods known in the art and further described herein. Interaction of a Dkk protein with a Dkk receptor can result in transduction of a signal from the cell surface to the nucleus. The signal transduced can be, an increase in intracellular calcium, an increase in

phosphatidylinositol or other molecule, and can result in, *e.g.*, in phosphorylation of specific proteins, a modulation of gene transcription and any of the other biological activities set forth herein.

- In a preferred embodiment, a Dkk activity is at least one or more of the following
- 5 activities: (i) interaction of a Dkk protein with and/or binding to a second molecule, (*e.g.*, a protein, such as a Dkk receptor, a soluble form of a Dkk receptor, a receptor for a member of the *wnt* family of signaling proteins, or a non-Dkk signaling molecule, for example, a lipid included in a cell membrane); (ii) interaction of a Dkk protein with an intracellular protein *via* a membrane-bound Dkk receptor; (iii) complex formation
- 10 between a soluble Dkk protein and a second soluble Dkk binding partner (*e.g.*, a non-Dkk protein molecule or a second Dkk protein molecule); (iv) interaction with other extracellular proteins (*e.g.*, regulation of *wnt*-dependent cellular adhesion to extracellular matrix components); (v) binding to and eliminating an undesirable molecule (*e.g.*, a detoxifying activity or defense function); and/or (vi) an enzymatic activity. In yet
- 15 another preferred embodiment, a Dkk activity is at least one or more of the following activities: (1) modulation of cellular signal transduction, either *in vitro* or *in vivo* (*e.g.*, modulation, *e.g.*, antagonism, of the activity of members of the *wnt* family of secreted proteins or suppression of *wnt*-dependent signal transduction, for example suppression of Wnt 2b, Wnt3 and/or Wnt8-dependent signal transduction by hDkk-1 and/or hDkk-4);
- 20 (2) regulation of communication between cells (*e.g.*, regulation of *wnt*-dependent cell-cell interactions); (3) regulation of expression of genes whose expression is modulated by binding of Dkk (*e.g.*, hDkk-3) to a receptor; (4) regulation of gene transcription in a cell involved in development or differentiation, either *in vitro* or *in vivo* (*e.g.*, induction of cellular differentiation); (5) regulation of gene transcription in a cell involved in
- 25 development or differentiation, wherein at least one gene encodes a differentiation-specific protein; (6) regulation of gene transcription in a cell involved in development or differentiation, wherein at least one gene encodes a second secreted protein; (7) regulation of gene transcription in a cell involved in development or differentiation, wherein at least one gene encodes a signal transduction molecule; (8) regulation of
- 30 cellular proliferation, either *in vitro* or *in vivo* (*e.g.*, induction of cellular proliferation or inhibition of proliferation as in the case of suppression of tumorigenesis (*e.g.*, suppression

of glial cell tumor growth, for example, glioblastoma growth)); (9) formation and maintenance of ordered spatial arrangements of differentiated tissues in vertebrates, both adult and embryonic (*e.g.*, induction of head formation during vertebrate development or maintenance of hematopoietic progenitor cells); (10) modulation of cell death, such as stimulation of cell survival; (11) regulating cell migration; and/or (12) immune modulation.

As referred to herein, "differentiation-specific proteins" include proteins involved in the transition of a cell from the undifferentiated to the differentiated phenotype. For example, such proteins can be differentiation specific structural proteins or differentiation-specific transcription factors. Such differentiation-specific proteins are generally expressed at higher levels in cells which are making the transition from the undifferentiated to the differentiated phenotype (*e.g.*, during embryonic development or during regeneration of mature tissue in the adult animal), or are expressed at higher levels in fully-differentiated or terminally-differentiated cells as compared to their undifferentiated counterparts. Also, as referred to herein, "differentiation-specific genes" include nucleic acid molecules which encode differentiation-specific proteins.

Accordingly, another embodiment of the invention features isolated Dkk proteins and polypeptides having a Dkk activity. Preferred Dkk proteins have at least one cysteine-rich region and a Dkk activity. In another preferred embodiment, the Dkk protein has at least one cysteine-rich region, wherein the cysteine-rich region comprises at least one cysteine-rich domain, and a Dkk activity. In another preferred embodiment, the Dkk protein has at least one cysteine-rich region, wherein the cysteine-rich region comprises at least two cysteine-rich domains, and a Dkk activity. In yet another preferred embodiment, a Dkk protein further comprises a signal sequence. In still another preferred embodiment, a Dkk protein has a cysteine-rich region, a Dkk activity, and an amino acid sequence sufficiently homologous to an amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:21.

A preferred Dkk fragment comprises a carboxy-terminal cysteine-rich domain. In one embodiment, a Dkk fragment comprises a carboxy-terminal cysteine-rich domain and retains a biological activity of a Dkk protein. In yet another embodiment, a Dkk fragment lacks an amino-terminal cysteine-rich domain.

The human Dkk-3 cDNA, which is approximately 2479 nucleotides in length, encodes a protein which is approximately 350 amino acid residues in length. The human Dkk-3 protein contains an N-terminal signal sequence and a cysteine-rich region comprising two cysteine-rich domains. A Dkk cysteine-rich region can be found at least, for example, from about amino acids 147-284 of SEQ ID NO:2. The hDkk-3 cysteine-rich region comprises an amino-terminal cysteine-rich domain from about amino acids 147-195 of SEQ ID NO:2 and a carboxy-terminal cysteine-rich domain from about amino acids 201-284 of SEQ ID NO:2. The human Dkk-3 protein is a secreted protein which further contains a signal sequence at about amino acids 1-21, 1-22, 1-23, or 1-24 of SEQ ID NO:2. Accordingly, a mature human Dkk-3 protein begins at about amino acid residue 22, 23, 24, or 25 of SEQ ID NO:2 and is about 329, 328, 327, or 326 amino acids in length. The prediction of such a signal peptide can be made, for example, utilizing the computer algorithm SIGNALP (Nielsen, *et al.*, (1997) *Protein Engineering* 10:1-6).

The human Dkk-4 cDNA, which is approximately 848 nucleotides in length, encodes a protein which is approximately 224 amino acid residues in length. The human Dkk-4 protein contains an N-terminal signal sequence and a cysteine-rich region comprising two cysteine-rich domains. A Dkk cysteine-rich region can be found at least, for example, from about amino acids 41-218 of SEQ ID NO:5. The hDkk-4 cysteine-rich region comprises an amino-terminal cysteine-rich domain from about amino acids 41-90 of SEQ ID NO:5 and a carboxy-terminal cysteine-rich domain from about amino acids 138-218 of SEQ ID NO:5. The human Dkk-4 protein is a secreted protein which further contains a signal sequence at about amino acids 1-17, 1-18, 1-19, or 1-20 of SEQ ID NO:5. Accordingly, a mature human Dkk-4 protein begins at about amino acid residue 18, 19, 20, or 21 of SEQ ID NO:5 and is about 207, 206, 205, or 204 amino acids in length. A preferred fragment of hDkk-4 comprises amino acid residues 134-224 of SEQ ID NO:5. In another embodiment, a preferred fragment of hDkk-4 consists of amino acid residues 134-224 of SEQ ID NO:5.

The human Dkk-1 cDNA, which is approximately 1536 nucleotides in length, encodes a protein which is approximately 266 amino acid residues in length. The human Dkk-1 protein contains an N-terminal signal sequence and a cysteine-rich region

comprising two cysteine-rich domains. A Dkk cysteine-rich region can be found at least, for example, from about amino acids 85-263 of SEQ ID NO:8. The hDkk-1 cysteine-rich region comprises an amino-terminal cysteine-rich domain from about amino acids 85-138 of SEQ ID NO:8 and a carboxy-terminal cysteine-rich domain from about amino acids 182-263 of SEQ ID NO:8. The human Dkk-1 protein is a secreted protein which further contains a signal sequence at about amino acids 1-18, 1-19, 1-20, or 1-21 of SEQ ID NO:8. Accordingly, a mature human Dkk-1 protein begins at about amino acid residue 19, 20, 21, or 22 of SEQ ID NO:8 and is about 248, 247, 246, or 245 amino acids in length.

10 The human Dkk-2 cDNA, which is approximately 3687 nucleotides in length, encodes a protein which is approximately 259 amino acid residues in length. The human Dkk-2 protein contains a cysteine-rich region comprising two cysteine-rich domains. A Dkk cysteine-rich region can be found at least, for example, from about amino acids 78-256 of SEQ ID NO:21. The hDkk-2 cysteine-rich region comprises an amino-terminal cysteine-rich domain from about amino acids 78-127 of SEQ ID NO:21 and a carboxy-terminal cysteine-rich domain from about amino acids 176-256 of SEQ ID NO:21. The human Dkk-2 protein is a secreted protein which further contains a signal sequence at about amino acids 1-31, 1-32, 1-33, or 1-34 of SEQ ID NO:21. Accordingly, a mature human Dkk-2 protein begins at about amino acid residue 32, 33, 34, or 35 of SEQ ID NO:21 and is about 228, 227, 226, or 225 amino acids in length.

Dkk proteins of the present invention can be used to identify additional Dkk-related proteins or family members. For example, a protein having homology to hDkk-3 was identified using the nucleotide sequence encoding the N-terminal unique region of hDkk-3 to search a nucleotide sequence database. A human cDNA clone (Accession No.: AA397836) was identified from the dBEST database as having homology to hDkk-3 and was fully sequenced. The encoded protein is referred to herein as human "Soggy-1" or "Dkk-like-N". The nucleotide and predicted amino acid sequence of human Soggy-1 are depicted in Figure 7. The nucleotide sequence of human Soggy-1 (SEQ ID NO:13) encodes a protein having 242 amino acids (SEQ ID NO:14). The nucleotide sequence of human Soggy-1 includes a 5' untranslated region containing nucleotides 1-74 of SEQ ID NO:13, a coding region containing nucleotides 75-800 of SEQ ID NO:13

(corresponding to nucleotides 1-726 of SEQ ID NO:15), and a 3' untranslated region containing nucleotides 801-928 of SEQ ID NO:13. The Soggy-1 protein (amino acid residues 32-132) has 25% identity to an N-terminal domain of human Dkk-3 (consisting of amino acid residues 22-140) as determined by ALIGN, Myers and Miller, (1989)

5 *CABIOS*, utilizing a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4.

Two murine cDNA clones were further identified from the database and fully sequenced. Combining the sequence information from these two clones resulted in a full-length sequence for murine Soggy-1. The nucleotide sequence and predicted amino

10 acid sequence of murine Soggy-1 are depicted in Figure 8. The nucleotide sequence of murine Soggy-1 (SEQ ID NO:26) encodes a protein having 230 amino acids (SEQ ID NO:27). The nucleotide sequence of murine Soggy-1 includes a 5' untranslated region containing nucleotides 1-56 of SEQ ID NO:26, a coding region containing nucleotides 57-746 of SEQ ID NO:26 (corresponding to SEQ ID NO:26), and a 3' untranslated

15 region containing nucleotides 747-835 of SEQ ID NO:26. Human and murine Soggy-1 proteins display 59% overall identity. An alignment of human and murine Soggy proteins to human and murine Dkk-3 proteins is depicted in Figure 10.

In one embodiment, a Soggy protein is identified based on the presence of at least one soggy domain or "SGY" domain in the protein or corresponding nucleic acid

20 molecule. As defined herein, a "SGY domain" includes a protein domain of a Soggy protein (*e.g.*, hSoggy-1) having an amino acid sequence of about 45-56 amino acids and having at least about 25-40% identity with amino acid residues 90-140 of hDkk-3 (leu90-glu140 of SEQ ID NO:2). In another embodiment, a "SGY domain" has 46-55, preferably 47-54, more preferably about 48-53, and even more preferably about 49-52 or

25 50-51 amino acids, and has at least about 27-38%, preferably about 28-37%, more preferably about 29-36%, even more preferably about 30-35%, and even more preferably about 31-34%, or 32-33% identity with amino acid residues 90-140 of hDkk-3 (Leu90-Glu140 of SEQ ID NO:2). In yet another embodiment, a "SGY domain" has the following consensus sequence: L - P - X(3) - H - X - E - X(7) - G - N - X - T - X(3) - H -

30 X(4) - K - X - T - X - N - X(2) - G - X(4) - S - E - X - V - X(2) - S - X(4) - E (SEQ ID NO:29). For example, human Soggy-1 has a SGY domain from about amino acid

residues 81-131 (Leu 81-Glu131 of SEQ ID NO:14) having 33% identity with amino acid residues 90-140 of hDkk-3 (leu90-glu140 of SEQ ID NO:2). Likewise, murine Soggy-1 has a SGY domain from about amino acid residues 71-120 (Leu 71-Glu120 of SEQ ID NO:27) having 33% identity with amino acid residues 90-140 of hDkk-3
5 (leu90-glu140 of SEQ ID NO:2). The SGY domains of human and murine Soggy-1 are depicted by shaded boxes in Figure 10.

In another embodiment of the invention, a Soggy protein has at least one SGY domain and a signal sequence. For example, in one embodiment, a hSoggy-1 protein contains a signal sequence of about amino acids 1-29, 1-30, 1-31, or 1-32 of SEQ ID
10 NO:14. Accordingly, a mature hSoggy-1 protein begins at about amino acid residue 30, 31, 32, or 33 of SEQ ID NO:14 and is about 213, 212, 211, or 210 amino acids in length. In another embodiment, a mSoggy-1 protein contains a signal sequence of about amino acids 1-19, 1-20, 1-21, or 1-22 of SEQ ID NO:27. Accordingly, a mature mSoggy-1 protein begins at about amino acid residue 211, 210, 209, or 208 of SEQ ID NO:28 and
15 is about 213, 212, 211, or 210 amino acids in length.

Various aspects of the invention are described in further detail in the following subsections:

I. Isolated Nucleic Acid Molecules

20 One aspect of the invention pertains to isolated nucleic acid molecules that encode Dkk proteins or biologically active portions thereof, as well as nucleic acid fragments sufficient for use as hybridization probes to identify Dkk-encoding nucleic acids (*e.g.*, Dkk mRNA) and fragments for use as PCR primers for the amplification or mutation of Dkk nucleic acid molecules. As used herein, the term "nucleic acid
25 molecule" is intended to include DNA molecules (*e.g.*, cDNA or genomic DNA) and RNA molecules (*e.g.*, mRNA) and analogs of the DNA or RNA generated using nucleotide analogs. The nucleic acid molecule can be single-stranded or double-stranded, but preferably is double-stranded DNA.

An "isolated" nucleic acid molecule is one which is separated from other nucleic
30 acid molecules which are present in the natural source of the nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (*i.e.*,

sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated Dkk nucleic acid molecule can contain less than about 5 kb, 4kb, 3kb, 2kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived. An isolated chromosome is not an isolated nucleic acid molecule as defined herein. Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized.

A nucleic acid molecule of the present invention, *e.g.*, a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a portion thereof, can be isolated using standard molecular biology techniques and the sequence information provided herein. Using all or portion of the nucleic acid sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, as a hybridization probe, Dkk nucleic acid molecules can be isolated using standard hybridization and cloning techniques (*e.g.*, as described in Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989).

Moreover, a nucleic acid molecule encompassing all or a portion of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as

Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, can be isolated by the polymerase chain reaction (PCR) using synthetic oligonucleotide primers
5 designed based upon the sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as
10 Accession Number _____.

A nucleic acid of the invention can be amplified using cDNA, mRNA or alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis.
15 Furthermore, oligonucleotides corresponding to Dkk nucleotide sequences can be prepared by standard synthetic techniques, *e.g.*, using an automated DNA synthesizer.

In a preferred embodiment, an isolated nucleic acid molecule of the invention comprises the nucleotide sequence shown in SEQ ID NO:1. The sequence of SEQ ID NO:1 corresponds to the human Dkk-3 cDNA. This cDNA comprises sequences
20 encoding the human Dkk-3 protein (*i.e.*, "the coding region", from nucleotides 38-1087), as well as 5' untranslated sequences (nucleotides 1 to 37) and 3' untranslated sequences (nucleotides 1088 to 2479). Alternatively, the nucleic acid molecule can comprise only the coding region of SEQ ID NO:1 (*e.g.*, nucleotides 38 to 1087, corresponding to SEQ ID NO:3). A plasmid containing the full-length nucleotide sequence encoding hDkk-3
25 was deposited with the American Type Culture Collection (ATCC), presently in Manassas Virginia, on June 11, 1997 and assigned Accession Number 98452.

In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises the nucleotide sequence shown in SEQ ID NO:4. The sequence of SEQ ID NO:4 corresponds to the human Dkk-4 cDNA. This cDNA comprises
30 sequences encoding the human Dkk-4 protein (*i.e.*, "the coding region", from nucleotides 125-796), as well as 5' untranslated sequences (nucleotides 1 to 124) and 3'

untranslated sequences (nucleotides 797 to 848). Alternatively, the nucleic acid molecule can comprise only the coding region of SEQ ID NO:4 (*e.g.*, nucleotides 125 to 796, corresponding to SEQ ID NO:6).

5 In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises the nucleotide sequence shown in SEQ ID NO:7. The sequence of SEQ ID NO:7 corresponds to the human Dkk-1 cDNA. This cDNA comprises sequences encoding the human Dkk-1 protein (*i.e.*, "the coding region", from nucleotides 109-906), as well as 5' untranslated sequences (nucleotides 1 to 108) and 3' untranslated sequences (nucleotides 907-1536). Alternatively, the nucleic acid molecule
10 can comprise only the coding region of SEQ ID NO:7 (*e.g.*, nucleotides 109-906, corresponding to SEQ ID NO:9). A plasmid containing the full-length nucleotide sequence encoding hDkk-1 was deposited with the American Type Culture Collection (ATCC), presently in Manassas Virginia, on January 16, 1998 and assigned Accession Number 98633.

15 In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises the nucleotide sequence shown in SEQ ID NO:20. The sequence of SEQ ID NO:20 corresponds to the human Dkk-2 cDNA. This cDNA comprises sequences encoding the human Dkk-2 protein (*i.e.*, "the coding region", from nucleotides 724-1500), 5' untranslated sequences (nucleotides 1-723), as well as 3' untranslated sequences (nucleotides 1501-3687). Alternatively, the nucleic acid
20 molecule can comprise only the coding region of SEQ ID NO:20 (*e.g.*, nucleotides 724 to 1500, corresponding to SEQ ID NO:22). A plasmid, clone fthul33, containing the full-length nucleotide sequence encoding hDkk-2 was deposited with the American Type Culture Collection (ATCC), presently in Manassas Virginia, on March 2, 1999 and
25 assigned Accession Number _____.

In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises the nucleotide sequence shown in SEQ ID NO:13. The sequence of SEQ ID NO:13 corresponds to the human Soggy cDNA. This cDNA comprises sequences encoding the human Soggy protein (*i.e.*, "the coding region", from
30 nucleotides 75 to 800), as well as 5' untranslated sequences (nucleotides 1 to 74) and 3' untranslated sequences (nucleotides 801 to 928). Alternatively, the nucleic acid

molecule can comprise only the coding region of SEQ ID NO:13 (*e.g.*, nucleotides 75 to 800, corresponding to SEQ ID NO:15).

In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises a nucleic acid molecule which is a complement of the nucleotide
5 sequence shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide
10 sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a portion of any of these nucleotide sequences. A nucleic acid molecule which is complementary to the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as
15 Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, is one which is sufficiently complementary to the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide
20 sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, such that it can hybridize to the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20, or the nucleotide sequence of the DNA insert of the plasmid deposited
25 with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, thereby forming a stable duplex.

In still another preferred embodiment, an isolated nucleic acid molecule of the
30 present invention comprises a nucleotide sequence which is at least about 30-35%, preferably about 40-45%, more preferably about 50-55%, even more preferably about

60-65%, and even more preferably at least about 70-75%, 80-85%, 90-95% or more homologous to the nucleotide sequences shown in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:20, or SEQ ID NO:22, the nucleotide sequence of the DNA insert of the
5 plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a portion of any of these nucleotide sequences.

In one aspect, the present invention features isolated nucleic acid molecules
10 which are linear (*e.g.*, linear fragments of double-stranded DNA, linear strands of single-stranded DNA, single-stranded RNA molecules, and oligonucleotides). Another aspect of the present invention features circular nucleic acid molecules (*e.g.*, double-stranded DNA molecules, for example, plasmid molecules including the nucleotide sequences shown in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7,
15 SEQ ID NO:9, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:20, or SEQ ID NO:22, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a
20 portion of any of these nucleotide sequences).

In one embodiment, the isolated nucleic acid molecules of the present invention are DNA molecules which are in a form suitable for expression (*e.g.*, suitable for expression of corresponding messenger RNA or mRNA). In another embodiment, the isolated nucleic acid molecules are DNA molecules which are in a form suitable for
25 expression of corresponding protein (*e.g.*, in a form, for example, in a vector, which is capable of expressing protein, *e.g.*, in the appropriate orientation for expression from regulatory elements and/or in-frame with appropriate regulatory elements). In another embodiment, the isolated nucleic acids are in a form suitable for determination of nucleic acid sequence (*e.g.*, in a form suitable for sequencing, for example, is a
30 sequencing vector including a M13, T7, T3 and SP6 promoter. Examples of sequencing vectors include, but are not limited to pBluescript (Stratagene™), pT7T3D (Pharmacia™)

and pCR2.1 (InVitrogen). In yet another embodiment, the isolated nucleic acid molecules are free from vector sequences. In a preferred embodiment, an isolated nucleic acid molecule is free from sequencing vector sequences.

Moreover, the nucleic acid molecule of the invention can comprise only a portion
5 of the nucleic acid sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID
NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid
deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA
insert of the plasmid deposited with ATCC as Accession Number 98633, or the
nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as
10 Accession Number _____, for example a fragment which can be used as a probe or
primer or a fragment encoding a biologically active portion of a Dkk protein or Dkk-
related protein. The nucleotide sequence determined from the cloning of the human Dkk
genes allows for the generation of probes and primers designed for use in identifying
and/or cloning Dkk homologues in other cell types, *e.g.*, from other tissues, as well as
15 Dkk homologues from other mammals and Dkk-related proteins. The probe/primer
typically comprises substantially purified oligonucleotide. The oligonucleotide typically
comprises a region of nucleotide sequence that hybridizes under stringent conditions to
at least about 12, preferably about 25, more preferably about 40, 50 or 75 consecutive
nucleotides of a sense sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ
20 ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid
deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA
insert of the plasmid deposited with ATCC as Accession Number 98633, or the
nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as
Accession Number _____, of an anti-sense sequence of SEQ ID NO:1, SEQ ID NO:4,
25 SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA
insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide
sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number
98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with
ATCC as Accession Number _____, or of a naturally occurring mutant of SEQ ID
30 NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide
sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number

98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____. In an exemplary embodiment, a nucleic acid molecule of the present invention comprises a nucleotide
5 sequence which hybridizes under stringent hybridization conditions to a nucleic acid molecule consisting of nucleotides 470-2479 of SEQ ID NO:1 or to a nucleic acid molecule consisting of nucleotides 1-475 of SEQ ID NO:4.

Probes based on human nucleotide sequences (*e.g.*, the human Dkk nucleotide sequence) can be used to detect transcripts or genomic sequences encoding the same or
10 homologous proteins. For instance, primers based on the nucleic acid represented in SEQ ID NOs:1 or 3 can be used in PCR reactions to clone Dkk homologs (*e.g.*, hDkk-3 homologues). In a preferred embodiment of the invention, Dkk homologs are cloned by PCR amplification (*e.g.*, RT-PCR) using primers hybridizing to a portion of the nucleotide sequence encoding the Dkk cysteine rich domain. Likewise, probes based on
15 the subject Dkk sequences can be used to detect transcripts or genomic sequences encoding the same or homologous proteins. In preferred embodiments, the probe further comprises a label group attached thereto, *e.g.*, the label group can be a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used as a part of a diagnostic test kit for identifying cells or tissue which misexpress a Dkk
20 protein, such as by measuring a level of a Dkk-encoding nucleic acid in a sample of cells from a subject *e.g.*, detecting Dkk mRNA levels or determining whether a genomic Dkk gene has been mutated or deleted.

A nucleic acid fragment encoding a "biologically active portion of a Dkk or Dkk-related protein" can be prepared by isolating a portion of SEQ ID NO:1, SEQ ID NO:4,
25 SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, which encodes a polypeptide having a biological
30 activity (the biological activities of the Dkk and Dkk-related proteins have previously

been described), expressing the encoded portion of the protein (*e.g.*, by recombinant expression *in vitro*) and assessing the activity of the encoded portion of the protein.

The invention further encompasses nucleic acid molecules that differ from the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, due to degeneracy of the genetic code and thus encode the same proteins as those encoded by the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____. In another embodiment, an isolated nucleic acid molecule of the invention has a nucleotide sequence encoding a protein having an amino acid sequence shown in SEQ ID NO:2, SEQ ID NO: 5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21.

In addition to the human nucleotide sequences shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, it will be appreciated by those skilled in the art that DNA sequence polymorphisms that lead to changes in the amino acid sequences of the Dkk or Dkk-related proteins may exist within a population (*e.g.*, the human population). Such genetic polymorphism in the Dkk or Dkk-related genes may exist among individuals within a population due to natural allelic variation. As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules comprising an open reading frame encoding a protein, preferably a mammalian Dkk or Dkk-related protein. Such natural allelic variations can typically result in 1-5% variance

in the nucleotide sequence of a Dkk or Dkk-related gene. Any and all such nucleotide variations and resulting amino acid polymorphisms in genes that are the result of natural allelic variation and that do not alter the functional activity of a Dkk or Dkk-related protein are intended to be within the scope of the invention.

5 Moreover, nucleic acid molecules encoding Dkk or Dkk-related proteins from other species, and thus which have a nucleotide sequence which differs from the human sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid
10 deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, are intended to be within the scope of the invention. For example, a murine Dkk-3 cDNA has been identified based of the nucleotide sequence of human Dkk-3. The nucleotide sequence of murine Dkk-3 (SEQ ID NO:16) encodes a hDkk-3 protein having 349
15 amino acids. The nucleotide and amino acid sequences of murine Dkk-3 are depicted in Figure 5. The coding region of murine Dkk-3 is represented by SEQ ID NO:18. A plasmid containing the full-length nucleotide sequence encoding mDkk-3 was deposited with the American Type Culture Collection (ATCC), presently in Manassas, Virginia, on January 16, 1998 and assigned Accession Number 98634. Likewise, a murine Dkk-
20 related protein (Soggy-1) has been identified based of the nucleotide sequence of human Dkk-3. The nucleotide sequence of murine Soggy-1 (SEQ ID NO:26) encodes a protein having 230 amino acids (SEQ ID NO:27). The nucleotide and amino acid sequences of murine Soggy-1 are depicted in Figure 8. The coding region of murine Soggy-1 is represented by SEQ ID NO:28.

25 Nucleic acid molecules corresponding to natural allelic variants and homologues of the Dkk or Dkk-related cDNAs of the invention can be isolated based on their homology to the human nucleic acids disclosed herein using the human cDNA, or a portion thereof, as a hybridization probe according to standard hybridization techniques under stringent hybridization conditions. Examples of tissues and/or libraries suitable
30 for isolation of the subject nucleic acids include brain, spinal chord and heart tissue. cDNA encoding a Dkk protein (*e.g.*, a hDkk-3 protein) can be obtained by isolating total

mRNA from a cell, *e.g.*, a vertebrate cell, a mammalian cell, or a human cell, including embryonic cells. Double stranded cDNAs can then be prepared from the total mRNA, and subsequently inserted into a suitable plasmid or bacteriophage vector using any one of a number of known techniques. The gene encoding a hDkk-3 protein can also be
5 cloned using established polymerase chain reaction techniques in accordance with the nucleotide sequence information provided by the invention. The nucleic acid of the invention can be DNA or RNA or analogs thereof.

Accordingly, in another embodiment, an isolated nucleic acid molecule of the invention is at least 15 nucleotides in length and hybridizes under stringent conditions to
10 the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid
15 deposited with ATCC as Accession Number _____. In another embodiment, the nucleic acid is at least 30, 50, 100, 250, 300, 400 or 500 nucleotides in length. As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences at least 60% homologous to each other typically remain hybridized to each other. Preferably, the conditions are
20 such that sequences at least about 70%, more preferably at least about 80%, even more preferably at least about 85% or 90% homologous to each other typically remain hybridized to each other. Such stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. A preferred, non-limiting example of stringent hybridization
25 conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65°C. Preferably, an isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to the sequence of SEQ ID NO:1 corresponds to a naturally-occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an
30 RNA or DNA molecule having a nucleotide sequence that occurs in nature (*e.g.*, encodes a natural protein).

In addition to naturally-occurring allelic variants of the Dkk or Dkk-related sequences that may exist in the population, the skilled artisan will further appreciate that changes can be introduced by mutation into the nucleotide sequences of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, thereby leading to changes in the amino acid sequence of the encoded Dkk proteins, without altering the functional ability of the Dkk proteins. For example, nucleotide substitutions leading to amino acid substitutions (particularly conservative amino acid substitutions) at "non-essential" amino acid residues can be made in the sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____. A "non-essential" amino acid residue is a residue that can be altered from the wild-type sequence of Dkk (or wild-type Dkk-related sequence) (*e.g.*, the sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21) without altering the biological activity, whereas an "essential" amino acid residue is required for biological activity. For example, amino acid residues that are conserved among the Dkk or Dkk-related proteins of the present invention (*e.g.*, cysteine residues within cysteine-rich domains), are predicted to be particularly unamenable to alteration. Furthermore, amino acid residues that are conserved between Dkk protein and other proteins having cysteine-rich domains are not likely to be amenable to alteration.

Accordingly, another aspect of the invention pertains to nucleic acid molecules encoding Dkk or Dkk-related proteins that contain changes in amino acid residues that are not essential for activity. Such proteins differ in amino acid sequence from SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21 yet retain biological activity. In one embodiment, the isolated nucleic acid molecule comprises a

nucleotide sequence encoding a protein, wherein the protein comprises an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8,, SEQ ID NO:14, or SEQ ID NO:21. Preferably, the protein encoded by the nucleic acid molecule is at least about 65-70% homologous to

5 SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21, more preferably at least about 75-80% homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21, even more preferably at least about 85-90% homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21, and most preferably at least about 95% homologous to SEQ ID NO:2, SEQ

10 ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21.

An isolated nucleic acid molecule encoding a Dkk protein homologous to the protein of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21 can be created by introducing one or more nucleotide substitutions, additions or deletions into the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7,

15 SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, such that one or more amino acid substitutions, additions or

20 deletions are introduced into the encoded protein. Mutations can be introduced into SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the

25 DNA insert of the plasmid deposited with ATCC as Accession Number _____, by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Preferably, conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar

30 side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (*e.g.*,

lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*,
5 threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine). Thus, for example, a predicted nonessential amino acid residue in a Dkk protein (*e.g.*, one not located in a cysteine-rich domain) is preferably replaced with another amino acid residue from the same side chain family. Alternatively, in another embodiment, mutations can be introduced randomly along all or part of a Dkk or
10 Dkk-related coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for biological activity to identify mutants that retain activity. Following mutagenesis of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of
15 the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, the encoded protein can be expressed recombinantly and the activity of the protein can be determined.

In a preferred embodiment, a mutant Dkk or Dkk-related protein can be assayed
20 for intracellular calcium, an increase in phosphatidylinositol or other molecule, and can result, *e.g.*, in phosphorylation of specific proteins, a modulation of gene transcription and any of the other biological activities set forth herein.

In a preferred embodiment, a mutant Dkk or Dkk-related protein can also be assayed for the ability to (1) modulate cellular signal transduction, either *in vitro* or *in*
25 *vivo*; (2) regulate communication between cells; (3) regulate expression of genes whose expression is modulated by binding of Dkk (*e.g.*, hDkk-3) to a receptor; (4) regulate gene transcription in a cell involved in development or differentiation, either *in vitro* or *in vivo*; (5) regulate cellular proliferation, either *in vitro* or *in vivo*; (6) form and/or maintain ordered spatial arrangements of differentiated tissues in vertebrates; (7)
30 modulate cell death (*e.g.* cell survival); (8) regulate cell migration; and/or (9) modulate immune system function.

In addition to the nucleic acid molecules encoding Dkk or Dkk-related proteins described above, another aspect of the invention pertains to isolated nucleic acid molecules which are antisense thereto. An "antisense" nucleic acid comprises a nucleotide sequence which is complementary to a "sense" nucleic acid encoding a protein, *e.g.*, complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence. Accordingly, an antisense nucleic acid can hydrogen bond to a sense nucleic acid. The antisense nucleic acid can be complementary to an entire Dkk coding strand, or to only a portion thereof. In one embodiment, an antisense nucleic acid molecule is antisense to a "coding region" of the coding strand of a nucleotide sequence encoding Dkk. The term "coding region" refers to the region of the nucleotide sequence comprising codons which are translated into amino acid residues (*e.g.*, the coding region of human Dkk-3 corresponds to SEQ ID NO:3, the coding region of human Dkk-4 corresponds to SEQ ID NO:6, the coding region of human Dkk-1 corresponds to SEQ ID NO:9, the coding region of human Dkk-2 corresponds to SEQ ID NO:22, and the coding region of human Soggy corresponds to SEQ ID NO:15). In another embodiment, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence encoding a Dkk or Dkk-related protein. The term "noncoding region" refers to 5' and 3' sequences which flank the coding region that are not translated into amino acids (*i.e.*, also referred to as 5' and 3' untranslated regions).

Given the coding strand sequences disclosed herein (*e.g.*, SEQ ID NO:3, SEQ ID NO:6, SEQ ID NO:9, SEQ ID NO:15, or SEQ ID NO:22), antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick base pairing. The antisense nucleic acid molecule can be complementary to the entire coding region of a Dkk or Dkk-related mRNA, but more preferably is an oligonucleotide which is antisense to only a portion of the coding or noncoding region of the mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of Dkk mRNA. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, or 90 nucleotides in length. In a preferred embodiment, an oligonucleotide is about 30-90, preferably about 40-80, more preferably about 50-70 nucleotides in length and is antisense to a portion of

SEQ ID NO:1 from about nucleotides 1-150. In another embodiment, an oligonucleotide is antisense to a portion of SEQ ID NO:4 from about nucleotides 25-225. In another embodiment, an oligonucleotide is antisense to a portion of SEQ ID NO:7 from about nucleotides 1-200. In another embodiment, an oligonucleotide is antisense to a portion of SEQ ID NO:20 from about nucleotides 625-825. In yet another embodiment, an oligonucleotide is antisense to a portion of SEQ ID NO:13 from about nucleotides 1-175.

An antisense nucleic acid of the invention can be constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (*e.g.*, an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, *e.g.*, phosphorothioate derivatives, acridine substituted nucleotides, can be used. Alternatively, the antisense nucleic acid molecule can be synthesized to increase transport across cellular membranes, *e.g.*, methylphosphonate derivatives. The antisense molecules can include a 3'-terminal cap (*e.g.*, a 3'-aminopropyl modification), a biotin moiety, or even a 3'-3' terminal linkage.

Examples of modified nucleotides which can be used to generate the antisense nucleic acid include 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (*v*), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (*v*), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)*w*, and 2,6-diaminopurine. Alternatively, the

antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (*i.e.*, RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

- 5 The antisense nucleic acid molecules of the invention are typically administered to a subject or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a Dkk or Dkk-related protein to thereby inhibit expression of the protein, *e.g.*, by inhibiting transcription and/or translation. The hybridization can be by conventional nucleotide complementarity to form a stable
- 10 duplex, or, for example, in the case of an antisense nucleic acid molecule which binds to DNA duplexes, through specific interactions in the major groove of the double helix. An example of a route of administration of antisense nucleic acid molecules of the invention include direct injection at a tissue site. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then administered systemically.
- 15 For example, for systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface, *e.g.*, by linking the antisense nucleic acid molecules to peptides or antibodies which bind to cell surface receptors or antigens. The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular
- 20 concentrations of the antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

- In yet another embodiment, the antisense nucleic acid molecule of the invention is an α -anomeric nucleic acid molecule. An α -anomeric nucleic acid molecule forms
- 25 specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gaultier *et al.*, (1987) *Nucleic Acids Res.* 15:6625-6641). The antisense nucleic acid molecule can also comprise a 2'-*o*-methylribonucleotide (Inoue *et al.*, (1987) *Nucleic Acids Res.* 15:6131-6148) or a chimeric RNA-DNA analogue (Inoue *et al.*, (1987) *FEBS Lett.* 215:327-330).

In still another embodiment, an antisense nucleic acid of the invention is a ribozyme. Ribozymes are catalytic RNA molecules with ribonuclease activity which are capable of cleaving a single-stranded nucleic acid, such as an mRNA, to which they have a complementary region. Thus, ribozymes (*e.g.*, hammerhead ribozymes
5 (described in Haselhoff and Gerlach (1988) *Nature* 334:585-591)) can be used to catalytically cleave Dkk or Dkk-related mRNA transcripts to thereby inhibit translation of Dkk or Dkk-related mRNA. A ribozyme having specificity for a Dkk- or Dkk-related-encoding nucleic acid can be designed based upon the nucleotide sequence of a Dkk or Dkk-related cDNA disclosed herein (*i.e.*, SEQ ID NO:1, SEQ ID NO:4, SEQ ID
10 NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633). For example, a derivative of a *Tetrahymena* L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be
15 cleaved in a Dkk-encoding mRNA. See, *e.g.*, Cech *et al.*, U.S. Patent No. 4,987,071; and Cech *et al.*, U.S. Patent No. 5,116,742. Alternatively, Dkk (or Dkk-related) mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See, *e.g.*, Bartel, D. and Szostak, J.W. (1993) *Science* 261:1411-1418.

20 Alternatively, gene expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region of the Dkk or Dkk-related gene (*e.g.*, the promoter and/or enhancers) to form triple helical structures that prevent transcription of the gene in target cells. See generally, Helene, C. (1991) *Anticancer Drug Des.* 6(6):569-84; Helene, C. *et al.*, (1992) *Ann. N.Y. Acad. Sci.* 660:27-36; and Maher, L.J.
25 (1992) *Bioassays* 14(12):807-15.

In yet another embodiment, the nucleic acid molecules of the present invention can be modified at the base moiety, sugar moiety or phosphate backbone to improve, *e.g.*, the stability, hybridization, or solubility of the molecule. For example, the deoxyribose phosphate backbone of the nucleic acid molecules can be modified to
30 generate peptide nucleic acids (see Hyrup B. *et al.* (1996) *Bioorganic & Medicinal Chemistry* 4 (1): 5-23). As used herein, the terms "peptide nucleic acids" or "PNAs"

refer to nucleic acid mimics, *e.g.*, DNA mimics, in which the deoxyribose phosphate backbone is replaced by a pseudopeptide backbone and only the four natural nucleobases are retained. The neutral backbone of PNAs has been shown to allow for specific hybridization to DNA and RNA under conditions of low ionic strength. The synthesis
5 of PNA oligomers can be performed using standard solid phase peptide synthesis protocols as described in Hyrup B. *et al.* (1996) *supra*; Perry-O'Keefe *et al.* PNAS 93: 14670-675.

PNAs of Dkk or Dkk-related nucleic acid molecules can be used therapeutic and diagnostic applications. For example, PNAs can be used as antisense or antigene agents
10 for sequence-specific modulation of gene expression by, for example, inducing transcription or translation arrest or inhibiting replication. PNAs of Dkk or Dkk-related nucleic acid molecules can also be used in the analysis of single base pair mutations in a gene, (*e.g.*, by PNA-directed PCR clamping); as 'artificial restriction enzymes' when used in combination with other enzymes, (*e.g.*, S1 nucleases (Hyrup B. (1996) *supra*));
15 or as probes or primers for DNA sequencing or hybridization (Hyrup B. *et al.* (1996) *supra*; Perry-O'Keefe *supra*).

In another embodiment, PNAs of Dkk can be modified, (*e.g.*, to enhance their stability or cellular uptake), by attaching lipophilic or other helper groups to PNA, by the formation of PNA-DNA chimeras, or by the use of liposomes or other techniques of
20 drug delivery known in the art. For example, PNA-DNA chimeras of Dkk nucleic acid molecules can be generated which may combine the advantageous properties of PNA and DNA. Such chimeras allow DNA recognition enzymes, (*e.g.*, RNase H and DNA polymerases), to interact with the DNA portion while the PNA portion would provide high binding affinity and specificity. PNA-DNA chimeras can be linked using linkers of
25 appropriate lengths selected in terms of base stacking, number of bonds between the nucleobases, and orientation (Hyrup B. (1996) *supra*). The synthesis of PNA-DNA chimeras can be performed as described in Hyrup B. (1996) *supra* and Finn P.J. *et al.* (1996) *Nucleic Acids Res.* 24 (17): 3357-63. For example, a DNA chain can be synthesized on a solid support using standard phosphoramidite coupling chemistry and
30 modified nucleoside analogs, *e.g.*, 5'-(4-methoxytrityl)amino-5'-deoxy-thymidine phosphoramidite, can be used as a between the PNA and the 5' end of DNA (Mag, M. *et*

al. (1989) *Nucleic Acid Res.* 17: 5973-88). PNA monomers are then coupled in a stepwise manner to produce a chimeric molecule with a 5' PNA segment and a 3' DNA segment (Finn P.J. *et al.* (1996) *supra*). Alternatively, chimeric molecules can be synthesized with a 5' DNA segment and a 3' PNA segment (Peterser, K.H. *et al.* (1975) 5 *Bioorganic Med. Chem. Lett.* 5: 1119-11124).

In other embodiments, the oligonucleotide may include other appended groups such as peptides (*e.g.*, for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (see, *e.g.*, Letsinger *et al.* (1989) *Proc. Natl. Acad. Sci. US.* 86:6553-6556; Lemaitre *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:648-652; 10 PCT Publication No. W088/09810, published December 15, 1988) or the blood-brain barrier (see, *e.g.*, PCT Publication No. W089/10134, published April 25, 1988). In addition, oligonucleotides can be modified with hybridization-triggered cleavage agents (See, *e.g.*, Krol *et al.* (1988) *BioTechniques* 6:958-976) or intercalating agents. (See, *e.g.*, Zon (1988) *Pharm. Res.* 5:539-549). To this end, the oligonucleotide may be 15 conjugated to another molecule, (*e.g.*, a peptide, hybridization triggered cross-linking agent, transport agent, or hybridization-triggered cleavage agent).

II. Isolated Dkk Proteins and Anti-Dkk Antibodies

One aspect of the invention pertains to isolated Dkk proteins, Dkk-related 20 proteins and biologically active portions thereof, as well as polypeptide fragments suitable for use as immunogens to raise antibodies. In one embodiment, native Dkk or Dkk-related proteins can be isolated from cells or tissue sources by an appropriate purification scheme using standard protein purification techniques. In another embodiment, proteins are produced by recombinant DNA techniques. Alternative to 25 recombinant expression, a Dkk or Dkk-related protein or polypeptide can be synthesized chemically using standard peptide synthesis techniques.

An "isolated" or "purified" protein or biologically active portion thereof is substantially free of cellular material or other contaminating proteins from the cell or tissue source from which the Dkk or Dkk-related protein is derived, or substantially free 30 from chemical precursors or other chemicals when chemically synthesized. The language "substantially free of cellular material" includes preparations of protein in

which the protein is separated from cellular components of the cells from which it is isolated or recombinantly produced. In one embodiment, the language "substantially free of cellular material" includes preparations of Dkk or Dkk-related protein having less than about 30% (by dry weight) of non-Dkk protein or non-Dkk-related protein (also referred to herein as a "contaminating protein"), more preferably less than about 20% of non-Dkk protein or non-Dkk-related protein, still more preferably less than about 10% of non-Dkk protein or non-Dkk-related protein, and most preferably less than about 5% non-Dkk protein or non-Dkk-related protein. When the Dkk or Dkk-related protein or biologically active portion thereof is recombinantly produced, it is also preferably substantially free of culture medium, *i.e.*, culture medium represents less than about 20%, more preferably less than about 10%, and most preferably less than about 5% of the volume of the protein preparation.

The language "substantially free of chemical precursors or other chemicals" includes preparations of Dkk or Dkk-related protein in which the protein is separated from chemical precursors or other chemicals which are involved in the synthesis of the protein. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of Dkk protein having less than about 30% (by dry weight) of chemical precursors, non-Dkk chemicals, or non-Dkk-related chemicals, more preferably less than about 20% chemical precursors, non-Dkk chemicals, or non-Dkk-related chemicals, still more preferably less than about 10% chemical precursors, non-Dkk chemicals, or non-Dkk-related chemicals, and most preferably less than about 5% chemical precursors, non-Dkk chemicals, or non-Dkk-related chemicals.

Biologically active portions of a Dkk or Dkk-related protein include peptides comprising amino acid sequences sufficiently homologous to or derived from the amino acid sequence of the Dkk or Dkk-related protein, *e.g.*, the amino acid sequence shown in SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21, which include less amino acids than the full length proteins, and exhibit at least one activity of a Dkk or Dkk-related protein. Typically, biologically active portions comprise a domain or motif with at least one activity of the Dkk or Dkk-related protein. A biologically active portion of a protein can be a polypeptide which is, for example, 10, 25, 50, 100 or more amino acids in length.

In one embodiment, a biologically active portion of a Dkk protein comprises at least a cysteine-rich region. In another embodiment, a biologically active portion of a Dkk protein comprises at least a cysteine-rich region, wherein the cysteine-rich region includes at least one cysteine-rich domain. In yet another embodiment, a biologically
5 active portion of a Dkk protein comprises at least a signal sequence.

In another embodiment, a biologically active portion of a Dkk-related protein (*e.g.*, a Soggy protein) comprises at least a Soggy domain. In yet another embodiment, a biologically active portion of a Dkk-related protein comprises at least a signal sequence.

In an alternative embodiment, a biologically active portion of a Dkk or Dkk-
10 related protein comprises an amino acid sequence lacking a signal sequence.

It is to be understood that a preferred biologically active portion of a Dkk or Dkk-related protein of the present invention may contain at least one of the above-identified structural domains. A more preferred biologically active portion of a Dkk or Dkk-related protein may contain at least two of the above-identified structural domains.
15 An even more preferred biologically active portion of a protein may contain at least three of the above-identified structural domains. A particularly preferred biologically active portion of a protein of the present invention may contain at least four of the above-identified structural domains.

Moreover, other biologically active portions, in which other regions of the
20 protein are deleted, can be prepared by recombinant techniques and evaluated for one or more of the functional activities of a native Dkk or Dkk-related protein.

In a preferred embodiment, the Dkk protein has an amino acid sequence shown in SEQ ID NO:2 or an amino acid sequence at least about 55% homologous to SEQ ID NO:2. In another preferred embodiment, the Dkk protein has an amino acid sequence
25 shown in SEQ ID NO:5 or an amino acid sequence at least about 35% homologous to SEQ ID NO:5. In another preferred embodiment, the Dkk protein has an amino acid sequence shown in SEQ ID NO:8 or an amino acid sequence at least about 85% homologous to SEQ ID NO:8. In another preferred embodiment, the Dkk protein has an amino acid sequence shown in SEQ ID NO:21 or an amino acid sequence at least about
30 35% homologous to SEQ ID NO:21. In another preferred embodiment, the protein has an amino acid sequence shown in SEQ ID NO:14 or an amino acid sequence at least

- 50 -

about 60% homologous to SEQ ID NO:14. In still another preferred embodiment, a protein of the present invention comprises an amino acid sequence which is at least about 30-35%, preferably about 40-45%, more preferably about 50-55%, even more preferably about 60-65%, and even more preferably at least about 70-75%, 80-85%, 90-
5 95% or more homologous to the amino acid sequences shown in SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21.

In other embodiments, the protein is substantially homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21, and, preferably, retains the functional activity of the protein of SEQ ID NO:2, SEQ ID NO:5, SEQ ID
10 NO:8, SEQ ID NO:14, or SEQ ID NO:21, yet differs in amino acid sequence due to natural allelic variation or mutagenesis, as described in detail in subsection I above. Accordingly, in another embodiment, the protein is a protein which comprises an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21 and, preferably,
15 retains the functional activity of the proteins of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:11, respectively. Preferably, the protein is at least about 70% homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21, more preferably at least about 80% homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21, even more
20 preferably at least about 90% homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21, and most preferably at least about 95% or more homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21.

To determine the percent homology of two amino acid sequences or of two
25 nucleic acids, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence and non-homologous sequences can be disregarded for comparison purposes). In one embodiment, an alignment is a global alignment, *e.g.*, an overall sequence alignment. In another embodiment, an
30 alignment is a local alignment. In a preferred embodiment, the length of a sequence aligned for comparison purposes is at least 30%, preferably at least 40%, more

preferably at least 50%, even more preferably at least 60%, and even more preferably at least 70%, 80%, or 90% of the length of the reference sequence to which it is aligned (e.g., when aligning a second sequence to the Dkk amino acid sequence of SEQ ID NO:2, at least 105, preferably at least 145, more preferably at least 175, even more preferably at least 210, and even more preferably at least 245, 280 or 315 amino acid residues are aligned). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid or nucleic acid "identity" is equivalent to amino acid or nucleic acid "homology"). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. In a embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch (*J. Mol. Biol.* (48):444-453 (1970)) algorithm which has been incorporated into the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossom 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, 5, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (available at <http://www.gcg.com>), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. In another embodiment, the percent identity between two amino acid or nucleotide sequences is determined using the algorithm of E. Meyers and W. Miller (*CABIOS*, 4:11-17 (1989)) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

The nucleic acid and protein sequences of the present invention can further be used as a "query sequence" to perform a search against public databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, *et al.* (1990) *J. Mol. Biol.* 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to Dkk nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to Dkk protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, (1997) *Nucleic Acids Res.* 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (*e.g.*, XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>.

The invention also provides Dkk or Dkk-related chimeric or fusion proteins. As used herein, a "chimeric protein" or "fusion protein" comprises a Dkk or Dkk-related polypeptide operatively linked to a non-Dkk polypeptide or non-Dkk-related polypeptide. A "Dkk polypeptide" or "Dkk-related polypeptide" refers to a polypeptide having an amino acid sequence corresponding to Dkk or a Dkk-related protein, whereas a "non-Dkk polypeptide" or "non-Dkk-related polypeptide" refers to a polypeptide having an amino acid sequence corresponding to a protein which is not substantially homologous to the Dkk or Dkk-related protein, *e.g.*, a protein which is different from the Dkk or Dkk-related protein and which is derived from the same or a different organism. Within a Dkk or Dkk-related fusion protein the Dkk or Dkk-related polypeptide can correspond to all or a portion of a Dkk or Dkk-related protein. In a preferred embodiment, a Dkk or Dkk-related fusion protein comprises at least one biologically active portion of a Dkk protein. In another preferred embodiment, a Dkk or Dkk-related fusion protein comprises at least two biologically active portions of a Dkk or Dkk-related protein. In another preferred embodiment, a Dkk or Dkk-related fusion protein comprises at least three biologically active portions of a Dkk or Dkk-related protein. Within the fusion protein, the term "operatively linked" is intended to indicate that the Dkk or Dkk-related polypeptide and the non-Dkk or non-Dkk-related polypeptide are

fused in-frame to each other. The non-Dkk or non-Dkk-related polypeptide can be fused to the N-terminus or C-terminus of the Dkk or Dkk-related polypeptide.

For example, in one embodiment, the fusion protein is a GST-Dkk fusion protein in which the Dkk sequences are fused to the C-terminus of the GST sequences. Such
5 fusion proteins can facilitate the purification of recombinant Dkk.

In another embodiment, the fusion protein is a Dkk or Dkk-related protein containing a heterologous signal sequence at its N-terminus. For example, the native Dkk signal sequence (i.e., about amino acids 1 to 23 of SEQ ID NO:2) can be removed and replaced with a signal sequence from another protein. In certain host cells (*e.g.*,
10 mammalian host cells), expression and/or secretion of Dkk or Dkk-related proteins can be increased through use of a heterologous signal sequence.

In yet another embodiment, the fusion protein is a Dkk-immunoglobulin fusion protein in which the Dkk sequences comprising primarily the Dkk cysteine-rich regions are fused to sequences derived from a member of the immunoglobulin protein family.
15 Soluble derivatives have also been made of cell surface glycoproteins in the immunoglobulin gene superfamily consisting of an extracellular domain of the cell surface glycoprotein fused to an immunoglobulin constant (Fc) region (see *e.g.*, Capon, *et al.* (1989) *Nature* 337:525-531 and Capon U.S. Patents 5,116,964 and 5,428,130 [CD4-IgG1 constructs]; Linsley, P.S. *et al.* (1991) *J. Exp. Med.* 173:721-730 [a CD28-IgG1 construct and a B7-1-IgG1 construct]; and Linsley, P.S. *et al.* (1991) *J. Exp. Med.* 174:561-569 and U.S. Patent 5,434,131[a CTLA4-IgG1]). Such fusion proteins have proven useful for modulating receptor-ligand interactions. Soluble derivatives of cell surface proteins of the tumor necrosis factor receptor (TNFR) superfamily proteins have been made consisting of an extracellular domain of the cell surface receptor fused to an
20 immunoglobulin constant (Fc) region (See for example Moreland *et al.* (1997) *N. Engl. J. Med.* 337(3):141-147; van der Poll *et al.* (1997) *Blood* 89(10):3727-3734; and Ammann *et al.* (1997) *J. Clin. Invest.* 99(7):1699-1703.)

The Dkk-immunoglobulin fusion proteins of the invention can be incorporated into pharmaceutical compositions and administered to a subject to inhibit an interaction
30 between a Dkk ligand and a Dkk receptor on the surface of a cell, to thereby suppress Dkk-mediated signal transduction *in vivo*. The Dkk-immunoglobulin fusion proteins

can be used to affect the bioavailability of a Dkk cognate receptor. Inhibition of the Dkk ligand/Dkk interaction may be useful therapeutically for both the treatment of differentiative or proliferative disorders, as well as modulating (*e.g.*, promoting or inhibiting) developmental responses, cell adhesion, and/or cell fate. Moreover, the Dkk-immunoglobulin fusion proteins of the invention can be used as immunogens to produce anti-Dkk antibodies in a subject, to purify Dkk ligands and in screening assays to identify molecules which inhibit the interaction of Dkk with a Dkk ligand.

Preferably, a Dkk or Dkk-related chimeric or fusion protein of the invention is produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques, for example by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and reamplified to generate a chimeric gene sequence (see, for example, *Current Protocols in Molecular Biology*, eds. Ausubel *et al.*, John Wiley & Sons: 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (*e.g.*, a GST polypeptide). A Dkk-encoding nucleic acid or nucleic acid encoding a Dkk-related protein can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the protein.

The present invention also pertains to variants of the Dkk or Dkk-related proteins which function as either agonists (mimetics) or as antagonists. Variants of the Dkk or Dkk-related proteins can be generated by mutagenesis, *e.g.*, discrete point mutation or truncation of a Dkk or Dkk-related protein. An agonist of the Dkk or Dkk-related proteins can retain substantially the same, or a subset, of the biological activities of the naturally occurring form of a Dkk or Dkk-related protein. An antagonist of a Dkk or Dkk-related protein can inhibit one or more of the activities of the naturally occurring

form of the protein by, for example, competitively binding to a downstream or upstream member of a cellular signaling cascade which includes the Dkk or Dkk-related protein. Thus, specific biological effects can be elicited by treatment with a variant of limited function. In one embodiment, treatment of a subject with a variant having a subset of
5 the biological activities of the naturally occurring form of the protein has fewer side effects in a subject relative to treatment with the naturally occurring form of the Dkk or Dkk-related protein.

In one embodiment, variants of a Dkk or Dkk-related protein which function as either agonists (mimetics) or as antagonists can be identified by screening combinatorial
10 libraries of mutants, *e.g.*, truncation mutants, of a Dkk or Dkk-related protein for protein agonist or antagonist activity. In one embodiment, a variegated library of Dkk or Dkk-related variants is generated by combinatorial mutagenesis at the nucleic acid level and is encoded by a variegated gene library. A variegated library of Dkk or Dkk-related
15 oligonucleotides into gene sequences such that a degenerate set of potential Dkk or Dkk-related sequences is expressible as individual polypeptides, or alternatively, as a set of larger fusion proteins (*e.g.*, for phage display) containing the set of Dkk or Dkk-related sequences therein. There are a variety of methods which can be used to produce
20 libraries of potential Dkk or Dkk-related variants from a degenerate oligonucleotide sequence. Chemical synthesis of a degenerate gene sequence can be performed in an automatic DNA synthesizer, and the synthetic gene then ligated into an appropriate expression vector. Use of a degenerate set of genes allows for the provision, in one mixture, of all of the sequences encoding the desired set of potential Dkk or Dkk-related
25 sequences. Methods for synthesizing degenerate oligonucleotides are known in the art (see, *e.g.*, Narang, S.A. (1983) *Tetrahedron* 39:3; Itakura *et al.*, (1984) *Annu. Rev. Biochem.* 53:323; Itakura *et al.*, (1984) *Science* 198:1056; Ike *et al.*, (1983) *Nucleic Acid Res.* 11:477.

In addition, libraries of fragments of a Dkk or Dkk-related protein coding sequence can be used to generate a variegated population of Dkk or Dkk-related
30 fragments for screening and subsequent selection of variants of a Dkk or Dkk-related protein. In one embodiment, a library of coding sequence fragments can be generated by

- 56 -

treating a double stranded PCR fragment of a Dkk coding sequence with a nuclease under conditions wherein nicking occurs only about once per molecule, denaturing the double stranded DNA, renaturing the DNA to form double stranded DNA which can include sense/antisense pairs from different nicked products, removing single stranded portions from reformed duplexes by treatment with S1 nuclease, and ligating the resulting fragment library into an expression vector. By this method, an expression library can be derived which encodes N-terminal, C-terminal and internal fragments of various sizes of the Dkk protein.

Several techniques are known in the art for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene products having a selected property. Such techniques are adaptable for rapid screening of the gene libraries generated by the combinatorial mutagenesis of Dkk or Dkk-related proteins. The most widely used techniques, which are amenable to high through-put analysis, for screening large gene libraries typically include cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and expressing the combinatorial genes under conditions in which detection of a desired activity facilitates isolation of the vector encoding the gene whose product was detected. Recursive ensemble mutagenesis (REM), a new technique which enhances the frequency of functional mutants in the libraries, can be used in combination with the screening assays to identify Dkk variants (Arkin and Yourvan (1992) *PNAS* 89:7811-7815; Delgrave *et al.*, (1993) *Protein Engineering* 6(3):327-331).

In one embodiment, cell based assays can be exploited to analyze a variegated Dkk or Dkk-related library. For example, a library of expression vectors can be transfected into a cell line which ordinarily responds to a particular ligand in a Dkk-dependent manner. The transfected cells are then contacted with the ligand and the effect of expression of the mutant on signaling by the ligand can be detected, *e.g.*, by measuring any of a number of immune cell responses. Plasmid DNA can then be recovered from the cells which score for inhibition, or alternatively, potentiation of ligand induction, and the individual clones further characterized.

An isolated Dkk protein, Dkk-related protein, or a portion or fragment thereof, can be used as an immunogen to generate antibodies that bind Dkk or Dkk-related proteins using standard techniques for polyclonal and monoclonal antibody preparation. A full-length Dkk or Dkk-related protein can be used or, alternatively, the invention
5 provides antigenic peptide fragments for use as immunogens. The antigenic peptide of Dkk comprises at least 8 amino acid residues of the amino acid sequence shown in SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21 and encompasses an epitope of Dkk or Dkk-related protein such that an antibody raised
10 against the peptide forms a specific immune complex with the protein. Preferably, the antigenic peptide comprises at least 10 amino acid residues, more preferably at least 15 amino acid residues, even more preferably at least 20 amino acid residues, and most preferably at least 30 amino acid residues.

Preferred epitopes encompassed by the antigenic peptide are regions of Dkk or Dkk-related proteins that are located on the surface of the protein, *e.g.*, hydrophilic
15 regions.

A Dkk or Dkk-related immunogen typically is used to prepare antibodies by immunizing a suitable subject, (*e.g.*, rabbit, goat, mouse or other mammal) with the immunogen. An appropriate immunogenic preparation can contain, for example, recombinantly expressed Dkk or Dkk-related protein or a chemically synthesized Dkk or
20 Dkk-related polypeptide. The preparation can further include an adjuvant, such as Freund's complete or incomplete adjuvant, or similar immunostimulatory agent. Immunization of a suitable subject with an immunogenic Dkk preparation, for example, induces a polyclonal anti-Dkk antibody response.

Accordingly, another aspect of the invention pertains to anti-Dkk antibodies as
25 well as antibodies to Dkk-related proteins. The term "antibody" as used herein refers to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, *i.e.*, molecules that contain an antigen binding site which specifically binds (immunoreacts with) an antigen, such as Dkk or Dkk-related antigens. Examples of immunologically active portions of immunoglobulin molecules include F(ab) and
30 F(ab')₂ fragments which can be generated by treating the antibody with an enzyme such as pepsin. The invention provides polyclonal and monoclonal antibodies that bind Dkk

or Dkk-related polypeptides. The term "monoclonal antibody" or "monoclonal antibody composition", as used herein, refers to a population of antibody molecules that contain only one species of an antigen binding site capable of immunoreacting with a particular epitope of Dkk or a or Dkk-related protein. A monoclonal antibody composition thus typically displays a single binding affinity for a particular Dkk or Dkk-related protein with which it immunoreacts.

Polyclonal antibodies can be prepared as described above by immunizing a suitable subject with a Dkk or Dkk-related immunogen. The antibody titer in the immunized subject can be monitored over time by standard techniques, such as with an enzyme linked immunosorbent assay (ELISA) using immobilized Dkk or Dkk-related protein. If desired, the antibody molecules directed against Dkk or Dkk-related protein can be isolated from the mammal (*e.g.*, from the blood) and further purified by well known techniques, such as protein A chromatography to obtain the IgG fraction. At an appropriate time after immunization, *e.g.*, when the antibody titers are highest, antibody-producing cells can be obtained from the subject and used to prepare monoclonal antibodies by standard techniques, such as the hybridoma technique originally described by Kohler and Milstein (1975) *Nature* 256:495-497) (see also, Brown *et al.*, (1981) *J. Immunol.* 127:539-46; Brown *et al.*, (1980) *J. Biol. Chem.* 255:4980-83; Yeh *et al.*, (1976) *PNAS* 76:2927-31; and Yeh *et al.*, (1982) *Int. J. Cancer* 29:269-75), the more recent human B cell hybridoma technique (Kozbor *et al.*, (1983) *Immunol Today* 4:72), the EBV-hybridoma technique (Cole *et al.*, (1985), *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96) or trioma techniques. The technology for producing monoclonal antibody hybridomas is well known (see generally R. H. Kenneth, in *Monoclonal Antibodies: A New Dimension In Biological Analyses*, Plenum Publishing Corp., New York, New York (1980); E. A. Lerner (1981) *Yale J. Biol. Med.*, 54:387-402; M.L. Gefter *et al.*, (1977) *Somatic Cell Genet.* 3:231-36). Briefly, an immortal cell line (typically a myeloma) is fused to lymphocytes (typically splenocytes) from a mammal immunized with a Dkk or Dkk-related immunogen as described above, and the culture supernatants of the resulting hybridoma cells are screened to identify a hybridoma producing a monoclonal antibody that binds Dkk or Dkk-related protein.

Any of the many well known protocols used for fusing lymphocytes and immortalized cell lines can be applied for the purpose of generating a monoclonal antibody (see, *e.g.*, G. Galfre *et al.*, (1977) *Nature* 266:55052; Gefter *et al.*, *Somatic Cell Genet.*, cited *supra*; Lerner, *Yale J. Biol. Med.*, cited *supra*; Kenneth, *Monoclonal Antibodies*, cited *supra*). Moreover, the ordinarily skilled worker will appreciate that there are many variations of such methods which also would be useful. Typically, the immortal cell line (*e.g.*, a myeloma cell line) is derived from the same mammalian species as the lymphocytes. For example, murine hybridomas can be made by fusing lymphocytes from a mouse immunized with an immunogenic preparation of the present invention with an immortalized mouse cell line. Preferred immortal cell lines are mouse myeloma cell lines that are sensitive to culture medium containing hypoxanthine, aminopterin and thymidine ("HAT medium"). Any of a number of myeloma cell lines can be used as a fusion partner according to standard techniques, *e.g.*, the P3-NS1/1-Ag4-1, P3-x63-Ag8.653 or Sp2/O-Ag14 myeloma lines. These myeloma lines are available from ATCC. Typically, HAT-sensitive mouse myeloma cells are fused to mouse splenocytes using polyethylene glycol ("PEG"). Hybridoma cells resulting from the fusion are then selected using HAT medium, which kills unfused and unproductively fused myeloma cells (unfused splenocytes die after several days because they are not transformed). Hybridoma cells producing a monoclonal antibody of the invention are detected by screening the hybridoma culture supernatants for antibodies that bind Dkk or Dkk-related protein, *e.g.*, using a standard ELISA assay.

Alternative to preparing monoclonal antibody-secreting hybridomas, a monoclonal antibody can be identified and isolated by screening a recombinant combinatorial immunoglobulin library (*e.g.*, an antibody phage display library) with Dkk or Dkk-related protein to thereby isolate immunoglobulin library members that bind Dkk or Dkk-related protein. Kits for generating and screening phage display libraries are commercially available (*e.g.*, the Pharmacia *Recombinant Phage Antibody System*, Catalog No. 27-9400-01; and the Stratagene *SurfZAP™ Phage Display Kit*, Catalog No. 240612). Additionally, examples of methods and reagents particularly amenable for use in generating and screening antibody display library can be found in, for example, Ladner *et al.*, U.S. Patent No. 5,223,409; Kang *et al.*, PCT International Publication No.

WO 92/18619; Dower *et al.*, PCT International Publication No. WO 91/17271; Winter *et al.*, PCT International Publication WO 92/20791; Markland *et al.*, PCT International Publication No. WO 92/15679; Breitling *et al.*, PCT International Publication WO 93/01288; McCafferty *et al.*, PCT International Publication No. WO 92/01047; Garrard
5 *et al.*, PCT International Publication No. WO 92/09690; Ladner *et al.*, PCT International Publication No. WO 90/02809; Fuchs *et al.*, (1991) *Bio/Technology* 9:1370-1372; Hay *et al.*, (1992) *Hum. Antibod. Hybridomas* 3:81-85; Huse *et al.*, (1989) *Science* 246:1275-1281; Griffiths *et al.*, (1993) *EMBO J* 12:725-734; Hawkins *et al.*, (1992) *J. Mol. Biol.* 226:889-896; Clarkson *et al.*, (1991) *Nature* 352:624-628; Gram *et al.*, (1992) *PNAS*
10 89:3576-3580; Garrad *et al.*, (1991) *Bio/Technology* 9:1373-1377; Hoogenboom *et al.*, (1991) *Nuc. Acid Res.* 19:4133-4137; Barbas *et al.*, (1991) *PNAS* 88:7978-7982; and McCafferty *et al.*, *Nature* (1990) 348:552-554.

Additionally, recombinant antibodies, such as chimeric and humanized monoclonal antibodies, comprising both human and non-human portions, which can be
15 made using standard recombinant DNA techniques, are within the scope of the invention. Such chimeric and humanized monoclonal antibodies can be produced by recombinant DNA techniques known in the art, for example using methods described in Robinson *et al.*, International Application No. PCT/US86/02269; Akira, *et al.*, European Patent Application 184,187; Taniguchi, M., European Patent Application 171,496;
20 Morrison *et al.*, European Patent Application 173,494; Neuberger *et al.*, PCT International Publication No. WO 86/01533; Cabilly *et al.*, U.S. Patent No. 4,816,567; Cabilly *et al.*, European Patent Application 125,023; Better *et al.*, (1988) *Science* 240:1041-1043; Liu *et al.*, (1987) *PNAS* 84:3439-3443; Liu *et al.*, (1987) *J. Immunol.* 139:3521-3526; Sun *et al.*, (1987) *PNAS* 84:214-218; Nishimura *et al.*, (1987) *Canc.*
25 *Res.* 47:999-1005; Wood *et al.*, (1985) *Nature* 314:446-449; and Shaw *et al.*, (1988) *J. Natl. Cancer Inst.* 80:1553-1559; Morrison, S. L. (1985) *Science* 229:1202-1207; Oi *et al.*, (1986) *BioTechniques* 4:214; Winter U.S. Patent 5,225,539; Jones *et al.*, (1986) *Nature* 321:552-525; Verhoeyan *et al.*, (1988) *Science* 239:1534; and Beidler *et al.*, (1988) *J. Immunol.* 141:4053-4060.

An antibody (*e.g.*, monoclonal antibody) can be used to isolate Dkk or Dkk-related protein by standard techniques, such as affinity chromatography or immunoprecipitation. An antibody can facilitate the purification of natural Dkk or Dkk-related protein from cells and of recombinantly produced Dkk or Dkk-related protein expressed in host cells. Moreover, an antibody can be used to detect Dkk or Dkk-related protein (*e.g.*, in a cellular lysate or cell supernatant) in order to evaluate the abundance and pattern of expression of the Dkk or Dkk-related protein. Antibodies can be used diagnostically to monitor protein levels in tissue as part of a clinical testing procedure, *e.g.*, to, for example, determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling (*i.e.*, physically linking) the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

III. Recombinant Expression Vectors and Host Cells

Another aspect of the invention pertains to vectors, preferably expression vectors, containing a nucleic acid encoding Dkk or a nucleic acid encoding a Dkk-related protein (or a portion thereof). As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (*e.g.*, bacterial vectors having a bacterial origin of replication and episomal

- 62 -

mammalian vectors). Other vectors (*e.g.*, non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "expression vectors". In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" can be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (*e.g.*, replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

The recombinant expression vectors of the invention comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, which is operatively linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, "operably linked" is intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner which allows for expression of the nucleotide sequence (*e.g.*, in an *in vitro* transcription/translation system or in a host cell when the vector is introduced into the host cell). The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (*e.g.*, polyadenylation signals). Such regulatory sequences are described, for example, in Goeddel; *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Regulatory sequences include those which direct constitutive expression of a nucleotide sequence in many types of host cell and those which direct expression of the nucleotide sequence only in certain host cells (*e.g.*, tissue-specific regulatory sequences). It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. The expression vectors of the invention can be introduced into host cells to thereby produce proteins or peptides, including fusion proteins or peptides, encoded by nucleic acids as

described herein (*e.g.*, Dkk proteins, Dkk-related proteins, mutant forms of Dkk or Dkk-related proteins, fusion proteins, etc.).

The recombinant expression vectors of the invention can be designed for expression of Dkk or Dkk-related proteins in prokaryotic or eukaryotic cells. For example, Dkk can be expressed in bacterial cells such as *E. coli*, insect cells (using baculovirus expression vectors) yeast cells or mammalian cells. Suitable host cells are discussed further in Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Alternatively, the recombinant expression vector can be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

Expression of proteins in prokaryotes is most often carried out in *E. coli* with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein encoded therein, usually to the amino terminus of the recombinant protein. Such fusion vectors typically serve three purposes: 1) to increase expression of recombinant protein; 2) to increase the solubility of the recombinant protein; and 3) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase. Typical fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith, D.B. and Johnson, K.S. (1988) *Gene* 67:31-40), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein.

Purified fusion proteins can be utilized in activity assays, in ligand binding (*e.g.*, direct assays or competitive assays described in detail below), to generate antibodies specific for Dkk or Dkk-related proteins, as examples. In a preferred embodiment, a Dkk or Dkk-related fusion expressed in a retroviral expression vector of the present invention can be utilized to infect bone marrow cells which are subsequently

transplanted into irradiated recipients. The pathology of the subject recipient is then examined after sufficient time has passed (e.g six (6) weeks).

Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amann *et al.*, (1988) *Gene* 69:301-315) and pET 11d (Studier *et al.*, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 60-89). Target gene expression from the pTrc vector relies on host RNA polymerase transcription from a hybrid trp-lac fusion promoter. Target gene expression from the pET 11d vector relies on transcription from a T7 gn10-lac fusion promoter mediated by a coexpressed viral RNA polymerase (T7 gn1). This viral polymerase is supplied by host strains BL21(DE3) or HMS174(DE3) from a resident λ prophage harboring a T7 gn1 gene under the transcriptional control of the lacUV 5 promoter.

One strategy to maximize recombinant protein expression in *E. coli* is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant protein (Gottesman, S., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 119-128). Another strategy is to alter the nucleic acid sequence of the nucleic acid to be inserted into an expression vector so that the individual codons for each amino acid are those preferentially utilized in *E. coli* (Wada *et al.*, (1992) *Nucleic Acids Res.* 20:2111-2118). Such alteration of nucleic acid sequences of the invention can be carried out by standard DNA synthesis techniques.

In another embodiment, the Dkk or Dkk-related expression vector is a yeast expression vector. Examples of vectors for expression in yeast *S. cerevisiae* include pYepSec1 (Baldari, *et al.*, (1987) *Embo J.* 6:229-234), pMFa (Kurjan and Herskowitz, (1982) *Cell* 30:933-943), pJRY88 (Schultz *et al.*, (1987) *Gene* 54:113-123), pYES2 (Invitrogen Corporation, San Diego, CA), and picZ (Invitrogen Corp, San Diego, CA).

Alternatively, Dkk or Dkk-related protein can be expressed in insect cells using baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., Sf 9 cells) include the pAc series (Smith *et al.*, (1983) *Mol. Cell Biol.* 3:2156-2165) and the pVL series (Lucklow and Summers (1989) *Virology* 170:31-39).

In yet another embodiment, a nucleic acid of the invention is expressed in mammalian cells using a mammalian expression vector. Examples of mammalian expression vectors include pCDM8 (Seed, B. (1987) *Nature* 329:840) and pMT2PC (Kaufman *et al.*, (1987) *EMBO J.* 6:187-195). When used in mammalian cells, the expression vector's control functions are often provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma, Adenovirus 2, cytomegalovirus and Simian Virus 40. For other suitable expression systems for both prokaryotic and eukaryotic cells see chapters 16 and 17 of Sambrook, J., Fritsch, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.

In another embodiment, the recombinant mammalian expression vector is capable of directing expression of the nucleic acid preferentially in a particular cell type (*e.g.*, tissue-specific regulatory elements are used to express the nucleic acid). Tissue-specific regulatory elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert *et al.*, (1987) *Genes Dev.* 1:268-277), lymphoid-specific promoters (Calame and Eaton (1988) *Adv. Immunol.* 43:235-275), in particular promoters of T cell receptors (Winoto and Baltimore (1989) *EMBO J.* 8:729-733) and immunoglobulins (Banerji *et al.*, (1983) *Cell* 33:729-740; Queen and Baltimore (1983) *Cell* 33:741-748), neuron-specific promoters (*e.g.*, the neurofilament promoter; Byrne and Ruddle (1989) *PNAS* 86:5473-5477), pancreas-specific promoters (Edlund *et al.*, (1985) *Science* 230:912-916), and mammary gland-specific promoters (*e.g.*, milk whey promoter; U.S. Patent No. 4,873,316 and European Application Publication No. 264,166). Developmentally-regulated promoters are also encompassed, for example the murine hox promoters (Kessel and Gruss (1990) *Science* 249:374-379) and the α -fetoprotein promoter (Campes and Tilghman (1989) *Genes Dev.* 3:537-546).

The invention further provides a recombinant expression vector comprising a DNA molecule of the invention cloned into the expression vector in an antisense orientation. That is, the DNA molecule is operatively linked to a regulatory sequence in a manner which allows for expression (by transcription of the DNA molecule) of an

RNA molecule which is antisense to Dkk mRNA or a Dkk-related mRNA. Regulatory sequences operatively linked to a nucleic acid cloned in the antisense orientation can be chosen which direct the continuous expression of the antisense RNA molecule in a variety of cell types, for instance viral promoters and/or enhancers, or regulatory sequences can be chosen which direct constitutive, tissue specific or cell type specific expression of antisense RNA. The antisense expression vector can be in the form of a recombinant plasmid, phagemid or attenuated virus in which antisense nucleic acids are produced under the control of a high efficiency regulatory region, the activity of which can be determined by the cell type into which the vector is introduced. For a discussion of the regulation of gene expression using antisense genes see Weintraub, H. *et al.*, Antisense RNA as a molecular tool for genetic analysis, *Reviews - Trends in Genetics*, Vol. 1(1) 1986.

Another aspect of the invention pertains to host cells into which a recombinant expression vector of the invention has been introduced. The terms "host cell" and "recombinant host cell" are used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

A host cell can be any prokaryotic or eukaryotic cell. For example, Dkk protein can be expressed in bacterial cells such as *E. coli*, insect cells, yeast or mammalian cells (such as Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to those skilled in the art.

Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (*e.g.*, DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation. Suitable methods for transforming or transfecting host cells can be found in Sambrook, *et al.*, (*Molecular Cloning: A*

Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989), and other laboratory manuals.

For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may
5 integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (*e.g.*, resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Preferred selectable markers include those which confer resistance to drugs, such as G418, hygromycin and methotrexate. Nucleic acid encoding a selectable marker can be
10 introduced into a host cell on the same vector as that encoding Dkk or can be introduced on a separate vector. Cells stably transfected with the introduced nucleic acid can be identified by drug selection (*e.g.*, cells that have incorporated the selectable marker gene will survive, while the other cells die).

A host cell of the invention, such as a prokaryotic or eukaryotic host cell in
15 culture, can be used to produce (*i.e.*, express) a Dkk or Dkk-related protein. Accordingly, the invention further provides methods for producing Dkk or Dkk-related proteins using the host cells of the invention. In one embodiment, the method comprises culturing the host cell of invention (into which a recombinant expression vector encoding Dkk or a Dkk-related protein has been introduced) in a suitable medium such
20 that protein is produced. In another embodiment, the method further comprises isolating Dkk or a Dkk-related protein from the medium or the host cell.

The host cells of the invention can also be used to produce nonhuman transgenic animals. For example, in one embodiment, a host cell of the invention is a fertilized oocyte or an embryonic stem cell into which Dkk-coding sequences (or Dkk-related
25 coding sequences) have been introduced. Such host cells can then be used to create non-human transgenic animals in which exogenous Dkk sequences (or Dkk-related sequences) have been introduced into their genome or homologous recombinant animals in which endogenous Dkk sequences (or Dkk-related sequences) have been altered. Such animals are useful for studying the function and/or activity of Dkk or Dkk-related
30 proteins and for identifying and/or evaluating modulators of Dkk or Dkk-related protein activity. As used herein, a "transgenic animal" is a non-human animal, preferably a

mammal, more preferably a rodent such as a rat or mouse, in which one or more of the cells of the animal includes a transgene. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, amphibians, etc. A transgene is exogenous DNA which is integrated into the genome of a cell from which a transgenic animal develops and which remains in the genome of the mature animal, thereby directing the expression of an encoded gene product in one or more cell types or tissues of the transgenic animal. As used herein, a "homologous recombinant animal" is a non-human animal, preferably a mammal, more preferably a mouse, in which an endogenous Dkk or Dkk-related gene has been altered by homologous recombination between the endogenous gene and an exogenous DNA molecule introduced into a cell of the animal, e.g., an embryonic cell of the animal, prior to development of the animal.

A transgenic animal of the invention can be created, for example, by introducing Dkk-encoding nucleic acid into the male pronuclei of a fertilized oocyte, e.g., by microinjection, retroviral infection, and allowing the oocyte to develop in a pseudopregnant female foster animal. The human Dkk cDNA sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, or SEQ ID NO:20 can be introduced as a transgene into the genome of a non-human animal. Alternatively, a nonhuman homologue of a human Dkk gene, such as a mouse Dkk gene, can be isolated based on hybridization to the human Dkk cDNA (described further in subsection I above) and used as a transgene.

Intronic sequences and polyadenylation signals can also be included in the transgene to increase the efficiency of expression of the transgene. A tissue-specific regulatory sequence(s) can be operably linked to the Dkk transgene to direct expression of Dkk protein to particular cells. Methods for generating transgenic animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009, both by Leder *et al.*, U.S. Patent No. 4,873,191 by Wagner *et al.*, and in Hogan, B., *Manipulating the Mouse Embryo*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986). Similar methods are used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of the Dkk transgene in its genome and/or expression of Dkk mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional

animals carrying the transgene. Moreover, transgenic animals carrying a transgene encoding Dkk can further be bred to other transgenic animals carrying other transgenes.

To create a homologous recombinant animal, a vector is prepared which contains at least a portion of a Dkk gene into which a deletion, addition or substitution has been introduced to thereby alter, *e.g.*, functionally disrupt, the Dkk gene. The Dkk gene can be a human gene (*e.g.*, the cDNA of SEQ ID NO:3, SEQ ID NO: 6, SEQ ID NO:9 or SEQ ID NO:22), but more preferably, is a non-human homologue of a human Dkk gene. For example, a mouse Dkk gene of SEQ ID NO:16 can be used to construct a homologous recombination vector suitable for altering an endogenous Dkk gene in the mouse genome. In a preferred embodiment, the vector is designed such that, upon homologous recombination, the endogenous Dkk gene is functionally disrupted (*i.e.*, no longer encodes a functional protein; also referred to as a "knock out" vector). Alternatively, the vector can be designed such that, upon homologous recombination, the endogenous Dkk gene is mutated or otherwise altered but still encodes functional protein (*e.g.*, the upstream regulatory region can be altered to thereby alter the expression of the endogenous Dkk protein). In the homologous recombination vector, the altered portion of the Dkk gene is flanked at its 5' and 3' ends by additional nucleic acid of the Dkk gene to allow for homologous recombination to occur between the exogenous Dkk gene carried by the vector and an endogenous Dkk gene in an embryonic stem cell. The additional flanking Dkk nucleic acid is of sufficient length for successful homologous recombination with the endogenous gene. Typically, several kilobases of flanking DNA (both at the 5' and 3' ends) are included in the vector (see *e.g.*, Thomas, K.R. and Capecchi, M. R. (1987) *Cell* 51:503 for a description of homologous recombination vectors). The vector is introduced into an embryonic stem cell line (*e.g.*, by electroporation) and cells in which the introduced Dkk gene has homologously recombined with the endogenous Dkk gene are selected (see *e.g.*, Li, E. *et al.*, (1992) *Cell* 69:915). The selected cells are then injected into a blastocyst of an animal (*e.g.*, a mouse) to form aggregation chimeras (see *e.g.*, Bradley, A. in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E.J. Robertson, ed. (IRL, Oxford, 1987) pp. 113-152). A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term. Progeny harboring the

homologously recombined DNA in their germ cells can be used to breed animals in which all cells of the animal contain the homologously recombined DNA by germline transmission of the transgene. Methods for constructing homologous recombination vectors and homologous recombinant animals are described further in Bradley, A.

5 (1991) *Current Opinion in Biotechnology* 2:823-829 and in PCT International Publication Nos.: WO 90/11354 by Le Mouellec *et al.*; WO 91/01140 by Smithies *et al.*; WO 92/0968 by Zijlstra *et al.*; and WO 93/04169 by Berns *et al.* It is also within the scope of the present invention to practice the above-described transgenic methodology utilizing nucleic acid molecules which encode Dkk-related proteins.

10 In another embodiment, transgenic non-humans animals can be produced which contain selected systems which allow for regulated expression of the transgene. One example of such a system is the *cre/loxP* recombinase system of bacteriophage P1. For a description of the *cre/loxP* recombinase system, see, *e.g.*, Lakso *et al.*, (1992) *PNAS* 89:6232-6236. Another example of a recombinase system is the FLP recombinase
15 system of *Saccharomyces cerevisiae* (O'Gorman *et al.*, (1991) *Science* 251:1351-1355. If a *cre/loxP* recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the *Cre* recombinase and a selected protein are required. Such animals can be provided through the construction of "double" transgenic animals, *e.g.*, by mating two transgenic animals, one containing a transgene encoding a
20 selected protein and the other containing a transgene encoding a recombinase.

Clones of the non-human transgenic animals described herein can also be produced according to the methods described in Wilmut, I. *et al.*, (1997) *Nature* 385:810-813. In brief, a cell, *e.g.*, a somatic cell, from the transgenic animal can be isolated and induced to exit the growth cycle and enter G₀ phase. The quiescent cell can
25 then be fused, *e.g.*, through the use of electrical pulses, to an enucleated oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops to morula or blastocyte and then transferred to pseudopregnant female foster animal. The offspring borne of this female foster animal will be a clone of the animal from which the cell, *e.g.*, the somatic cell, is
30 isolated.

IV. Pharmaceutical Compositions

The Dkk and Dkk-related nucleic acid molecules, Dkk and Dkk-related proteins, and anti-Dkk or anti-Dkk-related protein antibodies (also referred to herein as "active compounds") of the invention can be incorporated into pharmaceutical compositions
5 suitable for administration. Such compositions typically comprise the nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier. As used herein the language "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration.
10 The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

A pharmaceutical composition of the invention is formulated to be compatible
15 with its intended route of administration. Examples of routes of administration include parenteral, *e.g.*, intravenous, intradermal, subcutaneous, oral (*e.g.*, inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils,
20 polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases,
25 such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the
30 extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic

water, Cremophor EL™ (BASF, Parsippany, NJ) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as

5 bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of

10 surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be

15 brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound (*e.g.*, a Dkk protein, Dkk-related protein or antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as

20 required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a

25 powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and

30 used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is

applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as

5 microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

10 For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant, *e.g.*, a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be
15 permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in
20 the art.

The compounds can also be prepared in the form of suppositories (*e.g.*, with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

In one embodiment, the active compounds are prepared with carriers that will
25 protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art.
30 The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected

cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD50/ED50. Compounds which exhibit large therapeutic indices are preferred. While compounds that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC50 (*i.e.*, the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine

useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

The nucleic acid molecules of the invention can be inserted into vectors and used as gene therapy vectors. Gene therapy vectors can be delivered to a subject by, for example, intravenous injection, local administration (see U.S. Patent 5,328,470) or by stereotactic injection (see *e.g.*, Chen *et al.*, (1994) *PNAS* 91:3054-3057). The pharmaceutical preparation of the gene therapy vector can include the gene therapy vector in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from recombinant cells, *e.g.*, retroviral vectors, the pharmaceutical preparation can include one or more cells which produce the gene delivery system.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

V. Uses and Methods of the Invention

The molecules of the present invention (*e.g.*, nucleic acid molecules, proteins, protein homologues, and antibodies described herein) can be used in one or more of the following methods: a) screening assays; b) predictive medicine (*e.g.*, diagnostic assays, prognostic assays, monitoring clinical trials, and pharmacogenetics); and c) methods of treatment (*e.g.*, therapeutic and prophylactic). As described herein, a Dkk protein of the invention has one or more of the following activities: intracellular calcium, an increase in phosphatidylinositol or other molecule, and can result, *e.g.*, in phosphorylation of specific proteins, a modulation of gene transcription and any of the other biological activities set forth herein.

In a preferred embodiment, a Dkk activity is at least one or more of the following activities: (i) interaction of a Dkk protein with and/or binding to a second molecule, (*e.g.*, a protein, such as a Dkk (*e.g.*, hDkk-3) receptor, a soluble form of a Dkk receptor, a receptor for a member of the *wnt* family of signaling proteins, or a non-Dkk signaling molecule); (ii) interaction of a Dkk protein with an intracellular protein via a membrane-bound Dkk receptor; (iii) complex formation between a soluble Dkk protein and a

- 76 -

second soluble Dkk binding partner (*e.g.*, a non-Dkk protein molecule or a second Dkk protein molecule); (iv) interaction with other extracellular proteins (*e.g.*, regulation of *wnt*-dependent cellular adhesion to extracellular matrix components); (v) binding to and eliminating an undesirable molecule (*e.g.*, a detoxifying activity or defense function);
5 and/or (vi) an enzymatic activity, and can thus be used in, for example, (1) modulation of cellular signal transduction, either *in vitro* or *in vivo* (*e.g.*, antagonism of the activity of members of the *wnt* family of secreted proteins or suppression of *wnt*-dependent signal transduction); (2) regulation of communication between cells (*e.g.*, regulation of *wnt*-dependent cell-cell interactions); (3) regulation of expression of genes
10 whose expression is modulated by binding of Dkk (*e.g.*, hDkk-3) to a receptor; (4) regulation of gene transcription in a cell involved in development or differentiation, either *in vitro* or *in vivo* (*e.g.*, induction of cellular differentiation); (5) regulation of gene transcription in a cell involved in development or differentiation, wherein at least one gene encodes a differentiation-specific protein; (6) regulation of gene transcription
15 in a cell involved in development or differentiation, wherein at least one gene encodes a second secreted protein; (7) regulation of gene transcription in a cell involved in development or differentiation, wherein at least one gene encodes a signal transduction molecule; (8) regulation of cellular proliferation, either *in vitro* or *in vivo* (*e.g.*, induction of cellular proliferation or inhibition of proliferation, for example, inhibition of
20 tumorigenesis (*e.g.*, inhibition of glioblastoma proliferation)); (9) formation and maintenance of ordered spatial arrangements of differentiated tissues in vertebrates, both adult and embryonic (*e.g.*, induction of head formation during vertebrate development or maintenance of hematopoietic progenitor cells); (10) modulation of cell death, such as stimulation of cell survival; (11) regulating cell migration; and/or (12) immune
25 modulation.

Accordingly one embodiment of the present invention involves a method of use (*e.g.*, a diagnostic assay, prognostic assay, or a prophylactic/therapeutic method of treatment) wherein a molecule of the present invention (*e.g.*, a Dkk protein, Dkk nucleic acid, or a Dkk modulator) is used, for example, to diagnose, prognose and/or treat a
30 disease and/or condition in which any of the aforementioned activities (*i.e.*, activities (i) - (vi) and (1) - (12) in the above paragraph) is indicated. In another embodiment, the

present invention involves a method of use (*e.g.*, a diagnostic assay, prognostic assay, or a prophylactic/therapeutic method of treatment) wherein a molecule of the present invention (*e.g.*, a Dkk protein, Dkk nucleic acid, or a Dkk modulator) is used, for example, for the diagnosis, prognosis, and/or treatment of subjects, preferably a human subject, in which any of the aforementioned activities is pathologically perturbed. In a preferred embodiment, the methods of use (*e.g.*, diagnostic assays, prognostic assays, or prophylactic/therapeutic methods of treatment) involve administering to a subject, preferably a human subject, a molecule of the present invention (*e.g.*, a Dkk protein, Dkk nucleic acid, or a Dkk modulator) for the diagnosis, prognosis, and/or therapeutic treatment. In another embodiment, the methods of use (*e.g.*, diagnostic assays, prognostic assays, or prophylactic/therapeutic methods of treatment) involve administering to a human subject a molecule of the present invention (*e.g.*, a Dkk protein, Dkk nucleic acid, or a Dkk modulator).

Other embodiments of the invention pertain to the use of isolated nucleic acid molecules of the invention can be used, for example, to express Dkk or Dkk-related protein (*e.g.*, via a recombinant expression vector in a host cell in gene therapy applications), to detect Dkk or Dkk-related mRNA (*e.g.*, in a biological sample) or a genetic alteration in a Dkk or Dkk-related gene, and to modulate Dkk or Dkk-related activity, as described further below. In addition, the Dkk or Dkk-related proteins can be used to screen drugs or compounds which modulate the Dkk activity as well as to treat disorders characterized by insufficient or excessive production of Dkk or Dkk-related protein or production of Dkk or Dkk-related protein forms which have decreased or aberrant activity compared to Dkk or Dkk-related wild type protein (*e.g.*, developmental disorders or proliferative diseases such as cancer as well as diseases, ocular disorders (*e.g.*, blindness) conditions or disorders characterized by abnormal cell differentiation and/or survival, an abnormal extracellular structure, or an abnormality in a defense mechanism). Moreover, the antibodies of the invention can be used to detect and isolate Dkk or Dkk-related proteins, regulate the bioavailability of Dkk or Dkk-related proteins, and modulate Dkk or Dkk-related activity. The term "an aberrant activity", as applied to an activity of a protein such as Dkk (*e.g.*, hDkk-3), refers to an activity which differs from the activity of the wild-type or native protein or which differs from the activity of

the protein in a healthy subject. An activity of a protein can be aberrant because it is stronger than the activity of its native counterpart. Alternatively, an activity can be aberrant because it is weaker or absent related to the activity of its native counterpart. An aberrant activity can also be a change in an activity. For example an aberrant protein
5 can interact with a different protein relative to its native counterpart. A cell can have an aberrant Dkk (*e.g.*, hDkk-3) activity due to overexpression or underexpression of the gene encoding Dkk.

A. Screening Assays:

10 The invention provides a method (also referred to herein as a "screening assay") for identifying modulators, *i.e.*, candidate or test compounds or agents (*e.g.*, peptides, peptidomimetics, small molecules or other drugs) which bind to Dkk or Dkk-related proteins or have a stimulatory or inhibitory effect on, for example, Dkk or Dkk-related expression or activity. Modulators can include, for example, agonists and/or
15 antagonists. The term "agonist", as used herein, is meant to refer to an agent that mimics or upregulates (*e.g.* potentiates or supplements) a Dkk or Dkk-related (*e.g.*, hDkk-3) bioactivity. An agonist can be a compound which mimics a bioactivity of a Dkk or Dkk-related protein, such as transduction of a signal from a Dkk receptor, by, *e.g.*, interacting with a hDkk-3 receptor. An agonist can also be a compound that upregulates
20 expression of a Dkk or Dkk-related gene. An agonist can also be a compound which modulates the expression or activity of a protein which is located downstream, for example, of a Dkk receptor, thereby mimicking or enhancing the effect of binding of Dkk to a Dkk receptor.

"Antagonist" as used herein is meant to refer to an agent that inhibits, decreases
25 or suppresses a bioactivity (*e.g.*, hDkk-3). An antagonist can be a compound which decreases signalling from a Dkk or Dkk-related protein, *e.g.*, a compound that is capable of binding to hDkk-3 or to a hDkk-3 receptor. A preferred antagonist inhibits the interaction between a Dkk or Dkk-related protein and another molecule, such as a Dkk receptor. Alternatively, an antagonist can be a compound that downregulates expression
30 of a Dkk or Dkk-related gene. An antagonist can also be a compound which modulates

the expression or activity of a protein which is located downstream of a Dkk receptor, thereby antagonizing the effect of binding of Dkk to a Dkk receptor.

In one embodiment, the invention provides assays for screening candidate or test compounds which bind to or modulate the activity of a Dkk or Dkk-related protein or polypeptide or biologically active portion thereof. In another embodiment, the invention provides assays for screening candidate or test compounds which bind to or modulate the activity of a Dkk receptor. The test compounds of the present invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the 'one-bead one-compound' library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds (Lam, K.S. (1997) *Anticancer Drug Des.* 12:145).

Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt *et al.* (1993) *Proc. Natl. Acad. Sci. U.S.A.* 90:6909; Erb *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:11422; Zuckermann *et al.* (1994). *J. Med. Chem.* 37:2678; Cho *et al.*, (1993) *Science* 261:1303; Carrell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2059; Carrell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2061; and in Gallop *et al.* (1994) *J. Med. Chem.* 37:1233.

Libraries of compounds may be presented in solution (*e.g.*, Houghten (1992) *Biotechniques* 13:412-421), or on beads (Lam (1991) *Nature* 354:82-84), chips (Fodor (1993) *Nature* 364:555-556), bacteria (Ladner USP 5,223,409), spores (Ladner USP '409), plasmids (Cull *et al.* (1992) *Proc Natl Acad Sci USA* 89:1865-1869) or on phage (Scott and Smith (1990) *Science* 249:386-390); (Devlin (1990) *Science* 249:404-406); (Cwirla *et al.* (1990) *Proc. Natl. Acad. Sci.* 87:6378-6382); (Felici (1991) *J. Mol. Biol.* 222:301-310); (Ladner *supra.*).

In one embodiment, an assay is a cell-based assay in which a cell which expresses a Dkk receptor on the cell surface is contacted with a test compound and the ability of the test compound to bind to a Dkk receptor determined. The cell, for

example, can be of mammalian origin or a yeast cell. Determining the ability of the test compound to bind to a Dkk receptor can be accomplished, for example, by coupling the test compound with a radioisotope or enzymatic label such that binding of the test compound to the Dkk receptor can be determined by detecting the labeled compound in
5 a complex. For example, test compounds can be labeled with ^{125}I , ^{35}S , ^{14}C , or ^3H , either directly or indirectly, and the radioisotope detected by direct counting of radioemmission or by scintillation counting. Alternatively, test compounds can be enzymatically labeled with, for example, horseradish peroxidase, alkaline phosphatase, or luciferase, and the enzymatic label detected by determination of conversion of an
10 appropriate substrate to product.

It is also within the scope of this invention to determine the ability of a test compound to interact with a Dkk receptor without the labeling of any of the interactants. For example, a microphysiometer can be used to detect the interaction of a test compound with a Dkk receptor without the labeling of either the test compound or the
15 receptor. McConnell, H. M. *et al.* (1992) *Science* 257:1906-1912. As used herein, a "microphysiometer" (*e.g.*, Cytosensor™) is an analytical instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indicator of the interaction between ligand and receptor.

20 In a preferred embodiment, the assay comprises contacting a cell which expresses a Dkk receptor on the cell surface with a Dkk protein or biologically-active portion thereof, to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with a Dkk receptor, wherein determining the ability of the test compound to interact with a Dkk
25 receptor comprises determining the ability of the test compound to preferentially bind to the Dkk receptor as compared to the ability of Dkk, or a biologically active portion thereof, to bind to the receptor.

In another embodiment, an assay is a cell-based assay comprising contacting a cell expressing a Dkk target molecule with a test compound and determining the ability
30 of the test compound to modulate (*e.g.* stimulate or inhibit) the activity of the Dkk target molecule. Determining the ability of the test compound to modulate the activity of a

Dkk target molecule can be accomplished, for example, by determining the ability of the Dkk protein to bind to or interact with the Dkk target molecule.

Determining the ability of the Dkk protein to bind to or interact with a Dkk target molecule can be accomplished by one of the methods described above for determining
5 direct binding. In a preferred embodiment, determining the ability of the Dkk protein to bind to or interact with a Dkk target molecule can be accomplished by determining the activity of the target molecule. For example, the activity of the target molecule can be determined by detecting induction of a cellular second messenger of the target (*i.e.* intracellular Ca^{2+} , diacylglycerol, IP_3 , etc.), detecting catalytic/enzymatic activity of
10 the target an appropriate substrate, detecting the induction of a reporter gene (comprising a Dkk-responsive regulatory element operatively linked to a nucleic acid encoding a detectable marker, *e.g.*, luciferase), or detecting a cellular response, for example, development, differentiation or rate of proliferation.

In yet another embodiment, an assay of the present invention is a cell-free assay
15 in which a Dkk or Dkk-related protein or biologically active portion thereof is contacted with a test compound and the ability of the test compound to bind to the Dkk or Dkk-related protein or biologically active portion thereof is determined. Binding of the test compound to the Dkk or Dkk-related protein can be determined either directly or indirectly as described above. In a preferred embodiment, the assay includes contacting
20 the Dkk or Dkk-related protein or biologically active portion thereof with a known compound which binds Dkk or the Dkk-related protein to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with a Dkk or Dkk-related protein, wherein determining the ability of the test compound to interact with a Dkk or Dkk-related protein comprises
25 determining the ability of the test compound to preferentially bind to Dkk or a Dkk-related protein or biologically active portion thereof as compared to the known compound.

In another embodiment, the assay is a cell-free assay in which a Dkk or Dkk-related protein or biologically active portion thereof is contacted with a test compound
30 and the ability of the test compound to modulate (*e.g.*, stimulate or inhibit) the activity of the Dkk or Dkk-related protein or biologically active portion thereof is determined.

Determining the ability of the test compound to modulate the activity of a Dkk or Dkk-related protein can be accomplished, for example, by determining the ability of the Dkk or Dkk-related protein to bind to a target molecule (*e.g.*, a Dkk-target molecule) by one of the methods described above for determining direct binding. Determining the ability
5 of the Dkk or Dkk-related protein to bind to a target molecule can also be accomplished using a technology such as real-time Biomolecular Interaction Analysis (BIA).

Sjolander, S. and Urbaniczky, C. (1991) *Anal. Chem.* 63:2338-2345 and Szabo *et al.* (1995) *Curr. Opin. Struct. Biol.* 5:699-705. As used herein, "BIA" is a technology for studying biospecific interactions in real time, without labeling any of the interactants
10 (*e.g.*, BIAcore™). Changes in the optical phenomenon surface plasmon resonance (SPR) can be used as an indication of real-time reactions between biological molecules.

In an alternative embodiment, determining the ability of the test compound to modulate the activity of a Dkk or Dkk-related protein can be accomplished by determining the ability of the Dkk or Dkk-related protein to further modulate the activity
15 of a target molecule (*e.g.*, a Dkk-target molecule). For example, the catalytic/enzymatic activity of the target molecule on an appropriate substrate can be determined as previously described.

In yet another embodiment, the cell-free assay involves contacting a Dkk or Dkk-related protein or biologically active portion thereof with a known compound which
20 binds the Dkk or Dkk-related protein to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with the Dkk or Dkk-related protein, wherein determining the ability of the test compound to interact with the Dkk or Dkk-related protein comprises determining the ability of the Dkk or Dkk-related protein to preferentially bind to or modulate the
25 activity of a target molecule (*e.g.*, a Dkk target molecule).

In many drug screening programs which test libraries of compounds and natural extracts, high throughput assays are desirable in order to maximize the number of compounds surveyed in a given period of time. Assays which are performed in cell-free systems, such as may be derived with purified or semi-purified proteins, are often
30 preferred as "primary" screens in that they can be generated to permit rapid development and relatively easy detection of an alteration in a molecular target which is mediated by

a test compound. Moreover, the effects of cellular toxicity and/or bioavailability of the test compound can be generally ignored in the *in vitro* system, the assay instead being focused primarily on the effect of the drug on the molecular target as may be manifest in an alteration of binding affinity with upstream or downstream elements. Accordingly, in an exemplary screening assay of the present invention, the compound of interest is contacted with a Dkk (*e.g.*, hDkk-3) protein or a Dkk (*e.g.*, hDkk-3) binding partner, *e.g.*, a receptor. The receptor can be soluble or the receptor can be present on a cell surface. To the mixture of the compound and the Dkk protein or Dkk binding partner is then added a composition containing a Dkk binding partner or a Dkk protein, respectively. Detection and quantification of complexes of Dkk proteins and Dkk binding partners provide a means for determining a compound's efficacy at inhibiting (or potentiating) complex formation between Dkk and a binding partner. The efficacy of the compound can be assessed by generating dose response curves from data obtained using various concentrations of the test compound. Moreover, a control assay can also be performed to provide a baseline for comparison. In the control assay, isolated and purified Dkk polypeptide or binding partner is added to a composition containing the Dkk binding partner or Dkk polypeptide, and the formation of a complex is quantitated in the absence of the test compound.

The cell-free assays of the present invention are amenable to use of both soluble and/or membrane-bound forms of isolated proteins (*e.g.* Dkk proteins or biologically active portions thereof or Dkk target molecules). In the case of cell-free assays in which a membrane-bound form an isolated protein is used (*e.g.*, a Dkk target molecule or receptor) it may be desirable to utilize a solubilizing agent such that the membrane-bound form of the isolated protein is maintained in solution. Examples of such solubilizing agents include non-ionic detergents such as n-octylglucoside, n-dodecylglucoside, n-dodecylmaltoside, octanoyl-N-methylglucamide, decanoyl-N-methylglucamide, Triton[®] X-100, Triton[®] X-114, Thesit[®], Isotridecypoly(ethylene glycol ether)_n, 3-[(3-cholamidopropyl)dimethylamminio]-1-propane sulfonate (CHAPS), 3-[(3-cholamidopropyl)dimethylamminio]-2-hydroxy-1-propane sulfonate (CHAPSO), or N-dodecyl=N,N-dimethyl-3-ammonio-1-propane sulfonate.

In more than one embodiment of the above assay methods of the present invention, it may be desirable to immobilize either Dkk, a Dkk-related protein or a target molecule to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test compound to a Dkk or Dkk-related protein, or interaction of a Dkk or Dkk-related protein with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtitre plates, test tubes, and micro-centrifuge tubes. In one embodiment, a fusion protein can be provided which adds a domain that allows one or both of the proteins to be bound to a matrix. For example, glutathione-S-transferase/Dkk fusion proteins or glutathione-S-transferase/target fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtitre plates, which are then combined with the test compound or the test compound and either the non-adsorbed target protein or Dkk protein, and the mixture incubated under conditions conducive to complex formation (*e.g.*, at physiological conditions for salt and pH). Following incubation, the beads or microtitre plate wells are washed to remove any unbound components, the matrix immobilized in the case of beads, complex determined either directly or indirectly, for example, as described above. Alternatively, the complexes can be dissociated from the matrix, and the level of Dkk binding or activity determined using standard techniques.

Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either a Dkk protein, Dkk-related protein, or a Dkk target molecule can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated protein or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well known in the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, IL), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies reactive with Dkk, Dkk-related protein, or target molecules but which do not interfere with binding of the protein to its target molecule can be derivatized to the wells of the plate, and unbound target, Dkk, or Dkk-related protein trapped in the wells by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized

complexes, include immunodetection of complexes using antibodies reactive with the Dkk or Dkk-related protein or target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the Dkk or Dkk-related protein or target molecule.

5 In another embodiment, modulators of Dkk or Dkk-related expression are identified in a method wherein a cell is contacted with a candidate compound and the expression of Dkk or Dkk-related mRNA or protein in the cell is determined. The level of expression of mRNA or protein in the presence of the candidate compound is compared to the level of expression of mRNA or protein in the absence of the candidate
10 compound. The candidate compound can then be identified as a modulator of Dkk or Dkk-related expression based on this comparison. For example, when expression of Dkk mRNA or protein is greater (statistically significantly greater) in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of Dkk mRNA or protein expression. Alternatively, when expression of Dkk
15 mRNA or protein is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of Dkk mRNA or protein expression. The level of Dkk or Dkk-related mRNA or protein expression in the cells can be determined by methods described herein for detecting Dkk mRNA or protein.

20 In yet another aspect of the invention, the Dkk or Dkk-related proteins can be used as "bait proteins" in a two-hybrid assay or three-hybrid assay (see, *e.g.*, U.S. Patent No. 5,283,317; Zervos *et al.* (1993) *Cell* 72:223-232; Madura *et al.* (1993) *J. Biol. Chem.* 268:12046-12054; Bartel *et al.*, (1993) *Biotechniques* 14:920-924; Iwabuchi *et al.* (1993) *Oncogene* 8:1693-1696; and Brent WO94/10300), to identify other proteins,
25 which bind to or interact with Dkk or Dkk-related proteins ("binding proteins" or "bp") and modulate Dkk or Dkk-related activity. Such binding proteins are also likely to be involved in the propagation of signals by the Dkk or Dkk-related proteins as, for example, downstream elements of a Dkk-mediated signaling pathway. Alternatively, such binding proteins are likely to be cell-surface molecules associated with non-Dkk
30 expressing cells, wherein such binding proteins are involved in signal transduction.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. In one construct, the gene that codes for a Dkk protein is fused to a gene encoding the DNA binding domain of a known transcription factor (*e.g.*, GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact, *in vivo*, forming a Dkk-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (*e.g.*, LacZ) which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene which encodes the protein which interacts with the Dkk or Dkk-related protein.

This invention further pertains to novel agents identified by the above-described screening assays and to processes for producing such agents by use of these assays. Accordingly, in one embodiment, the present invention includes a compound or agent obtainable by a method comprising the steps of any one of the aforementioned screening assays (*e.g.*, cell-based assays or cell-free assays). For example, in one embodiment, the invention includes a compound or agent obtainable by a method comprising contacting a cell which expresses a target molecule with a test compound and the determining the ability of the test compound to bind to, or modulate the activity of, the target molecule. In another embodiment, the invention includes a compound or agent obtainable by a method comprising contacting a cell which expresses a target molecule with a Dkk or Dkk-related protein or biologically-active portion thereof, to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with, or modulate the activity of, the target molecule. In another embodiment, the invention includes a compound or agent obtainable by a method comprising contacting a Dkk or Dkk-related protein or biologically active portion thereof with a test compound and determining the ability of the test compound to

bind to, or modulate (*e.g.*, stimulate or inhibit) the activity of, the Dkk or Dkk-related protein or biologically active portion thereof. In yet another embodiment, the present invention includes a compound or agent obtainable by a method comprising contacting a Dkk or Dkk-related protein or biologically active portion thereof with a known

- 5 compound which binds the Dkk or Dkk-related protein to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with, or modulate the activity of the Dkk or Dkk-related protein.

- Accordingly, it is within the scope of this invention to further use an agent
10 identified as described herein in an appropriate animal model. For example, an agent identified as described herein (*e.g.*, a Dkk modulating agent, an antisense Dkk nucleic acid molecule, a Dkk-specific antibody, or a Dkk-binding partner) can be used in an animal model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal
15 model to determine the mechanism of action of such an agent.

- The present invention also pertains to uses of novel agents identified by the above-described screening assays for diagnoses, prognoses, and treatments as described herein. Accordingly, it is within the scope of the present invention to use such agents in the design, formulation, synthesis, manufacture, and/or production of a drug or
20 pharmaceutical composition for use in diagnosis, prognosis, or treatment, as described herein. For example, in one embodiment, the present invention includes a method of synthesizing or producing a drug or pharmaceutical composition by reference to the structure and/or properties of a compound obtainable by one of the above-described screening assays. For example, a drug or pharmaceutical composition can be
25 synthesized based on the structure and/or properties of a compound obtained by a method in which a cell which expresses a target molecule (*e.g.*, a Dkk target molecule) is contacted with a test compound and the ability of the test compound to bind to, or modulate the activity of, the target molecule is determined. In another exemplary embodiment, the present invention includes a method of synthesizing or producing a
30 drug or pharmaceutical composition based on the structure and/or properties of a compound obtainable by a method in which a Dkk or Dkk-related protein or biologically

active portion thereof is contacted with a test compound and the ability of the test compound to bind to, or modulate (*e.g.*, stimulate or inhibit) the activity of, the Dkk or Dkk-related protein or biologically active portion thereof is determined.

5 B. Detection Assays

Portions or fragments of the cDNA sequences identified herein (and the corresponding complete gene sequences) can be used in numerous ways as polynucleotide reagents. For example, these sequences can be used to: (i) map their respective genes on a chromosome; and, thus, locate gene regions associated with
10 genetic disease; (ii) identify an individual from a minute biological sample (tissue typing); and (iii) aid in forensic identification of a biological sample. These applications are described in the subsections below.

1. Chromosome Mapping

15 Once the sequence (or a portion of the sequence) of a gene has been isolated, this sequence can be used to map the location of the gene on a chromosome. This process is called chromosome mapping. Accordingly, portions or fragments of the Dkk or Dkk-related nucleotide sequences, described herein, can be used to map the location of the Dkk or Dkk-related genes on a chromosome. The mapping of the Dkk or Dkk-related
20 sequences to chromosomes is an important first step in correlating these sequences with genes associated with disease.

Briefly, Dkk or Dkk-related genes can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp in length) from the Dkk or Dkk-related nucleotide sequences. Computer analysis of the Dkk or Dkk-related sequences can be used to
25 predict primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers can then be used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the Dkk or Dkk-related sequences will yield an amplified fragment.

Somatic cell hybrids are prepared by fusing somatic cells from different mammals (*e.g.*, human and mouse cells). As hybrids of human and mouse cells grow and divide, they gradually lose human chromosomes in random order, but retain the mouse chromosomes. By using media in which mouse cells cannot grow, because they
5 lack a particular enzyme, but human cells can, the one human chromosome that contains the gene encoding the needed enzyme, will be retained. By using various media, panels of hybrid cell lines can be established. Each cell line in a panel contains either a single human chromosome or a small number of human chromosomes, and a full set of mouse chromosomes, allowing easy mapping of individual genes to specific human
10 chromosomes. (D'Eustachio P. *et al.*, (1983) *Science* 220:919-924). Somatic cell hybrids containing only fragments of human chromosomes can also be produced by using human chromosomes with translocations and deletions.

PCR mapping of somatic cell hybrids is a rapid procedure for assigning a particular sequence to a particular chromosome. Three or more sequences can be
15 assigned per day using a single thermal cycler. Using the Dkk or Dkk-related nucleotide sequences to design oligonucleotide primers, sublocalization can be achieved with panels of fragments from specific chromosomes. Other mapping strategies which can similarly be used to map a 9o, 1p, or 1v sequence to its chromosome include *in situ* hybridization (described in Fan, Y. *et al.*, (1990) *PNAS*, 87:6223-27), pre-screening with
20 labeled flow-sorted chromosomes, and pre-selection by hybridization to chromosome specific cDNA libraries.

Fluorescence *in situ* hybridization (FISH) of a DNA sequence to a metaphase chromosomal spread can further be used to provide a precise chromosomal location in one step. Chromosome spreads can be made using cells whose division has been
25 blocked in metaphase by a chemical such as colcemid that disrupts the mitotic spindle. The chromosomes can be treated briefly with trypsin, and then stained with Giemsa. A pattern of light and dark bands develops on each chromosome, so that the chromosomes can be identified individually. The FISH technique can be used with a DNA sequence as short as 500 or 600 bases. However, clones larger than 1,000 bases have a higher
30 likelihood of binding to a unique chromosomal location with sufficient signal intensity for simple detection. Preferably 1,000 bases, and more preferably 2,000 bases will

suffice to get good results at a reasonable amount of time. For a review of this technique, see Verma *et al.*, Human Chromosomes: A Manual of Basic Techniques (Pergamon Press, New York 1988).

Reagents for chromosome mapping can be used individually to mark a single
5 chromosome or a single site on that chromosome, or panels of reagents can be used for marking multiple sites and/or multiple chromosomes. Reagents corresponding to noncoding regions of the genes actually are preferred for mapping purposes. Coding sequences are more likely to be conserved within gene families, thus increasing the chance of cross hybridizations during chromosomal mapping.

10 Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. (Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man, available on-line through Johns Hopkins University Welch Medical Library). The relationship between a gene and a disease, mapped to the same chromosomal region, can
15 then be identified through linkage analysis (co-inheritance of physically adjacent genes), described in, for example, Egeland, J. *et al.* (1987) *Nature*, 325:783-787.

Moreover, differences in the DNA sequences between individuals affected and unaffected with a disease associated with a Dkk or Dkk-related gene, can be determined. If a mutation is observed in some or all of the affected individuals but not in any
20 unaffected individuals, then the mutation is likely to be the causative agent of the particular disease. Comparison of affected and unaffected individuals generally involves first looking for structural alterations in the chromosomes, such as deletions or translocations that are visible from chromosome spreads or detectable using PCR based on that DNA sequence. Ultimately, complete sequencing of genes from several
25 individuals can be performed to confirm the presence of a mutation and to distinguish mutations from polymorphisms.

2. Tissue Typing

The Dkk or Dkk-related sequences of the present invention can also be used to
30 identify individuals from minute biological samples. The United States military, for example, is considering the use of restriction fragment length polymorphism (RFLP) for

identification of its personnel. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identification. This method does not suffer from the current limitations of "Dog Tags" which can be lost, switched, or stolen, making positive identification
5 difficult. The sequences of the present invention are useful as additional DNA markers for RFLP (described in U.S. Patent 5,272,057).

Furthermore, the sequences of the present invention can be used to provide an alternative technique which determines the actual base-by-base DNA sequence of selected portions of an individual's genome. Thus, the Dkk or Dkk-related nucleotide
10 sequences described herein can be used to prepare two PCR primers from the 5' and 3' ends of the sequences. These primers can then be used to amplify an individual's DNA and subsequently sequence it.

Panels of corresponding DNA sequences from individuals, prepared in this manner, can provide unique individual identifications, as each individual will have a
15 unique set of such DNA sequences due to allelic differences. The sequences of the present invention can be used to obtain such identification sequences from individuals and from tissue. The Dkk or Dkk-related nucleotide sequences of the invention uniquely represent portions of the human genome. Allelic variation occurs to some degree in the coding regions of these sequences, and to a greater degree in the noncoding regions. It
20 is estimated that allelic variation between individual humans occurs with a frequency of about once per each 500 bases. Each of the sequences described herein can, to some degree, be used as a standard against which DNA from an individual can be compared for identification purposes. Because greater numbers of polymorphisms occur in the noncoding regions, fewer sequences are necessary to differentiate individuals. The
25 noncoding sequences of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20, can comfortably provide positive individual identification with a panel of perhaps 10 to 1,000 primers which each yield a noncoding amplified sequence of 100 bases. If predicted coding sequences, such as those in SEQ ID NO:3, SEQ ID NO:6, SEQ ID NO:9, SEQ ID NO:15 or SEQ ID NO:22 are used, a more appropriate
30 number of primers for positive individual identification would be 500-2,000.

If a panel of reagents from Dkk or Dkk-related nucleotide sequences described herein is used to generate a unique identification database for an individual, those same reagents can later be used to identify tissue from that individual. Using the unique identification database, positive identification of the individual, living or dead, can be made from extremely small tissue samples.

3. Use of Partial Dkk or Dkk-related Sequences in Forensic Biology

DNA-based identification techniques can also be used in forensic biology.

Forensic biology is a scientific field employing genetic typing of biological evidence found at a crime scene as a means for positively identifying, for example, a perpetrator of a crime. To make such an identification, PCR technology can be used to amplify DNA sequences taken from very small biological samples such as tissues, *e.g.*, hair or skin, or body fluids, *e.g.*, blood, saliva, or semen found at a crime scene. The amplified sequence can then be compared to a standard, thereby allowing identification of the origin of the biological sample.

The sequences of the present invention can be used to provide polynucleotide reagents, *e.g.*, PCR primers, targeted to specific loci in the human genome, which can enhance the reliability of DNA-based forensic identifications by, for example, providing another "identification marker" (*i.e.* another DNA sequence that is unique to a particular individual). As mentioned above, actual base sequence information can be used for identification as an accurate alternative to patterns formed by restriction enzyme generated fragments. Sequences targeted to noncoding regions of SEQ ID NOs:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20 are particularly appropriate for this use as greater numbers of polymorphisms occur in the noncoding regions, making it easier to differentiate individuals using this technique. Examples of polynucleotide reagents include the Dkk nucleotide sequences or portions thereof, *e.g.*, fragments derived from the noncoding regions of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20, having a length of at least 20 bases, preferably at least 30 bases.

The Dkk or Dkk-related nucleotide sequences described herein can further be used to provide polynucleotide reagents, *e.g.*, labeled or labelable probes which can be used in, for example, an *in situ* hybridization technique, to identify a specific tissue, *e.g.*, brain tissue. This can be very useful in cases where a forensic pathologist is presented
5 with a tissue of unknown origin. Panels of such Dkk or Dkk-related probes can be used to identify tissue by species and/or by organ type.

In a similar fashion, these reagents, *e.g.*, Dkk or Dkk-related primers or probes can be used to screen tissue culture for contamination (*i.e.* screen for the presence of a mixture of different types of cells in a culture).

10

C. Predictive Medicine:

The present invention also pertains to the field of predictive medicine in which diagnostic assays, prognostic assays, and monitoring clinical trials are used for prognostic (predictive) purposes to thereby treat an individual prophylactically.

15 Accordingly, one aspect of the present invention relates to diagnostic assays for determining Dkk or Dkk-related protein and/or nucleic acid expression as well as Dkk or Dkk-related activity, in the context of a biological sample (*e.g.*, blood, serum, cells, tissue) to thereby determine whether an individual is afflicted with a disease or disorder, or is at risk of developing a disorder, associated with aberrant Dkk or Dkk-related
20 expression or activity, such as aberrant cell proliferation, differentiation, and/or survival resulting for example in a neurodegenerative disease (*e.g.*, Alzheimer's disease, Parkinson's disease, Huntington's chorea, amyotrophic lateral sclerosis and the like, as well as spinocerebellar degenerations) or cancer (for example, cancers of the epithelia (*e.g.*, carcinomas of the pancreas, stomach, liver, secretory glands (*e.g.*,
25 adenocarcinoma) bladder, lung, breast, skin (*e.g.*, malignant melanoma), reproductive tract including prostate gland, ovary, cervix and uterus); cancers of the hematopoietic and immune system (*e.g.*, leukemias and lymphomas); cancers of the central nervous, brain system and eye (*e.g.*, gliomas, glioblastoma, neuroblastoma and retinoblastoma); and cancers of connective tissues, bone, muscles and vasculature (*e.g.*, sarcomas)). The
30 invention also provides for prognostic (or predictive) assays for determining whether an individual is at risk of developing a disorder associated with Dkk or Dkk-related protein,

nucleic acid expression or activity. For example, mutations in a Dkk or Dkk-related gene can be assayed in a biological sample. Such assays can be used for prognostic or predictive purpose to thereby prophylactically treat an individual prior to the onset of a disorder characterized by or associated with Dkk or Dkk-related protein, nucleic acid expression or activity.

Another aspect of the invention pertains to monitoring the influence of agents (*e.g.*, drugs, compounds) on the expression or activity of Dkk or Dkk-related in clinical trials.

These and other agents are described in further detail in the following sections.

1. Diagnostic Assays

An exemplary method for detecting the presence or absence of Dkk or Dkk-related protein or nucleic acid in a biological sample involves obtaining a biological sample from a test subject and contacting the biological sample with a compound or an agent capable of detecting Dkk or Dkk-related protein or nucleic acid (*e.g.*, mRNA, genomic DNA) that encodes Dkk or Dkk-related protein such that the presence of Dkk or Dkk-related protein or nucleic acid is detected in the biological sample. A preferred agent for detecting Dkk or Dkk-related mRNA or genomic DNA is a labeled nucleic acid probe capable of hybridizing to Dkk or Dkk-related mRNA or genomic DNA. The nucleic acid probe can be, for example, a full-length Dkk nucleic acid, such as the nucleic acid of SEQ ID NO: 1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a portion thereof, such as an oligonucleotide of at least 15, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically hybridize under stringent conditions to Dkk or Dkk-related mRNA or genomic DNA. Other suitable probes for use in the diagnostic assays of the invention are described herein.

A preferred agent for detecting Dkk or Dkk-related protein is an antibody capable of binding to the protein, preferably an antibody with a detectable label.

Antibodies can be polyclonal, or more preferably, monoclonal. An intact antibody, or a fragment thereof (*e.g.*, Fab or F(ab')₂) can be used. The term "labeled", with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (*i.e.*, physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include detection of a primary antibody using a fluorescently labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently labeled streptavidin. The term "biological sample" is intended to include tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject. That is, the detection method of the invention can be used to detect Dkk or Dkk-related mRNA, protein, or genomic DNA in a biological sample *in vitro* as well as *in vivo*. For example, *in vitro* techniques for detection of Dkk or Dkk-related mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro* techniques for detection of Dkk or Dkk-related protein include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence. *In vitro* techniques for detection of Dkk or Dkk-related genomic DNA include Southern hybridizations. Furthermore, *in vivo* techniques for detection of Dkk or Dkk-related protein include introducing into a subject a labeled antibody. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques.

In one embodiment, the biological sample contains protein molecules from the test subject. Alternatively, the biological sample can contain mRNA molecules from the test subject or genomic DNA molecules from the test subject. A preferred biological sample is a serum sample isolated by conventional means from a subject.

In another embodiment, the methods further involve obtaining a control biological sample from a control subject, contacting the control sample with a compound or agent capable of detecting Dkk or Dkk-related protein, mRNA, or genomic DNA, such that the presence of Dkk or Dkk-related protein, mRNA or genomic DNA is

detected in the biological sample, and comparing the presence of Dkk or Dkk-related protein, mRNA or genomic DNA in the control sample with the presence of Dkk or Dkk-related protein, mRNA or genomic DNA in the test sample.

The invention also encompasses kits for detecting the presence of Dkk or a Dkk-related protein in a biological sample. For example, the kit can comprise a labeled compound or agent capable of detecting Dkk or Dkk-related protein or mRNA in a biological sample; means for determining the amount of Dkk or Dkk-related protein or mRNA in the sample; and means for comparing the amount of Dkk or Dkk-related protein or mRNA in the sample with a standard. The compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect Dkk or Dkk-related protein or nucleic acid.

2. Prognostic Assays

The diagnostic methods described herein can furthermore be utilized to identify subjects having or at risk of developing a disease or disorder associated with aberrant Dkk expression or activity. For example, the assays described herein, such as the preceding diagnostic assays or the following assays, can be utilized to identify a subject having or at risk of developing a disorder associated with Dkk or Dkk-related protein, nucleic acid expression or activity such as a proliferative disorder, a differentiative or developmental disorder, a hematopoietic disorder as well as diseases, conditions or disorders characterized by abnormal cell survival, abnormal extracellular structure, or an abnormality in a defense mechanism. Alternatively, the prognostic assays can be utilized to identify a subject having or at risk for developing a differentiative or proliferative disease (*e.g.*, cancer). Thus, the present invention provides a method for identifying a disease or disorder associated with aberrant Dkk or Dkk-related expression or activity in which a test sample is obtained from a subject and Dkk or Dkk-related protein or nucleic acid (*e.g.*, mRNA, genomic DNA) is detected, wherein the presence of Dkk or Dkk-related protein or nucleic acid is diagnostic for a subject having or at risk of developing a disease or disorder associated with aberrant Dkk or Dkk-related expression or activity. As used herein, a "test sample" refers to a biological sample obtained from a

subject of interest. For example, a test sample can be a biological fluid (*e.g.*, serum), cell sample, or tissue.

Furthermore, the prognostic assays described herein can be used to determine whether a subject can be administered an agent (*e.g.*, an agonist, antagonist, peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate) to treat a disease or disorder associated with aberrant Dkk or Dkk-related expression or activity. For example, such methods can be used to determine whether a subject can be effectively treated with an agent for a disorder, such as a proliferative disorder, a differentiative or developmental disorder, a hematopoietic disorder, as well disorders characterized by abnormal cell survival, an abnormal extracellular structure, or an abnormality in a defense mechanism. Alternatively, such methods can be used to determine whether a subject can be effectively treated with an agent for a differentiative or proliferative disease (*e.g.*, cancer). Thus, the present invention provides methods for determining whether a subject can be effectively treated with an agent for a disorder associated with aberrant Dkk or Dkk-related expression or activity in which a test sample is obtained and Dkk or Dkk-related protein or nucleic acid expression or activity is detected (*e.g.*, wherein the abundance of Dkk or Dkk-related protein or nucleic acid expression or activity is diagnostic for a subject that can be administered the agent to treat a disorder associated with aberrant Dkk or Dkk-related expression or activity.)

The methods of the invention can also be used to detect genetic alterations in a Dkk or Dkk-related gene, thereby determining if a subject with the altered gene is at risk for a disorder characterized by aberrant development, aberrant cellular differentiation, aberrant cellular proliferation or an aberrant hematopoietic response. In preferred embodiments, the methods include detecting, in a sample of cells from the subject, the presence or absence of a genetic alteration characterized by at least one of an alteration affecting the integrity of a gene encoding a Dkk or Dkk-related-protein, or the mis-expression of the Dkk or Dkk-related gene. For example, such genetic alterations can be detected by ascertaining the existence of at least one of 1) a deletion of one or more nucleotides from a Dkk or Dkk-related gene; 2) an addition of one or more nucleotides to a Dkk or Dkk-related gene; 3) a substitution of one or more nucleotides of a Dkk or Dkk-related gene, 4) a chromosomal rearrangement of a Dkk or Dkk-related gene; 5) an

alteration in the level of a messenger RNA transcript of a Dkk or Dkk-related gene, 6) aberrant modification of a Dkk or Dkk-related gene, such as of the methylation pattern of the genomic DNA, 7) the presence of a non-wild type splicing pattern of a messenger RNA transcript of a Dkk or Dkk-related gene, 8) a non-wild type level of a Dkk or Dkk-related-protein, 9) allelic loss of a Dkk or Dkk-related gene, and 10) inappropriate post-translational modification of a Dkk or Dkk-related-protein. As described herein, there are a large number of assay techniques known in the art which can be used for detecting alterations in a Dkk or Dkk-related gene. A preferred biological sample is a tissue or serum sample isolated by conventional means from a subject.

10 In certain embodiments, detection of the alteration involves the use of a probe/primer in a polymerase chain reaction (PCR) (see, *e.g.*, U.S. Patent Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (see, *e.g.*, Landegran *et al.* (1988) *Science* 241:1077-1080; and Nakazawa *et al.* (1994) *PNAS* 91:360-364), the latter of which can be particularly
15 useful for detecting point mutations in the Dkk or Dkk-related-gene (see Abravaya *et al.* (1995) *Nucleic Acids Res.* 23:675-682). This method can include the steps of collecting a sample of cells from a patient, isolating nucleic acid (*e.g.*, genomic, mRNA or both) from the cells of the sample, contacting the nucleic acid sample with one or more primers which specifically hybridize to a Dkk or Dkk-related gene under conditions such
20 that hybridization and amplification of the Dkk or Dkk-related-gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. It is anticipated that PCR and/or LCR may be desirable to use as a preliminary amplification step in conjunction with any of the techniques used for detecting mutations described
25 herein.

Alternative amplification methods include: self sustained sequence replication (Guatelli, J.C. *et al.*, 1990, *Proc. Natl. Acad. Sci. USA* 87:1874-1878), transcriptional amplification system (Kwoh, D.Y. *et al.*, 1989, *Proc. Natl. Acad. Sci. USA* 86:1173-1177), Q-Beta Replicase (Lizardi, P.M. *et al.*, 1988, *Bio/Technology* 6:1197), or any
30 other nucleic acid amplification method, followed by the detection of the amplified molecules using techniques well known to those of skill in the art. These detection

schemes are especially useful for the detection of nucleic acid molecules if such molecules are present in very low numbers.

In an alternative embodiment, mutations in a Dkk or Dkk-related gene from a sample cell can be identified by alterations in restriction enzyme cleavage patterns. For example, sample and control DNA is isolated, amplified (optionally), digested with one or more restriction endonucleases, and fragment length sizes are determined by gel electrophoresis and compared. Differences in fragment length sizes between sample and control DNA indicates mutations in the sample DNA. Moreover, the use of sequence specific ribozymes (see, for example, U.S. Patent No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site.

In other embodiments, genetic mutations in a Dkk or Dkk-related gene can be identified by hybridizing a sample and control nucleic acids, *e.g.*, DNA or RNA, to high density arrays containing hundreds or thousands of oligonucleotides probes (Cronin, M.T. *et al.* (1996) *Human Mutation* 7: 244-255; Kozal, M.J. *et al.* (1996) *Nature Medicine* 2: 753-759). For example, genetic mutations in Dkk can be identified in two dimensional arrays containing light-generated DNA probes as described in Cronin, M.T. *et al. supra*. Briefly, a first hybridization array of probes can be used to scan through long stretches of DNA in a sample and control to identify base changes between the sequences by making linear arrays of sequential overlapping probes. This step allows the identification of point mutations. This step is followed by a second hybridization array that allows the characterization of specific mutations by using smaller, specialized probe arrays complementary to all variants or mutations detected. Each mutation array is composed of parallel probe sets, one complementary to the wild-type gene and the other complementary to the mutant gene.

In yet another embodiment, any of a variety of sequencing reactions known in the art can be used to directly sequence the Dkk or Dkk-related gene and detect mutations by comparing the sequence of the sample Dkk or Dkk-related sequence with the corresponding wild-type (control) sequence. Examples of sequencing reactions include those based on techniques developed by Maxim and Gilbert ((1977) *PNAS* 74:560) or Sanger ((1977) *PNAS* 74:5463). It is also contemplated that any of a variety

of automated sequencing procedures can be utilized when performing the diagnostic assays ((1995) *Biotechniques* 19:448), including sequencing by mass spectrometry (see, e.g., PCT International Publication No. WO 94/16101; Cohen *et al.* (1996) *Adv. Chromatogr.* 36:127-162; and Griffin *et al.* (1993) *Appl. Biochem. Biotechnol.* 38:147-159).

Other methods for detecting mutations in the Dkk or Dkk-related gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA heteroduplexes (Myers *et al.* (1985) *Science* 230:1242). In general, the art technique of "mismatch cleavage" starts by providing heteroduplexes of formed by hybridizing (labeled) RNA or DNA containing the wild-type Dkk or Dkk-related sequence with potentially mutant RNA or DNA obtained from a tissue sample. The double-stranded duplexes are treated with an agent which cleaves single-stranded regions of the duplex such as which will exist due to basepair mismatches between the control and sample strands. For instance, RNA/DNA duplexes can be treated with RNase and DNA/DNA hybrids treated with S1 nuclease to enzymatically digesting the mismatched regions. In other embodiments, either DNA/DNA or RNA/DNA duplexes can be treated with hydroxylamine or osmium tetroxide and with piperidine in order to digest mismatched regions. After digestion of the mismatched regions, the resulting material is then separated by size on denaturing polyacrylamide gels to determine the site of mutation. See, for example, Cotton *et al.* (1988) *Proc. Natl Acad Sci USA* 85:4397; Saleeba *et al.* (1992) *Methods Enzymol.* 217:286-295. In a preferred embodiment, the control DNA or RNA can be labeled for detection.

In still another embodiment, the mismatch cleavage reaction employs one or more proteins that recognize mismatched base pairs in double-stranded DNA (so called "DNA mismatch repair" enzymes) in defined systems for detecting and mapping point mutations in Dkk cDNAs obtained from samples of cells. For example, the mutY enzyme of *E. coli* cleaves A at G/A mismatches and the thymidine DNA glycosylase from HeLa cells cleaves T at G/T mismatches (Hsu *et al.* (1994) *Carcinogenesis* 15:1657-1662). According to an exemplary embodiment, a probe based on a Dkk sequence, e.g., a wild-type Dkk sequence, is hybridized to a cDNA or other DNA product from a test cell(s). The duplex is treated with a DNA mismatch repair enzyme,

and the cleavage products, if any, can be detected from electrophoresis protocols or the like. See, for example, U.S. Patent No. 5,459,039.

In other embodiments, alterations in electrophoretic mobility will be used to identify mutations in Dkk or Dkk-related genes. For example, single strand
5 conformation polymorphism (SSCP) may be used to detect differences in electrophoretic mobility between mutant and wild type nucleic acids (Orita *et al.* (1989) *Proc Natl. Acad. Sci USA*: 86:2766, see also Cotton (1993) *Mutat Res* 285:125-144; and Hayashi (1992) *Genet Anal Tech Appl* 9:73-79). Single-stranded DNA fragments of sample and control Dkk or Dkk-related nucleic acids will be denatured and allowed to renature. The
10 secondary structure of single-stranded nucleic acids varies according to sequence, the resulting alteration in electrophoretic mobility enables the detection of even a single base change. The DNA fragments may be labeled or detected with labeled probes. The sensitivity of the assay may be enhanced by using RNA (rather than DNA), in which the secondary structure is more sensitive to a change in sequence. In a preferred
15 embodiment, the subject method utilizes heteroduplex analysis to separate double stranded heteroduplex molecules on the basis of changes in electrophoretic mobility (Keen *et al.* (1991) *Trends Genet* 7:5).

In yet another embodiment the movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing
20 gradient gel electrophoresis (DGGE) (Myers *et al.* (1985) *Nature* 313:495). When DGGE is used as the method of analysis, DNA will be modified to insure that it does not completely denature, for example by adding a GC clamp of approximately 40 bp of high-melting GC-rich DNA by PCR. In a further embodiment, a temperature gradient is used in place of a denaturing gradient to identify differences in the mobility of control
25 and sample DNA (Rosenbaum and Reissner (1987) *Biophys Chem* 265:12753).

Examples of other techniques for detecting point mutations include, but are not limited to, selective oligonucleotide hybridization, selective amplification, or selective primer extension. For example, oligonucleotide primers may be prepared in which the known mutation is placed centrally and then hybridized to target DNA under conditions
30 which permit hybridization only if a perfect match is found (Saiki *et al.* (1986) *Nature* 324:163); Saiki *et al.* (1989) *Proc. Natl Acad. Sci USA* 86:6230). Such allele specific

- 102 -

oligonucleotides are hybridized to PCR amplified target DNA or a number of different mutations when the oligonucleotides are attached to the hybridizing membrane and hybridized with labeled target DNA.

Alternatively, allele specific amplification technology which depends on selective PCR amplification may be used in conjunction with the instant invention. Oligonucleotides used as primers for specific amplification may carry the mutation of interest in the center of the molecule (so that amplification depends on differential hybridization) (Gibbs *et al.* (1989) *Nucleic Acids Res.* 17:2437-2448) or at the extreme 3' end of one primer where, under appropriate conditions, mismatch can prevent, or reduce polymerase extension (Prossner (1993) *Tibtech* 11:238). In addition it may be desirable to introduce a novel restriction site in the region of the mutation to create cleavage-based detection (Gasparini *et al.* (1992) *Mol. Cell Probes* 6:1). It is anticipated that in certain embodiments amplification may also be performed using Taq ligase for amplification (Barany (1991) *Proc. Natl. Acad. Sci USA* 88:189). In such cases, ligation will occur only if there is a perfect match at the 3' end of the 5' sequence making it possible to detect the presence of a known mutation at a specific site by looking for the presence or absence of amplification.

The methods described herein may be performed, for example, by utilizing pre-packaged diagnostic kits comprising at least one probe nucleic acid or antibody reagent described herein, which may be conveniently used, *e.g.*, in clinical settings to diagnose patients exhibiting symptoms or family history of a disease or illness involving a Dkk gene.

Furthermore, any cell type or tissue in which Dkk or a Dkk-related sequence is expressed may be utilized in the prognostic assays described herein.

3. Monitoring of Effects During Clinical Trials

Monitoring the influence of agents (*e.g.*, drugs, compounds) on the expression or activity of Dkk or Dkk-related molecule (*e.g.*, modulation of cellular signal transduction, regulation of gene transcription in a cell involved in development or differentiation, regulation of cellular proliferation) can be applied not only in basic drug screening, but also in clinical trials. For example, the effectiveness of an agent determined by a

screening assay as described herein to increase Dkk or Dkk-related gene expression, protein levels, or upregulate Dkk or Dkk-related activity, can be monitored in clinical trials of subjects exhibiting decreased Dkk or Dkk-related gene expression, protein levels, or downregulated Dkk or Dkk-related activity. Alternatively, the effectiveness of an agent determined by a screening assay to decrease Dkk or Dkk-related gene expression, protein levels, or downregulate Dkk or Dkk-related activity, can be monitored in clinical trials of subjects exhibiting increased Dkk or Dkk-related gene expression, protein levels, or upregulated Dkk or Dkk-related activity. In such clinical trials, the expression or activity of Dkk or Dkk-related and, preferably, other genes that have been implicated in, for example, a proliferative disorder can be used as a "read out" or markers of the phenotype of a particular cell.

For example, and not by way of limitation, genes, including Dkk and Dkk-related genes, that are modulated in cells by treatment with an agent (*e.g.*, compound, drug or small molecule) which modulates Dkk or Dkk-related activity (*e.g.*, identified in a screening assay as described herein) can be identified. Thus, to study the effect of agents on proliferative disorders, developmental or differentiative disorder, hematopoietic disorder as well disorders characterized by abnormal cell differentiation and/or survival, an abnormal extracellular structure, or an abnormality in a defense mechanism, for example, in a clinical trial, cells can be isolated and RNA prepared and analyzed for the levels of expression of Dkk, Dkk-related, and other genes implicated in the proliferative disorder, developmental or differentiative disorder, hematopoietic disorder as well as disorders characterized by abnormal cell differentiation and/or survival, an abnormal extracellular structure, or an abnormality in a defense mechanism, respectively. The levels of gene expression (*i.e.*, a gene expression pattern) can be quantified by Northern blot analysis or RT-PCR, as described herein, or alternatively by measuring the amount of protein produced, by one of the methods as described herein, or by measuring the levels of activity of Dkk, Dkk-related, or other genes. In this way, the gene expression pattern can serve as a marker, indicative of the physiological response of the cells to the agent. Accordingly, this response state may be determined before, and at various points during treatment of the individual with the agent.

In a preferred embodiment, the present invention provides a method for monitoring the effectiveness of treatment of a subject with an agent (*e.g.*, an agonist, antagonist, peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate identified by the screening assays described herein) comprising the steps of (i) obtaining a pre-administration sample from a subject prior to administration of the agent; (ii) detecting the level of expression of a Dkk or Dkk-related protein, mRNA, or genomic DNA in the preadministration sample; (iii) obtaining one or more post-administration samples from the subject; (iv) detecting the level of expression or activity of the Dkk or Dkk-related protein, mRNA, or genomic DNA in the post-administration samples; (v) comparing the level of expression or activity of the Dkk or Dkk-related protein, mRNA, or genomic DNA in the pre-administration sample with the Dkk or Dkk-related protein, mRNA, or genomic DNA in the post administration sample or samples; and (vi) altering the administration of the agent to the subject accordingly. For example, increased administration of the agent may be desirable to increase the expression or activity of Dkk or Dkk-related nucleic acid or protein to higher levels than detected, *i.e.*, to increase the effectiveness of the agent. Alternatively, decreased administration of the agent may be desirable to decrease expression or activity of Dkk or Dkk-related nucleic acid or protein to lower levels than detected, *i.e.* to decrease the effectiveness of the agent. According to such an embodiment, Dkk or Dkk-related expression or activity may be used as an indicator of the effectiveness of an agent, even in the absence of an observable phenotypic response.

C. Methods of Treatment:

The present invention provides for both prophylactic and therapeutic methods of treating a subject at risk of (or susceptible to) a disorder or having a disorder associated with aberrant Dkk or Dkk-related expression or activity. With regards to both prophylactic and therapeutic methods of treatment, such treatments may be specifically tailored or modified, based on knowledge obtained from the field of pharmacogenomics. "Pharmacogenomics", as used herein, refers to the application of genomics technologies such as gene sequencing, statistical genetics, and gene expression analysis to drugs in clinical development and on the market. More specifically, the term refers the study of

how a patient's genes determine his or her response to a drug (*e.g.*, a patient's "drug response phenotype", or "drug response genotype".) Thus, another aspect of the invention provides methods for tailoring an individual's prophylactic or therapeutic treatment with either the Dkk or Dkk-related molecules of the present invention or Dkk or Dkk-related modulators according to that individual's drug response genotype. Pharmacogenomics allows a clinician or physician to target prophylactic or therapeutic treatments to patients who will most benefit from the treatment and to avoid treatment of patients who will experience toxic drug-related side effects.

1. Prophylactic Methods

In one aspect, the invention provides a method for preventing in a subject, a disease or condition associated with an aberrant Dkk or Dkk-related expression or activity, by administering to the subject an agent which modulates Dkk or Dkk-related expression or at least one Dkk or Dkk-related activity. Subjects at risk for a disease which is caused or contributed to by aberrant Dkk or Dkk-related expression or activity can be identified by, for example, any or a combination of diagnostic or prognostic assays as described herein. Administration of a prophylactic agent can occur prior to the manifestation of symptoms characteristic of the Dkk or Dkk-related aberrancy, such that a disease or disorder is prevented or, alternatively, delayed in its progression. Depending on the type of Dkk or Dkk-related aberrancy, for example, an agonist or antagonist agent can be used for treating the subject. The appropriate agent can be determined based on screening assays described herein. The prophylactic methods of the present invention are further discussed in the following subsections.

2. Therapeutic Methods

Another aspect of the invention pertains to methods of modulating Dkk or Dkk-related expression or activity for therapeutic purposes. The modulatory method of the invention involves contacting a cell with an agent that modulates one or more of the activities of Dkk or Dkk-related protein activity associated with the cell. An agent that modulates Dkk or Dkk-related protein activity can be an agent as described herein, such

as a nucleic acid or a protein, a naturally-occurring target molecule of a Dkk or Dkk-related protein, a peptide, a Dkk or Dkk-related peptidomimetic, or other small molecule. In one embodiment, the agent stimulates one or more Dkk or Dkk-related protein activity. Examples of such stimulatory agents include active Dkk or Dkk-related
5 protein and a nucleic acid molecule encoding Dkk or Dkk-related that has been introduced into the cell. In another embodiment, the agent inhibits one or more Dkk or Dkk-related protein activity. Examples of such inhibitory agents include antisense Dkk or Dkk-related nucleic acid molecules and antibodies. These modulatory methods can be performed *in vitro* (e.g., by culturing the cell with the agent) or, alternatively, *in vivo*
10 (e.g., by administering the agent to a subject). As such, the present invention provides methods of treating an individual afflicted with a disease or disorder characterized by aberrant expression or activity of a Dkk or Dkk-related protein or nucleic acid molecule. The present invention also provides methods of modulating the function, morphology, proliferation, and/or differentiation of cells in the tissues in which a Dkk or Dkk-related
15 protein or nucleic acid molecule is expressed. Alternatively, Dkk or Dkk-related polypeptides, nucleic acids, and modulators thereof, can be used to treat disorders associated with abnormal or aberrant metabolism or function of cells in the tissues in which the Dkk or Dkk-related protein or nucleic acid molecule is expressed.

For example, tissues in which Dkk-3 is expressed include embryonic eye, bone,
20 and cartilage, fetal brain, lung, and kidney, and adult heart (in particular, atrioventricular valves and atrial myocytes), eye (in particular, the integrating bipolar and ganglion cells of the retina, the ciliary body, and lens epithelium), brain (in particular, neurons of the cortex and hippocampus), placenta, lung, and skeletal muscle. Accordingly, Dkk-3 polypeptides, nucleic acids, or modulators thereof, can be used to treat cardiovascular
25 disorders, such as ischemic heart disease (e.g., angina pectoris, myocardial infarction, and chronic ischemic heart disease), hypertensive heart disease, pulmonary heart disease, valvular heart disease (e.g., rheumatic fever and rheumatic heart disease, endocarditis, mitral valve prolapse, and aortic valve stenosis), congenital heart disease (e.g., valvular and vascular obstructive lesions, atrial or ventricular septal defect, and
30 patent ductus arteriosus), or myocardial disease (e.g., myocarditis, congestive cardiomyopathy, and hypertrophic cardiomyopathy).

In another embodiment, Dkk-3 polypeptides, nucleic acids, or modulators thereof, can be used to treat optic disorders such as diseases associated with amaurosis (*e.g.*, a. fugax and a. albuminuric) diseases associated with amblyopia, glaucoma, optic neuropathy (*e.g.*, ischemic neuropathy, optic neuritis, and infiltrative neuropathy),
5 ophthalmia (*e.g.*, o. catarrhal, trachoma, o. neuroparalytic, and conjunctiva), visual disorders resulting from systemic disease or disorders of other tissues (*e.g.*, diabetes mellitus, hyperthyroidism, and vitamin A or riboflavin deficiency), or tumors, neoplasms, and metastases.

In another embodiment, Dkk-3 polypeptides, nucleic acids, or modulators thereof, can be used to treat disorders of the brain, such as cerebral edema, senile
10 dementia of the Alzheimer type, epilepsy, amnesia, hydrocephalus, brain herniations, iatrogenic disease (due to, *e.g.*, infection, toxins, or drugs), inflammations (*e.g.*, bacterial and viral meningitis, encephalitis, and cerebral toxoplasmosis), cerebrovascular diseases (*e.g.*, hypoxia, ischemia, and infarction, intracranial hemorrhage and vascular
15 malformations, and hypertensive encephalopathy), and tumors (*e.g.*, neuroglial tumors, neuronal tumors, tumors of pineal cells, meningeal tumors, primary and secondary lymphomas, intracranial tumors, and medulloblastoma), and to treat injury or trauma to the brain.

In another embodiment, Dkk-3 polypeptides, nucleic acids, or modulators thereof, can be used to treat placental disorders, such as toxemia of pregnancy (*e.g.*,
20 preeclampsia and eclampsia), placentitis, or spontaneous abortion.

In another embodiment, Dkk-3 polypeptides, nucleic acids, or modulators thereof, can be used to treat pulmonary disorders, such as atelectasis, pulmonary
congestion or edema, chronic obstructive airway disease (*e.g.*, emphysema, chronic
25 bronchitis, bronchial asthma, and bronchiectasis), diffuse interstitial diseases (*e.g.*, sarcoidosis, pneumoconiosis, hypersensitivity pneumonitis, Goodpasture's syndrome, idiopathic pulmonary hemosiderosis, pulmonary alveolar proteinosis, desquamative interstitial pneumonitis, chronic interstitial pneumonia, fibrosing alveolitis, hamman-rich syndrome, pulmonary eosinophilia, diffuse interstitial fibrosis, Wegener's
30 granulomatosis, lymphomatoid granulomatosis, and lipid pneumonia), or tumors (*e.g.*, bronchogenic carcinoma, bronchioloalveolar carcinoma, bronchial carcinoid,

hamartoma, and mesenchymal tumors).

In another embodiment, Dkk-3 polypeptides, nucleic acids, or modulators thereof, can be used to treat disorders of skeletal muscle, such as muscular atrophy (due to, *e.g.*, denervation, malnutrition, loss of blood supply, or neuromuscular disease, *e.g.*, amyotonia congenita, amyotrophic lateral sclerosis of Charcot, and progressive muscular atrophy of Aran-Duchenne), myositis (due to, *e.g.*, bacterial, viral, fungal or parasitic infection), muscular dystrophies (*e.g.*, Duchenne type, Becker type, facioscapulohumeral, limb-girdle, myotonic dystrophy, and ocular myopathy), myasthenia gravis, or tumors and tumor-like lesions of muscles (*e.g.*, traumatic myositis ossificans, desmoids, musculoaponeurotic fibromatosis, Dupuytren's contracture, nodular (pseudosarcomatous) fasciitis, rhabdomyoma, rhabdomyosarcoma, and granular cell myoblastomas).

Tissues in which Dkk-4 is expressed include cerebellum, activated human T-lymphocytes, lung, and esophagus. Accordingly, in one embodiment, Dkk-4 polypeptides, nucleic acids, or modulators thereof, can be used to treat disorders of the cerebellum, such as disturbances of synergy (*e.g.*, asynergia or limb ataxia, dysmetria, decomposition of movement, hypermetria, hypometria, dysdiadochokinesia, hypotonia, tremor, dysarthria, nystagmus), disturbances of equilibrium (due to, *e.g.*, a lesion involving the vestibulocerebellum), disturbances of gait stance, or tone (due to, *e.g.*, a lesion or degeneration of the spinocerebellum), or tumors (*e.g.*, astrocytoma and medulloblastoma).

In another embodiment, Dkk-4 polypeptides, nucleic acids, or modulators thereof, can be used to treat lymphocytic disorders, such as lymphopenia, lymphocytosis, acute and chronic lymphadenitis, malignant lymphomas (*e.g.*, Non-Hodgkin's lymphomas, Hodgkin's lymphomas, leukemias, multiple myeloma, histiocytoses, and angioimmunoblastic lymphadenopathy).

In another embodiment, Dkk-4 polypeptides, nucleic acids, or modulators thereof, can be used to treat pulmonary disorders, such as atelectasis, pulmonary congestion or edema, chronic obstructive airway disease (*e.g.*, emphysema, chronic bronchitis, bronchial asthma, and bronchiectasis), diffuse interstitial diseases (*e.g.*, sarcoidosis, pneumoconiosis, hypersensitivity pneumonitis, Goodpasture's syndrome,

idiopathic pulmonary hemosiderosis, pulmonary alveolar proteinosis, desquamative interstitial pneumonitis, chronic interstitial pneumonia, fibrosing alveolitis, hamman-rich syndrome, pulmonary eosinophilia, diffuse interstitial fibrosis, Wegener's granulomatosis, lymphomatoid granulomatosis, and lipid pneumonia), or tumors (*e.g.*,
5 bronchogenic carcinoma, bronchioloalveolar carcinoma, bronchial carcinoid, hamartoma, and mesenchymal tumors).

In another embodiment, Dkk-4 polypeptides, nucleic acids, or modulators thereof, can be used to treat esophageal disorders, such as neuromuscular disturbances (*e.g.*, achalasia, annular narrowings, Schatzki's rings, hiatal hernia, Mallory-Weiss
10 syndrome), esophagitis (due to *e.g.*, bacteremia, viremia, fungal infections, uremia, graft-versus-host disease, chemotherapy, radiation, and prolonged gastric intubation), diverticula (*e.g.*, Zenker's diverticulum), systemic sclerosis, varices (due to, *e.g.*, portal hypertension, systemic amyloidosis and sarcoidosis), or tumors or neoplasms (*e.g.*, leiomyoma, fibromas, lipomas, hemangiomas, lymphangiomas, squamous papillomas,
15 adenocarcinomas and undifferentiated carcinomas, and sarcomas).

Dkk-1 is highly expressed, for example, in placenta. Accordingly, Dkk-1 polypeptides, nucleic acids, or modulators thereof, can be used to treat placental disorders, such as toxemia of pregnancy (*e.g.*, preeclampsia and eclampsia), placentitis, or spontaneous abortion.

20 Tissues in which Dkk-2 is expressed include, for example, heart, brain, placenta, lung, and skeletal muscle. Accordingly, Dkk-2 polypeptides, nucleic acids, or modulators thereof, can be used to treat cardiovascular disorders, such as ischemic heart disease (*e.g.*, angina pectoris, myocardial infarction, and chronic ischemic heart disease), hypertensive heart disease, pulmonary heart disease, valvular heart disease (*e.g.*,
25 rheumatic fever and rheumatic heart disease, endocarditis, mitral valve prolapse, and aortic valve stenosis), congenital heart disease (*e.g.*, valvular and vascular obstructive lesions, atrial or ventricular septal defect, and patent ductus arteriosus), or myocardial disease (*e.g.*, myocarditis, congestive cardiomyopathy, and hypertrophic cardiomyopathy).

In another embodiment, Dkk-2 polypeptides, nucleic acids, or modulators thereof, can be used to treat disorders of the brain, such as cerebral edema, senile dementia of the Alzheimer type, epilepsy, amnesia, hydrocephalus, brain herniations, iatrogenic disease (due to, *e.g.*, infection, toxins, or drugs), inflammations (*e.g.*, bacterial
5 and viral meningitis, encephalitis, and cerebral toxoplasmosis), cerebrovascular diseases (*e.g.*, hypoxia, ischemia, and infarction, intracranial hemorrhage and vascular malformations, and hypertensive encephalopathy), and tumors (*e.g.*, neuroglial tumors, neuronal tumors, tumors of pineal cells, meningeal tumors, primary and secondary lymphomas, intracranial tumors, and medulloblastoma), and to treat injury or trauma to
10 the brain.

In another embodiment, Dkk-2 polypeptides, nucleic acids, or modulators thereof, can be used to treat placental disorders, such as toxemia of pregnancy (*e.g.*, preeclampsia and eclampsia), placentitis, or spontaneous abortion.

In another embodiment, Dkk-2 polypeptides, nucleic acids, or modulators
15 thereof, can be used to treat pulmonary disorders, such as atelectasis, pulmonary congestion or edema, chronic obstructive airway disease (*e.g.*, emphysema, chronic bronchitis, bronchial asthma, and bronchiectasis), diffuse interstitial diseases (*e.g.*, sarcoidosis, pneumoconiosis, hypersensitivity pneumonitis, Goodpasture's syndrome, idiopathic pulmonary hemosiderosis, pulmonary alveolar proteinosis, desquamative
20 interstitial pneumonitis, chronic interstitial pneumonia, fibrosing alveolitis, hamman-rich syndrome, pulmonary eosinophilia, diffuse interstitial fibrosis, Wegener's granulomatosis, lymphomatoid granulomatosis, and lipid pneumonia), or tumors (*e.g.*, bronchogenic carcinoma, bronchioloalveolar carcinoma, bronchial carcinoid, hamartoma, and mesenchymal tumors).

25 In another embodiment, Dkk-2 polypeptides, nucleic acids, or modulators thereof, can be used to treat disorders of skeletal muscle, such as muscular atrophy (due to, *e.g.*, denervation, malnutrition, loss of blood supply, or neuromuscular disease, *e.g.*, amyotonia congenita, amyotrophic lateral sclerosis of Charcot, and progressive muscular atrophy of Aran-Duchenne), myositis (due to, *e.g.*, bacterial, viral, fungal or parasitic
30 infection), muscular dystrophies (*e.g.*, Duchenne type, Becker type, facioscapulohumeral, limb-girdle, myotonic dystrophy, and ocular myopathy),

myasthenia gravis, or tumors and tumor-like lesions of muscles (*e.g.*, traumatic myositis ossificans, desmoids, musculoaponeurotic fibromatosis, Dupuytren's contracture, nodular (pseudosarcomatous) fasciitis, rhabdomyoma, rhabdomyosarcoma, and granular cell myoblastomas).

5 Soggy-1 is expressed in, for example, testis (*e.g.*, spermatogenic epithelium of the seminiferous tubules, spermatogonia) and in embryonic developing dorsal root ganglia, cartilage primordium of the nasal septum, and the eye. Accordingly, Soggy-1 polypeptides, nucleic acids, or modulators thereof, can be used to treat testicular disorders, such as unilateral testicular enlargement (*e.g.*, nontuberculous, granulomatous
10 orchitis), inflammatory diseases resulting in testicular dysfunction (*e.g.*, gonorrhea and mumps), and tumors (*e.g.*, germ cell tumors, interstitial cell tumors, androblastoma, testicular lymphoma and adenomatoid tumors). In another embodiment, Soggy-1 polypeptides, nucleic acids, or modulators thereof, can be used to treat infertility due to, for example, spermatogenetic failure.

15 In one aspect, the above-described methods involve administering an agent (*e.g.*, an agent identified by a screening assay described herein), or combination of agents that modulates (*e.g.*, upregulates or downregulates) Dkk or Dkk-related expression or activity. In another embodiment, the method involves administering a Dkk or Dkk-related protein or nucleic acid molecule as therapy to compensate for reduced or aberrant
20 Dkk or Dkk-related expression or activity.

A preferred embodiment of the present invention involves a method for treatment of a disease or disorder associated with a Dkk or Dkk-related protein which includes the step of administering a therapeutically effective amount of an antibody to a Dkk or Dkk-related protein to a subject. As defined herein, a therapeutically effective amount of
25 antibody (*i.e.*, an effective dosage) ranges from about 0.001 to 30 mg/kg body weight, preferably about 0.01 to 25 mg/kg body weight, more preferably about 0.1 to 20 mg/kg body weight, and even more preferably about 1 to 10 mg/kg, 2 to 9 mg/kg, 3 to 8 mg/kg, 4 to 7 mg/kg, or 5 to 6 mg/kg body weight. The skilled artisan will appreciate that certain factors may influence the dosage required to effectively treat a subject, including
30 but not limited to the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and other diseases present. Moreover, treatment of a

subject with a therapeutically effective amount of an antibody can include a single treatment or, preferably, can include a series of treatments. In a preferred example, a subject is treated with antibody in the range of between about 0.1 to 20 mg/kg body weight, one time per week for between about 1 to 10 weeks, preferably between 2 to 8 weeks, more preferably between about 3 to 7 weeks, and even more preferably for about 4, 5, or 6 weeks. It will also be appreciated that the effective dosage of antibody used for treatment may increase or decrease over the course of a particular treatment. Changes in dosage may result from the results of diagnostic assays as described herein.

Stimulation of Dkk or Dkk-related activity is desirable in situations in which Dkk or Dkk-related activity is abnormally downregulated and/or in which increased Dkk or Dkk-related activity is likely to have a beneficial effect. Likewise, inhibition of Dkk or Dkk-related activity is desirable in situations in which Dkk or Dkk-related activity is abnormally upregulated and/or in which decreased Dkk or Dkk-related activity is likely to have a beneficial effect. One example of such a situation is where a subject has a disorder characterized by aberrant development or cellular differentiation. Another example of such a situation is where the subject has a proliferative disease (*e.g.*, cancer) or a neurogenerative disorder. Yet another example of such a situation is where it is desirable to achieve tissue regeneration in a subject (*e.g.*, where a subject has undergone brain or spinal cord injury and it is desirable to regenerate neuronal tissue in a regulated manner.)

Accordingly, in one embodiment, the disease is a disease characterized by an abnormal cell proliferation, differentiation, and/or survival. For example, the disease can be a hyper- or hypoproliferative disease. The invention also provides methods for treating diseases characterized by an abnormal cell proliferation, differentiation, and/or survival in a subject, which are not characterized by an abnormal Dkk or Dkk-related activity (*e.g.*, hDkk-3 activity). In fact, since Dkk is likely to be capable of modulating the proliferative state of a cell (*i.e.*, state of proliferation, differentiation, and or survival of a cell), Dkk can regulate disease wherein the abnormal proliferative state of a cell results from a defect other than an abnormal Dkk activity.

Hyperproliferative diseases can be treated with Dkk or Dkk-related (*e.g.*, hDkk-3) therapeutics include neoplastic and hyperplastic diseases, such as various forms of cancers and leukemias, and fibroproliferative disorders. Other hyperproliferative diseases that can be treated or prevented with the subject Dkk or Dkk-related therapeutics (*e.g.* hDkk-3 therapeutics) include malignant conditions, premalignant conditions, and benign conditions. The condition to be treated or prevented can be a solid tumor, such as a tumor arising in an epithelial tissue. Accordingly, treatment of such a cancer could comprise administration to the subject of a Dkk or Dkk-related therapeutic decreasing the interaction of Dkk with a Dkk receptor. Other cancers that can be treated or prevented with a Dkk or Dkk-related protein include cancers of the epithelia (*e.g.*, carcinomas of the pancreas, kidney, stomach, colon, esophagus liver, secretory glands (*e.g.*, adenocarcinoma) bladder, lung, breast, skin (*e.g.*, malignant melanoma, seminoma squamous adenocarcinoma), reproductive tract including prostate gland, testis, ovary, cervix and uterus); cancers of the hematopoietic and immune system (*e.g.*, leukemias and lymphomas); cancers of the central nervous, brain system and eye (*e.g.*, malignant astrocytoma, gliomas, neuroblastoma and retinoblastoma); and cancers of connective tissues, bone, heart, muscles and vasculature (*e.g.*, sarcomas, for example, osteosarcoma). Additional solid tumors within the scope of the invention include those that can be found in a medical textbook.

The condition to be treated or prevented can also be a soluble tumor, such as leukemia, either chronic or acute, including chronic or acute myelogenous leukemia, chronic or acute lymphocytic leukemia, promyelocytic leukemia, monocytic leukemia, myelomonocytic leukemia, and erythroleukemia. Yet other proliferative disorders that can be treated with a Dkk or Dkk-related therapeutic of the invention include heavy chain disease, multiple myeloma, lymphoma, *e.g.*, Hodgkin's lymphoma and non-Hodgkin's lymphoma, and Waldenstroem's macroglobulemia.

Diseases or conditions characterized by a solid or soluble tumor can be treated by administering a Dkk or Dkk-related therapeutic either locally or systemically, such that aberrant cell proliferation is inhibited or decreased. Methods for administering the compounds of the invention are further described below.

The invention also provides methods for preventing the formation and/or development of tumors. For example, the development of a tumor can be preceded by the presence of a specific lesion, such as a pre-neoplastic lesion, *e.g.*, hyperplasia, metaplasia, and dysplasia, which can be detected, *e.g.*, by cytologic methods. Such
5 lesions can be found, *e.g.*, in epithelial tissue. Thus, the invention provides a method for inhibiting progression of such a lesion into a neoplastic lesion, comprising administering to the subject having a preneoplastic lesion an amount of a Dkk or Dkk-related therapeutic sufficient to inhibit progression of the preneoplastic lesion into a neoplastic lesion.

10 The invention also provides for methods for treating or preventing diseases or conditions in which proliferation of cells is desired. For example, Dkk or Dkk-related therapeutics can be used to stimulate tissue repair or wound healing, such as after surgery or to stimulate tissue healing from burns. Other diseases in which proliferation of cells is desired are hypoproliferative diseases, *i.e.*, diseases characterized by an
15 abnormally low proliferation of certain cells.

In yet another embodiment, the invention provides a method for treating or preventing diseases or conditions characterized by aberrant cell differentiation. Accordingly, the invention provides methods for stimulating cellular differentiation in conditions characterized by an inhibition of normal cell differentiation which may or
20 may not be accompanied by excessive proliferation. Alternatively, Dkk or Dkk-related therapeutics can be used to inhibit differentiation of specific cells.

In a preferred method, the aberrantly proliferating and/or differentiating cell is a cell present in the nervous system. A role for Dkk in the nervous system is suggested at least in part from the fact that human Dkk-3 is expressed in human fetal brain.

25 Accordingly, the invention provides methods for treating diseases or conditions associated with a central or peripheral nervous system. For example, the invention provides methods for treating lesions of the nervous system associated with an aberrant proliferation, differentiation or survival of any of the following cells: cells of the central nervous system including neurons and glial cells (*e.g.*, astrocytes and oligodendrocytes)
30 and supporting cells of peripheral neurons (*e.g.*, Schwann cells and satellite cells). Disorders of the nervous system include, but are not limited to: spinal cord injuries,

brain injuries, brain tumors (*e.g.*, astrocytic tumors, for example, astrocytomas and glioblastomas), lesions associated with surgery, ischemic lesions, malignant lesions, infectious lesions, degenerative lesions (*e.g.*, Parkinson's disease, Alzheimer's disease, Huntington's chorea, amyotrophic lateral sclerosis), demyelinating diseases (*e.g.*,
5 multiple sclerosis, human immunodeficiency associated myelopathy, transverse myelopathy, progressive multifocal leukoencephalopathy, pontine myelinolysis), motor neuron injuries, progressive spinal muscular atrophy, progressive bulbar palsy, primary lateral sclerosis, infantile and juvenile muscular atrophy, progressive bulbar paralysis of childhood (*i.e.*, Fazio-Londe syndrome), poliomyelitis, and hereditary motorsensory
10 neuropathy (*i.e.*, Charcot-Marie-Tooth disease).

In another embodiment, the invention provides a method for enhancing the survival and/or stimulating proliferation and/or differentiation of cells and tissues *in vitro*. In a preferred embodiment, Dkk or Dkk-related therapeutics are used to promote tissue regeneration and/or repair (*e.g.*, to treat nerve injury). For example, tissues from a
15 subject can be obtained and grown *in vitro* in the presence of a Dkk or Dkk-related therapeutic, such that the tissue cells are stimulated to proliferate and/or differentiate. The tissue can then be readministered to the subject.

Among the approaches which may be used to ameliorate disease symptoms involving an aberrant Dkk or Dkk-related activity and/or an abnormal cell proliferation,
20 differentiation, and/or survival, are, for example, antisense, ribozyme, and triple helix molecules described above. Examples of suitable compounds include the antagonists, agonists or homologues described in detail above.

Yet other Dkk or Dkk-related therapeutics consist of a first peptide comprising a Dkk or Dkk-related peptide capable of binding to a Dkk receptor, and a second peptide
25 which is cytotoxic. Such therapeutics can be used to specifically target and lyse cells expressing or overexpressing a receptor for Dkk.

3. Pharmacogenomics

The Dkk or Dkk-related molecules of the present invention, as well as agents, or
30 modulators which have a stimulatory or inhibitory effect on Dkk or Dkk-related activity (*e.g.*, Dkk or Dkk-related gene expression) as identified by a screening assay described

herein can be administered to individuals to treat (prophylactically or therapeutically) disorders (*e.g.*, proliferative or developmental disorders) associated with aberrant Dkk or Dkk-related activity. In conjunction with such treatment, pharmacogenomics (*i.e.*, the study of the relationship between an individual's genotype and that individual's response to a foreign compound or drug) may be considered. Differences in metabolism of therapeutics can lead to severe toxicity or therapeutic failure by altering the relation between dose and blood concentration of the pharmacologically active drug. Thus, a physician or clinician may consider applying knowledge obtained in relevant pharmacogenomics studies in determining whether to administer a Dkk or Dkk-related molecule or Dkk or Dkk-related modulator as well as tailoring the dosage and/or therapeutic regimen of treatment with a Dkk or Dkk-related molecule or Dkk or Dkk-related modulator.

Pharmacogenomics deals with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See *e.g.*, Eichelbaum, M., *Clin Exp Pharmacol Physiol*, 1996, 23(10-11):983-985 and Linder, M.W., *Clin Chem*, 1997, 43(2):254-266. In general, two types of pharmacogenetic conditions can be differentiated. Genetic conditions transmitted as a single factor altering the way drugs act on the body (altered drug action) or genetic conditions transmitted as single factors altering the way the body acts on drugs (altered drug metabolism). These pharmacogenetic conditions can occur either as rare genetic defects or as naturally-occurring polymorphisms. For example, glucose-6-phosphate dehydrogenase deficiency (G6PD) is a common inherited enzymopathy in which the main clinical complication is haemolysis after ingestion of oxidant drugs (anti-malarials, sulfonamides, analgesics, nitrofurans) and consumption of fava beans.

One pharmacogenomics approach to identifying genes that predict drug response, known as "a genome-wide association", relies primarily on a high-resolution map of the human genome consisting of already known gene-related markers (*e.g.*, a "bi-allelic" gene marker map which consists of 60,000-100,000 polymorphic or variable sites on the human genome, each of which has two variants.) Such a high-resolution genetic map can be compared to a map of the genome of each of a statistically significant number of patients taking part in a Phase II/III drug trial to identify markers

associated with a particular observed drug response or side effect. Alternatively, such a high resolution map can be generated from a combination of some ten-million known single nucleotide polymorphisms (SNPs) in the human genome. As used herein, a "SNP" is a common alteration that occurs in a single nucleotide base in a stretch of
5 DNA. For example, a SNP may occur once per every 1000 bases of DNA. A SNP may be involved in a disease process, however, the vast majority may not be disease-associated. Given a genetic map based on the occurrence of such SNPs, individuals can be grouped into genetic categories depending on a particular pattern of SNPs in their individual genome. In such a manner, treatment regimens can be tailored to groups of
10 genetically similar individuals, taking into account traits that may be common among such genetically similar individuals.

Alternatively, a method termed the "candidate gene approach", can be utilized to identify genes that predict drug response. According to this method, if a gene that encodes a drug's target is known (*e.g.*, a Dkk protein or Dkk receptor of the present
15 invention), all common variants of that gene can be fairly easily identified in the population and it can be determined if having one version of the gene versus another is associated with a particular drug response.

As an illustrative embodiment, the activity of drug metabolizing enzymes is a major determinant of both the intensity and duration of drug action. The discovery of
20 genetic polymorphisms of drug metabolizing enzymes (*e.g.*, N-acetyltransferase 2 (NAT 2) and cytochrome P450 enzymes CYP2D6 and CYP2C19) has provided an explanation as to why some patients do not obtain the expected drug effects or show exaggerated drug response and serious toxicity after taking the standard and safe dose of a drug. These polymorphisms are expressed in two phenotypes in the population, the extensive
25 metabolizer (EM) and poor metabolizer (PM). The prevalence of PM is different among different populations. For example, the gene coding for CYP2D6 is highly polymorphic and several mutations have been identified in PM, which all lead to the absence of functional CYP2D6. Poor metabolizers of CYP2D6 and CYP2C19 quite frequently experience exaggerated drug response and side effects when they receive standard doses.
30 If a metabolite is the active therapeutic moiety, PM show no therapeutic response, as demonstrated for the analgesic effect of codeine mediated by its CYP2D6-formed

- 118 -

metabolite morphine. The other extreme are the so called ultra-rapid metabolizers who do not respond to standard doses. Recently, the molecular basis of ultra-rapid metabolism has been identified to be due to CYP2D6 gene amplification.

Alternatively, a method termed the "gene expression profiling", can be utilized to
5 identify genes that predict drug response. For example, the gene expression of an animal dosed with a drug (*e.g.*, a Dkk molecule or Dkk modulator of the present invention) can give an indication whether gene pathways related to toxicity have been turned on.

Information generated from more than one of the above pharmacogenomics
10 approaches can be used to determine appropriate dosage and treatment regimens for prophylactic or therapeutic treatment an individual. This knowledge, when applied to dosing or drug selection, can avoid adverse reactions or therapeutic failure and thus enhance therapeutic or prophylactic efficiency when treating a subject with a Dkk molecule or Dkk or Dkk-related modulator, such as a modulator identified by one of the
15 exemplary screening assays described herein.

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application are hereby incorporated by reference.

20

EXAMPLES

The invention is based, at least in part, on the discovery of a family of genes encoding human cysteine-rich secreted proteins which are related to *Xenopus* Dickkopf (Dkk) proteins. This family includes hDkk-1, hDkk-2, hDkk-3, and hDkk-4. hDkks 1-4
25 contain two highly conserved cysteine-rich domains (CRDs), the most C-terminal of which demonstrates similarity to the colipase protein family. The invention is based also in part on the discovery of a family of Dkk-related proteins, referred to as Soggy proteins, as well as the genes encoding Soggy proteins. Soggy-1 is a novel secreted protein which is related to the N-terminal region of Dkk-3 but lacks CRDs. The
30 following examples illustrate the structure and function of each of these novel human secreted proteins.

Example 1: Isolation And Characterization of Human hDkk-3 cDNA

In this example, the isolation and characterization of the gene encoding human Dkk-3 (also referred to as "hDkk-3", "Cysteine Rich Secreted Protein-1", "CRSP-1"
5 "CRISPY-1" or "TANGO 59") is described.

Isolation of a Human Dkk-3 cDNA

The invention is based at least in part on the discovery of a human gene encoding a secreted protein, referred to herein as human Dickkopf-3 (hDkk-3). A partial cDNA
10 was isolated using a Signal Sequence Trap method. This methodology takes advantage of the fact that molecules such as Dkk have an amino terminal signal sequence which directs certain secreted and membrane-bound proteins through the cellular secretory apparatus.

Briefly, a randomly primed cDNA library using mRNA prepared from human
15 fetal brain tissue (Clontech, Palo Alto CA) was made by using the Stratagene-ZAP-cDNA Synthesis™ kit, (catalog #20041). The cDNA was ligated into the mammalian expression vector pTrap adjacent to a cDNA encoding placental alkaline phosphatase lacking a secretory signal. The plasmids were transformed into *E. coli* and DNA was prepared using the Wizard™ DNA purification kit (Promega). DNA was transfected into
20 COS-7 cells with lipofectamine™ (Gibco-BRL). After 48 hours incubation the COS cell supernatants were assayed for alkaline phosphatase on a Wallac Micro-Beta scintillation counter using the Phospha-Light™ kit (Tropix Inc. Catalog #BP300). The individual plasmid DNAs scoring positive in the COS cell Alkaline Phosphatase secretion assay were further analyzed by DNA sequencing using standard procedures.

25 Using a partial cDNA isolated by the above-described method (clone Amhb3c2), a full length cDNA encoding human Dkk-3 was isolated from a lambda Ziplox™ human fetal brain cDNA library using conventional hybridization techniques (Sambrook *et al.*, *supra*). The nucleotide sequence encoding the full length human Dkk-3 protein is shown in Figure 1 and is set forth as SEQ ID NO: 1. The full length protein encoded by
30 this nucleic acid is comprised of about 350 amino acids and has the amino acid sequence

shown in Figure 1 and set forth as SEQ ID NO:2. The coding portion (open reading frame) of SEQ ID NO:1 is set forth as SEQ ID NO:3. DNA for the clone Fmhb059 was deposited with the ATCC as Accession No. 98452.

5 Analysis of Human hDkk-3

Determination of the hydrophobicity profile of human Dkk-3 having the amino acid sequence set forth in SEQ ID NO:2 indicated the presence of a hydrophobic region from about amino acid 1 to about amino acid 22 of SEQ ID NO:2. Further analysis of the amino acid sequence SEQ ID NO:2 using a signal peptide prediction program
10 predicted the presence of a signal peptide from about amino acid 1 to about amino acid 22 of SEQ ID NO:2. Accordingly, the mature hDkk-3 protein includes about 328 amino acids spanning from about amino acid 23 to about amino acid 350 of SEQ ID NO:2. The presence of the signal sequence, in addition to the fact that hDkk-3 has been identified using a Signal Sequence Trap system, indicates that hDkk-3 is a secreted
15 protein. Furthermore, the prediction of such a signal peptide and signal peptide cleavage site can be made, for example, utilizing the computer algorithm SIGNALP (Nielsen, *et al.*, (1997) *Protein Engineering* 10:1-6).

Examination of the cDNA sequence depicted in Figure 1 shows that human Dkk-3 is particularly rich in cysteine residues. As shown in Figure 1, hDkk-3 contains 20
20 cysteine residues located between amino acid 147 and amino acid 284 of SEQ ID NO: 2. This region has been termed the cysteine-rich region. These cysteine residues can form 10 disulfide bridges.

A BLAST search (Altschul *et al.*, (1990) *J. Mol. Biol.* 215:403) of the nucleotide and the amino acid sequences of hDkk-3 has revealed that hDkk-3 is similar to a chicken
25 cDNA encoding a protein of unknown function having GenBank Accession No. D26311. This cDNA was isolated from a chicken lens cDNA library and was shown to be expressed in lens fibers and lens epithelium, but not in neural retina nor in liver cells. (Sawada *et al.*, (1996) *Int. J. Dev. Biol.* 40:531). hDkk-3 and the chicken protein have 56% amino acid sequence identity and 72% amino acid sequence similarity. The amino
30 acid sequence similarity between the chicken protein and human Dkk-3 is particularly high in the cysteine-rich domain of hDkk-3 which is located between amino acids 147

and 284 of SEQ ID NO: 2. In particular, the 20 cysteine residues of hDkk-3 located in this region are present in the chicken protein.

Two genes recently identified in a screen for suppressors of glioblastoma formation (Ligon *et al.* (1997) *Oncogene* 14:1075-1081) also show homology to hDkk-3. These genes, RIG ("Regulated In Glioblastoma") and RIG-like 7-1 (GenBank Accession Nos. U32331 and AF034208, respectively) were identified in a differential screen for mRNAs regulated by the introduction of a normal copy of chromosome 10 into a glioblastoma cell line harboring a deletion in chromosome 10 that promotes tumorigenesis. A schematic diagram summarizing the relationship between the sequences of the hDkk-3 and the RIG genes is presented as Figure 12. The indicated region of identity between hDkk-3 and RIG comprises a short portion of the 3' UTR of the human Dkk-3 mRNA (*e.g.*, RIG mRNA is ~100% identical to residues 2479 to 2153 of SEQ ID NO:1). RIG-like 7-1 is homologous to hDkk-3 across a longer region (*e.g.*, 97% identical from about nucleotides 316 to 2438 of SEQ ID NO:1) although the encoded RIG-like 7-1 protein lacks the Dkk N-terminal signal sequence and is not therefore predicted to be a secreted protein. These data associate hDkk-3 with human glioblastoma and suggest that hDkk-3 may be important in the suppression of the tumorigenic phenotype. A role in glioblastoma is also consistent with the high level of hDkk-3 mRNA expression observed in human brain tissue. In addition, the co-localization of the hDkk-3, RIG and RIG-like genes to a region of chromosome 11 (11p15.1) implicated in the development of human malignant astrocytoma (Ligon *et al.*, *supra*) further indicates a role for these genes in tumorigenesis.

Human hDkk-3 protein has also some amino acid sequence similarity to metallothionein, particularly in the cysteine-rich domain.

Tissue Distribution of hDkk-3 mRNA

For Northern blots, all hybridizations were to Clontech Multiple Tissue Northern Blots and were performed in ExpressHyb solution (Clontech) for 1-20 hours. All probes were prepared by random primed radiolabelling (Prime-It, Stratagene). Blots were washed sequentially to a final stringency of 0.2x SSC/0.2% SDS and exposed to autoradiographic film. Hybridizations of a control β -actin cDNA probe consistently

demonstrated even loading of the Northern blots. The results of hybridization of the probe to various mRNA samples are described below.

Hybridization of a Clontech Fetal Multiple Tissue Northern (MTN) blot (Clontech, LaJolla, CA) containing RNA from fetal brain, lung, liver, and kidney
5 indicated the presence of high levels of hDkk-3 mRNA (~2.5kb) in fetal brain, lung, and slightly lower levels of hDkk-3 mRNA in fetal kidney. However, no significant level of hDkk-3 mRNA was found in fetal liver.

Hybridization of a Clontech human Multiple Tissue Northern (MTN) blot (Clontech, LaJolla, CA) containing RNA from adult heart, brain, placenta, lung, liver,
10 skeletal muscle, kidney, and pancreas with a human Dkk-3 probe indicated the presence of high levels of hDkk-3 mRNA in heart, slightly lower levels in brain, and much lower levels in placenta and lung. Some hDkk-3 mRNA was also found in adult skeletal muscle. However, no significant levels of hDkk-3 mRNA was observed in adult liver,
15 kidney, or pancreas. Interestingly, the chicken gene which is homologous to hDkk-3 was not expressed at detectable levels in liver either (Sawada *et al.*, (1996) *Int. J. Dev. Biol.* 40:531).

Further hybridization of a Clontech human Multiple Tissue Northern (MTN) blot (Clontech, LaJolla, CA) including RNA from bone marrow, adrenal gland, trachea,
lymph node, spinal cord, thyroid, and stomach revealed high levels of expression of
20 hDkk-3 in mRNA isolated from adult spinal cord, and lower level expression in adrenal gland, trachea, thyroid, and stomach.

Thus, hDkk-3 is expressed in a tissue specific manner, with the strongest expression observed in brain, heart, and spinal cord.

25 **Example 2: Isolation And Characterization of mDkk-3 cDNA**

In this example, the isolation and characterization of the gene encoding murine Dkk-3 (also referred to as "mDkk-3", "murine Cysteine Rich Secreted Protein-1", "murine CRSP-1" or "murine CRISPY-1") is described.

Identification of a Murine Dkk-3 cDNA

A full length mDkk-3 cDNA was identified by comparison of the hDkk-3 sequence to a proprietary EST Database using the BLAST-X algorithm. A single clone identified in a adult mouse brain cDNA library was obtained and sequenced fully. DNA
5 for the clone Fmmb059s was deposited with the ATCC as Accession No. 98634. mDkk-3 is predicted to have a signal peptide from residues 1 to 23 of SEQ ID NO:17, cleavage of which results in a mature protein having 326 amino acids in length corresponding to amino acids 24 to 349 of SEQ ID NO:17.

10 Tissue Distribution of mDkk-3 mRNA

To determine the expression pattern of mDkk-3, *in situ* hybridization was performed as follows. Normal mouse embryos and adult mouse tissues were collected from C57BL/6 mice, embedded in TissueTek™ O.C.T Compound (Sakura Finetek U.S.A., Inc., Torrance, CA), frozen on dry ice, and stored at -80°C. Cryostat serial
15 sections (8µm) were thaw mounted on Superfrost Plus™ slides (VWR Scientific, West Chester, PA.) and air dried on a slide warmer at 40°C for 20 minutes. Sections were then fixed with 4% formaldehyde in DEPC treated 0.1 M phosphate-buffered saline (PBS, pH 7.5) at room temperature for 10 minutes and rinsed twice in DEPC-PBS. Sections were rinsed in 0.1 M triethanolamine-HCl (TEA, pH 8.0) , incubated in 0.25%
20 acetic anhydride-TEA for 10 minutes and rinsed in DEPC-2X SSC (standard sodium citrate). Sections were dehydrated through a series of graded ethanols, incubated in 100% chloroform for 5 minutes, rinsed in 100% and 95% ethanol for 1 minutes and air dried.

Antisense and sense RNA transcripts were prepared by *in vitro* transcription
25 (Riboprobe Gemini System™, Promega) of PCR amplified cDNA templates. Template amplification primers were as follows;

mDkk-3 forward 5'-CAGTGAGTGCTGTGGAGACC-3' (SEQ ID NO:30), and

reverse 5'-TCTTCAGTCAGGCTCCTCTC-3' (SEQ ID NO:31).

Probes were labeled with ³⁵S-UTP (NEN) and purified on G-25 spin columns
30 (Pharmacia). The hybridization cocktail contained: 50% formamide, 10% dextran

- 124 -

sulfate, 0.1% sodium dodecyl sulfate (SDS), 0.1% sodium thiosulfate, 1X Denhardt's solution, 0.6 M NaCl, 10 mM Tris (pH 7.5), 1 mM EDTA, 100 mM dithiothreitol (DTT), 0.1 mg/ml sheared salmon sperm, 50 µg/ml yeast tRNA, 0.5 mg/ml yeast total RNA, and ³⁵S-UTP labeled probe at a concentration of 5 X 10⁷ c.p.m./100µl of final hybridization solution; 100 µl of hybridization solution was put on each section. The sections were then covered with a glass coverslip and incubated in a humidified chamber at 55°C for 18 h. After hybridization, slides were washed with 2 X SSC. Sections were then sequentially incubated at 37°C in TNE (a solution containing 10 mM Tris-HCl (pH 7.6), 500 mM NaCl, and 1 mM EDTA), for 10 minutes, in TNE with 10ug/ml RNase A for 30 minutes, and finally in TNE for 10 minutes. Slides were then rinsed with 2 X SSC at room temperature, washed in 2 X SSC at 50°C for 1 h, 0.2 X SSC at 55°C for 1h, and 0.2 X SSC at 60°C for 1 h. Sections were dehydrated with a series of graded concentrations of ethanol 0.3 M ammonium acetate, air dried and exposed to Kodak Biomax MR™ scientific imaging film for 6 days at room temperature.

mDkk-3 expression in the brain was found to be highly localized to the cortex and hippocampus but was not observed in the dentate gyrus. Higher power magnification confirmed the mDkk-3 mRNA was localized to neurons within these structures. In the adult eye, mDkk-3 mRNA was found to be highly expressed in the retina, ciliary body, and lens epithelium. Expression in the retina was localized to the integrating bipolar and ganglion cells. In adult heart, mDkk-3 was detected in the atrioventricular valves and also in myocytes of the atria. Expression was highly restricted to the atria and noticeably absent from ventricular tissue. High level expression of mDkk3 mRNA was also observed in developing eye, bone and cartilage in day 14 embryos. These findings corroborate and extend the northern analysis of hDkk-3 mRNA expression in human tissues and also suggest that Dkk-3 may play a role in bone and ocular physiology in addition to functions in neural and cardiac tissues.

Example 3: Secretion and Post-Translational Modification of Dkk-3

This example describes the secretion and post-translational modification (*e.g.*, glycosylation and processing) of hDkk-3 as well as methods for small and large scale purification of hDkk-3.

5

hDkk-3 Expression Constructs

Expression constructs for two forms of hDkk-3 were prepared using the mammalian expression vector pMET-stop. Form-1 comprised a cDNA incorporating the complete 350aa hDkk-3 protein coding sequence (hDkk-3flag.long) and form-2
10 comprised the entire hDkk-3 protein coding sequence except for the final 18 amino acids (hDkk-3flag.short). A C-terminal sequence encoding the FLAG epitope (DYKDDDDK) (SEQ ID NO:19) was added to both hDkk-3 forms for ease of detection and purification. hDkk-3flag cDNAs were generated by PCR from a full length hDkk-3 cDNA template and ligated into pMET-stop using EcoR1 and Sal1 restriction sites.

15

Trial Transfection - Small Scale Expression

Expression constructs for hDkk-3flag.long and hDkk-3flag.short were transfected into 293T cells using 10 µl of lipofectamine (GIBCO/BRL) and 2 µg of DNA per well of a 6-well plate of cells which were 70-80% confluent. After 5 hours at
20 37°C, cells were fed with 1ml of 20%FCS/DMEM. After incubation overnight at 37°C, cells were conditioned in 1ml OptiMEM for 48 hours at 37°C. Samples of supernatant and cell pellets were solubilized in boiling SDS-PAGE gel buffer, run out on a 4-20% SDS-PAGE gel, transferred to a nylon membrane and probed with the anti-FLAG monoclonal antibody M2. Samples from both supernatant and pellet samples showed
25 significant immunoreactivity within a molecular weight range of 40-65 kDa on autoradiographic film using a HRP conjugated secondary antibody and ECL detection reagents. Thus, both forms of hDkk-3 tested are secreted from 293T cells thereby confirming experimentally that hDkk-3 is a secreted protein. It should be noted that the molecular weights of both forms of hDkk-3 tested are greater than predicted from the
30 amino acid sequence, suggesting that the hDkk-3 proteins secreted by 293T cells may be glycosylated. This is consistent with the presence of four potential sites for N-linked

glycosylation in the hDkk-3 protein (*e.g.*, at about amino acids 96-99, 106-109, 121-124, and 204-207 of SEQ ID NO:2).

Deglycosylation of hDkk-3

5 Given the heterogenous nature of secreted human Dkk-3, the effect of N-Glycanase treatment on the mobility of secreted flag-tagged hDkk-3 was studied. Briefly, 1mL samples of 293T cell supernatants collected 72 hours after transfection with the appropriate constructs were incubated with 50μL anti-flag M2 agarose beads (Sigma) for 16hrs at 4°C. Beads were washed with PBS (pH7.4) containing,
10 sequentially, 0.1%, 0.05% and 0.01% Triton X-100. The beads were resuspended in 20μL of 20mM sodium phosphate, pH 7.5, 50mM EDTA, 0.02% sodium azide, (incubation buffer) together with 0.5% SDS, 5% 2-mercaptoethanol and boiled for 2 minutes. The supernatant was split into equal 10μL aliquots which were diluted with 10μL incubation
15 enzyme buffer (20mM Tris-HCl, 1mM EDTA, 50mM NaCl, 0.02% sodium azide pH 7.5) or with enzyme buffer alone as control. After 18 hours at 37°C, samples were boiled in equal volumes of SDS-PAGE buffer and analyzed by SDS-PAGE and Western blotting. For western analysis, samples were electroblotted onto PVDF (Novex) after SDS-PAGE on 4-20% gradient gels, probed with M2 anti-flag antibody (1:500, Sigma)
20 followed by HRP conjugated sheep anti-mouse IgG (1:5000, Amersham), developed with chemiluminescent reagents (Renaissance, Dupont) and exposed to autoradiography film (Biomax MR2 film, Kodak).

Utilizing the above-described methodology, it was determined that hDkk-3 protein displayed a significant increase in mobility following N-Glycanase treatment.
25 The major 45-65 kD form of soluble hDkk-3 was observed as two species of 45-55 and 40 kD following deglycosylation. This finding is consistent with the presence of multiple potential sites of N-linked glycosylation in the hDkk-3 protein. The reason for the heterogeneity of deglycosylated hDkk-3 reflects either proteolytic processing or incomplete removal of carbohydrate from one or more attachment sites. A 30 kD hDkk3
30 species was also observed in these experiments, the mobility of which was unaltered by

N-Glycanase treatment. This form was only observed after overnight incubation of the samples and may be a non-specific degradation product.

Large Scale hDkk-3 Protein Production

5 For scale-up of hDkk-3flag.long protein expression, 30 x 150mm plates of 293T cells at 70-80% confluence were transfected with 27 µg DNA, 100 µl lipofectamine in 18ml OptiMEM for 5 hours at 37°C. 18 ml of 10%FCS/DMEM was added to each plate and incubated overnight at 37°C. 24 hours after the start of transfection, transfection supernatant was aspirated and 35 mls OptiMEM was added to each plate and the plates
10 incubated at 37°C for 72 hours. Conditioned medium was harvested, spun at 4000 rpm for 30 min. at 4°C, and filtered through a 0.45 micron filter unit. 1100 ml was passed over a 1.6 x 10 cm anti-FLAG M2 affinity column pre-equilibrated in PBS pH7.4 buffer at a flow rate of 2.0 ml per minute. After washing with 200 ml of PBS pH 7.4 buffer, bound material was eluted by a step of 200 mM Glycine pH 3.0 buffer and 0.5 ml
15 fractions collected. Upon elution, a significant protein peak was detected by absorbance at 280nm. Samples corresponding to conditioned medium, flow through and eluted fractions were analyzed by Coomassie blue and silver stained SDS-PAGE and by western blot analysis as described above. Significant immunoreactivity within a molecular weight range of 40-65 kDa was detected in conditioned medium and eluted
20 fractions but not in the flow through sample, indicating that the secreted hDkk-3flag.long protein bound to the affinity column specifically and was eluted efficiently by the described conditions. Coomassie blue staining of SDS-PAGE gels suggested that the predominant immunoreactive protein constituted >90% of the protein present in the bound and eluted protein peak. Peak fractions of eluted protein were pooled and
25 dialysed against Phosphate Buffered Saline resulting in a 4 ml volume of recombinant hDkk-3flag.long protein at a concentration of approximately 1mg/ml.

Example 4: Isolation and Characterization of hDkk-4

In this example, the isolation and characterization of the gene encoding human
30 Dkk-4 (also referred to as "hDkk-4", "Cysteine Rich Secreted Protein-2", "CRSP-2" or "CRISPY-2") is described.

Isolation and Analysis of a Human Dkk-4 cDNA

To identify novel proteins related to hDkk-3, the human Dkk-3 amino acid sequence was used to search the dbEST database using TBLASTN (WashUversion, 2.0, 5 BLOSUM62 search matrix). A dbEST clone with accession number AA565546 was identified as having homology to a portion of the hDkk-3 cDNA. This clone was obtained from the IMAGE consortium and sequenced fully to define the entire hDkk-4 sequence depicted in Figure 2.

Determination of the hydrophobicity profile of human Dkk-4 having the amino 10 acid sequence set forth in SEQ ID NO:5 indicated the presence of a hydrophobic region from about amino acid 1 to about amino acid 19 of SEQ ID NO:5. Further analysis of the amino acid sequence SEQ ID NO:5 using a signal peptide prediction program predicted the presence of a signal peptide from about amino acid 1 to about amino acid 19 of SEQ ID NO:5. Accordingly, the mature hDkk-4 protein includes about 205 amino 15 acids spanning from about amino acid 20 to about amino acid 224 of SEQ ID NO:5.

Tissue Distribution of hDkk-4

hDkk-4 mRNA was undetectable by Northern analysis in all adult and fetal human tissues examined. Accordingly, a survey was performed of a cDNA library panel 20 by PCR with hDkk-4 specific PCR primers. Using such primers, products were identified in libraries prepared from cerebellum, activated human T-lymphocytes, lung and esophagus.

Secretion and Post-Translational Modification of human Dkk-4

25 Flag epitope-tagged human Dkk-4 protein was transiently overexpressed in 293T cells and analyzed as described previously for hDkk-3. Soluble hDkk-4 was consistently detected as three major immunoreactive species of approximately 40 kD [form (i)], 30-32 kD [form (ii)] and 15-17 kD [form (iii)]. Neither form (i), (ii) or (iii) was significantly affected by N-glycanase treatment, consistent with the absence of N- 30 glycosylation sites from the protein.

To determine the possible cause of heterogeneity in the size of secreted hDkk-4, Edman N-terminal sequencing of anti-flag affinity purified material corresponding to bands (i), (ii) and (iii) was performed. Briefly, flag-tagged Dkk-4 protein was isolated by passing the conditioned media over an M2-biotin (Sigma)/streptavidin Poros column (2.1 X 30 mm, PE Biosystems); the column was then washed with PBS, pH 7.4 and flag-tagged protein eluted with 200 mM glycine, pH 3.0. Eluted fractions with 280nm absorbance greater than background were analyzed by SDS-PAGE and western blot. Purified Dkk-4 protein bound to PVDF membrane after SDS-PAGE and electroblotting was sequenced for N-terminal amino acid analysis on a PE Applied Biosystems Model 494 Procise instrument using Edman-based chemistry protein sequencing. The amino acid residues were analyzed by HPLC (Spherogel micro PTH 3-micron column) and determined by separation and peak height as compared to standards.

The N-terminal sequence of band (i) was found to be XVLDFNNIRS (SEQ ID NO:34) which corresponds exactly to the predicted signal peptide cleavage site (between Ala-18 and Leu-19). Because the same band is identified by anti-flag antibodies, which recognize the C-terminal epitope tag, band (i) was thus identified as the full length, mature hDkk-4 protein. The band (iii) N-terminal sequence was found to be SQGRKGQEGS (SEQ ID NO:35) which corresponds to CRD-2 cleaved at the dibasic site Lys132/Lys133 (*e.g.*, Lys113/Lys114 of the mature protein following cleavage of the a 19 amino acid signal sequence or Lys 114/Lys 115 following cleavage of a 18 amino acid signal sequence). These data obtained for bands (i) and (iii) indicate clearly that hDkk4 is proteolytically processed by 293T cells, resulting in the release of CRD-2 (a 91 amino acid biologically-active fragment) from the full length protein.

Moreover, the three major species migrated similarly on SDS-PAGE conducted under either reducing or non-reducing conditions. Thus, each of the major C-terminal (anti-flag immunoreactive) hDkk-4 species exist as independent proteolytic fragments that are not covalently linked via disulfide bonds to other subunits or complex components when secreted from 293T cells.

Example 5: Isolation and Characterization of hDkk-1

In this example, the isolation and characterization of the gene encoding human Dkk-1 (also referred to as "hDkk-1", "Cysteine Rich Secreted Protein-3", "CRSP-3" or "CRISPY-3") is described.

Identification of a Human Dkk-1 cDNA

Searching a proprietary database of EST information using the sequence of hDkk-3, an hDkk-1 partial sequence was found corresponding to a clone from a human fetal kidney cDNA library having the identification code jthKb075a10. This clone was sequenced further and to define the entire hDkk-1 sequence depicted in Figure 3. DNA for the clone jthKb075a10 was deposited with the ATCC as Accession No. 98633.

hDkk-1 has a predicted signal peptide from about amino acid residue 1 to 20 of SEQ ID NO:8, cleavage of which results in a mature protein having 246 amino acid residues in length and corresponding to amino acid residues 21 to 266 of SEQ ID NO:8.

Tissue Distribution of hDkk-1

Northern blot analysis of various tissues including heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas was performed as previously described using a probe specific for hDkk-1. A ~1.8 kb hDkk1 mRNA was detected at high levels in human placenta, but not in other tissues tested.

Secretion and Post-Translational Modification of hDkk-1

Flag epitope-tagged human Dkk-1 protein was transiently overexpressed in 293T cells and analyzed as described previously. hDkk-1 was efficiently secreted from mammalian cells and was readily detected in conditioned medium of transfected cells. Mature secreted hDkk-1 migrated with a molecular weight of approximately 42-50 kD. Treatment with N-Glycanase had no significant effect on the mobility of soluble hDkk-1. Although hDkk-1 contains one potential site of N-linked glycosylation at its extreme C-terminus (*e.g.*, at amino acids 256-259 of SEQ ID NO8), this site is not conserved in *Xenopus* Dkk-1 (Glinka *et al.*, *supra*) and appears not to be a major site of carbohydrate addition in 293T cells.

Example 6: Isolation and Characterization of hDkk-2

In this example, the isolation and characterization of the gene encoding human Dkk-2 (also referred to as "hDkk-2", "Cysteine Rich Secreted Protein-4", "CRSP-4" or "CRISPY-4") is described.

5

Isolation of a Human Dkk-2 cDNA

Using the hDkk-3 sequence to query the dbEST database, a clone having similarity to a portion of hDkk-3 was identified having Accession No. W55979. This clone was subsequently obtained from the IMAGE consortium and sequenced to define a partial hDkk-2 sequence set forth as SEQ ID NO:10. This cDNA comprises a coding region from nucleotides 1-537, as well as 3' untranslated sequences (nucleotides 538 to 702). The coding region alone is set forth as SEQ ID NO:12. The predicted amino acid sequence corresponds to amino acids 1 to 179 of SEQ ID NO:11. A cDNA encoding full length hDkk-2 was isolated from a human fetal lung lambda Ziplox libraries by conventional plaque hybridization (Sambrook *et al.*, 1989) and fully sequenced. The full-length nucleotide sequence is set forth as SEQ ID NO:20 and the predicted amino acid sequence is set forth as SEQ ID NO:21. The coding region alone is set forth as SEQ ID NO:22. The predicted amino acid sequence corresponds to amino acids 1 to 259 of SEQ ID NO:21. DNA for the clone fthul33 was deposited with the ATCC as Accession No. _____. hDkk-2 has a predicted signal peptide from about amino acid residue 1 to 33 of SEQ ID NO:21, cleavage of which results in a mature protein having 226 amino acid residues in length and corresponding to amino acid residues 34 to 259 of SEQ ID NO:21.

Tissue Distribution of hDkk-2

Northern blot analysis of various tissues (*e.g.*, heart, brain, skeletal muscle, colon, thymus, spleen, kidney, liver, small intestine, placenta, lung, and peripheral blood leukocytes) was performed as previously described using a probe specific for hDkk-2. Of the tissues tested, hDkk-2 mRNA expression was highest in heart, brain, placenta, lung, and skeletal muscle. hDkk-2 transcripts of approximately 4.0 and 4.5 kb were observed.

Secretion and Post-translational Modification of hDkk-2

Flag epitope-tagged human Dkk-2 protein was transiently overexpressed in 293T cells and analyzed as described previously. Soluble hDkk-2 was detected as a major species of 15-17 kD, closely similar in size to form (iii) of hDkk-4. Additional minor forms of hDkk-2 were also observed in certain experiments in the range of 20-21kD. Deglycosylation of hDkk-2 was not studied since the protein sequence lacks potential N-glycosylation sites. By comparison with the data presented in Example 4 regarding the dibasic proteolytic cleavage site in the hDkk-4 protein sequences, it is predicted that the major 15-17 kD form of hDkk-2 detected in these experiments corresponds to CRD-2, as was the case for hDkk-4.

Example 7: Isolation of Soggy proteins

In this example, the isolation and characterization of the gene encoding human and murine Soggy-1 (also referred to as "Cysteine Rich Secreted Protein-N" or "CRISP-N") is described.

Identification of a Human and Murine Soggy-1 cDNAs

Human Soggy-1 was identified as a novel protein with similarity to the N-terminal domain of hDkk3. A human partial sequence was identified in the dbEST database for a clone having the accession number AA397836. This clone was obtained from the IMAGE collection and sequenced fully to define the entire human Soggy-1 sequence depicted in Figure 7. Two murine partial sequences were likewise identified in the dBEST database. The clones were obtained from the IMAGE consortium and sequenced. The entire murine Soggy-1 sequence is depicted in Figure 8. Human and murine Soggy cDNAs encode proteins of 242aa and 230aa, respectively, and are predicted to be secreted owing to the presence of N-terminal signal peptides. hSoggy-1 has a predicted signal peptide from about amino acid residue 1 to 30 of SEQ ID NO:14, cleavage of which results in a mature protein having 194 amino acid residues in length and corresponding to amino acid residues 31 to 224 of SEQ ID NO:14. mSoggy-1 has a predicted signal peptide from about amino acid residue 1 to 20 of SEQ ID NO:27,

cleavage of which results in a mature protein having 210 amino acid residues in length and corresponding to amino acid residues 21 to 230 of SEQ ID NO:27. Human and murine Soggy proteins display 59% overall identity although significant amino acid identities are seen beyond this domain that extend into the CRDs of Dkk-3 (Figure 10).

5 However, cysteine residues are not conserved within these domains and the residues shared by Soggy and Dkk-3 are poorly conserved in other Dkks indicating that the sequence relationship between these proteins is unique. Homology is most obvious within a 51 amino acid region in which 33% identity is observed between hSoggy, mSoggy, hDkk-3 and mDkk-3. This 51 amino acid domain is referred to herein as an

10 SGY domain. Human and mouse Soggy-1 proteins each possess 2 sites of potential N-linked glycosylation which are within the SGY domain and are also conserved with Dkk3. (e.g., NNLT, corresponding to amino acid residues 97-100 of SEQ ID NO:14 and NKTG corresponding to amino acid residues 112-115 of SEQ ID NO:14). In contrast to other Dkks, the C-terminal domain of Soggy-1 shows no similarity to other protein

15 sequences in the public databases nor does it contain any cysteine residues. Soggy was so named in view of its lack of CRDs compared to hDkk-3, which had been previously designated Cysteine Rich Secreted Protein-1 ("CRISPY-1").

Tissue Distribution of Soggy-1

20 To investigate Soggy-1 mRNA expression, a mouse cDNA probe was used on murine Northern blots. A 1kb mSoggy-1 mRNA was expressed at very high levels in testis and, interestingly, demonstrated transient expression during mouse embryogenesis. Soggy-1 mRNA, which was undetectable at day 7 of gestation, was transiently expressed at day 11 and day 15, after which the expression level declined to

25 undetectable levels. Thus, mSoggy-1 displays a developmentally regulated pattern of expression.

In situ analysis was performed as described in Example 1. For detection of murine Soggy-1, the following primers were used:

mSoggy forward 5'-ACCTGCAATGTGTCGACTGAG-3' (SEQ ID NO:32), and

30 *reverse* 5'-CACTTACAGCTGTTGGGATG-3' (SEQ ID NO:33).

Consistent with the Northern analysis, very high level expression of Soggy-1 mRNA was observed by *in situ* analysis in adult testis. Upon high magnification, Soggy-1 mRNA was found to be expressed at high levels in the spermatogenic epithelium of the seminiferous tubules and in the spermatogonia at various stages of development. A series of sagittal sections of mouse embryos from E13.5 - E 17.5 and post-natal day 1.5 pups were also analyzed. In E15.5 embryos, Soggy-1 mRNA transcripts were localized to the developing dorsal root ganglia (DRGs) and also found in the cartilage primordium of the nasal septum. Soggy-1 expression was also seen in the eye from E13.5 to E16.5, as observed for mDkk-3. Expression of Soggy-1 mRNA at various stages of development is consistent with the northern analysis described above and suggests that Soggy-1 may play a role in multiple stages of development.

Secretion and Post-Translational Modification of Soggy Proteins

Flag epitope-tagged human Soggy-1 protein was transiently overexpressed in 293T cells and analysed as previously described. hSoggy was efficiently secreted from transfected 293T cells and migrated with a molecular weight of approximately 40-50 kD. Given the heterogenous nature of secreted human Soggy-1, the effect of N-Glycanase treatment on the mobility of secreted flag-tagged hSoggy-1 was studied. hSoggy displayed a 5-10 kD decrease in apparent molecular weight after N-Glycanase treatment, consistent with the presence of 2 potential sites of N-glycosylation in the protein.

Example 8: Structure of the Dkk Family proteins and Dkk-Related Proteins

The amino acid and nucleotide homology between Dkk family members and Dkk-related proteins is set forth in the following tables. Where indicated, mDkk-1 and xDkk-1 correspond to a murine and *Xenopus* proteins set forth in Glinks *et al.*, *supra*, and having Accession Nos: AF030433 and AF030434, respectively. Likewise cDkk-3 has Accession No. D26311

- 135 -

Table II sets forth overall sequence identities as determined using the ALIGN program, (version 2.0) using a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4:

	hDkk-3	hDkk-4	hDkk-1	hDkk-2	mDkk-1	xDkk-1	CLFEST
hDkk-3	100	16.0	18.6	15.1	18.5	16.5	53.0
hDkk-4		100	33.7	35.2	32.6	33.7	16.2
hDkk-1			100	33.1	80.2	53.5	17.4
hDkk-2				100	30.5	33.7	12.5

5

Table III sets forth nucleic acid identities as determined using the using the Wilbur Lipman DNA alignment program, Ktuple: 3; Gap Penalty: 3; Window: 20:

	hDkk-3	hDkk-4	hDkk-1	hDkk-2	mDkk-1	xDkk-1	CLFEST
hDkk-3	100	30.0	37.2	34.7	31.5	45.4	58.8
hDkk-4		100	43.0	35.9	38.8	38.4	36.7
hDkk-1			100	59.3	66.4	53.7	32.1
hDkk-2				100	38.8	38.4	36.7

- 136 -

Table IV sets forth local amino acid identities as determined using the Smith-Waterman algorithm as implemented in the program Bestfit of the GCG package, with gap penalties of 8 for opening and 1 for extending:

hDkk-1								
mDkk-1	82							
xDkk-1	64	63						
hDkk-2	50	48	47					
hDkk-3	39	37	37	37				
mDkk-3	36	33	38	40	83			
cDkk-3	34	31	35	36	61	60		
hDkk-4	45	43	47	46	40	39	34	
	hDkk-1	mDkk-1	xDkk-1	hDkk-2	hDkk-3	mDkk-3	cDkk-3	hDkk-4

5

A multiple alignment of the amino acid sequences of hDkk-1, hDkk-2, hDkk-3, hDkk-4, mDkk-1, mDkk-3, xDkk-1, and cDkk-3 is shown in Figure 6. Predicted signal peptides are underlined, N-glycosylation sites are indicated by a thick bar, CRD-1 by an open box and CRD-2 by a shaded box. The proteolytic cleavage site within hDkk4 is indicated by a double asterisk. The domain structure of the full length human Dkk proteins of the present invention as well as human Soggy are schematically illustrated in Figure 9. Signal peptides (darkened boxes), Cysteine Rich Domain 1 ("CRD-1") (also referred to as the "amino-terminal cysteine-rich domain"), Cysteine Rich Domain 2 ("CRD-2") (also referred to as the "carboxy-terminal cysteine-rich domain"), the soggy domain (SGY) within hDkk-3 and hSoggy-1, and sites of N-glycosylation are indicated.

As demonstrated at least in Figures 6 and 9, human Dkks 1 through 4 each possess an N-terminal signal peptide and contain two conserved cysteine-rich domains (CRDs) separated by a linker or spacer region. Each CRD possesses 10 conserved cysteine residues. The second CRD has elsewhere been described as a colipase-like domain because the positions of the ten conserved cysteines in this domain have been shown to be closely similar to those in proteins of the colipase family (Aravind and Koonin, *supra*). Conservation of CRD-1 and CRD-2 suggests important functions for

20

- 137 -

these domains. In contrast to the CRDs, the linker or spacer region that joins CRD-1 and CRD-2 is highly variable between hDkks, being notably larger in hDkk-1, -2 and -4 (50-55aa) as compared to Dkk-3 (12aa). Four potential sites of N-linked glycosylation are present in hDkk3 and are conserved in chicken and mouse Dkk-3. These sites are not conserved in other Dkk family members. hDkk1 possesses one potential N-glycosylation site located close to the C-terminus of the protein which is conserved in murine Dkk-1 but not in *Xenopus* Dkk-1 (Fig. 6). In addition, each hDkk possesses several potential dibasic proteolytic cleavage sites, suggesting the proteins may be subject to post-translational processing. hDkk3 is the most divergent of the four human Dkks and possesses an extended N-terminal unique region which precedes CRD-1 and an extended C-terminal unique region which is highly acidic.

Example 9: Effects of hDkks and Soggy on Wnt-induced axis duplication in *Xenopus* embryos

This Example describes the functional activities of the hDkk and Soggy proteins of the present invention.

Xenopus embryo culture and RNA microinjections

Eggs were obtained from *Xenopus* females injected with 700 units of human chorionic gonadotropin, fertilized *in vitro* and cultured in 0.1 x MMR (Newport and Kirschner (1982) *Cell* 30:675-686). Embryonic stages were determined according to Nieuwkoop and Faber (1967) *Normal table of Xenopus laevis* (Daudin) Amsterdam: North Holland Publ. All cDNAs were subcloned into pCS2 vector (Rupp *et al.* (1984) *Genes & Development* 8:1311-1323), and capped mRNAs were synthesized *in vitro* as described (Krieg and Melton (1984) *Nucleic Acids Res.* 12:7057-7070, using the Message Machine kit (Ambion). The following plasmids were used as templates for mRNA synthesis: hDkk-1-pCS2, hDkk-2-pCS2, hDkk-3-pCS2, hDkk-4-pCS2, hSoggy-pCS2, Xwnt8 (Christian *et al.*, (1991) *Development* 111:1045-1055), Xwnt2B (Landesman and Sokol (1997) *Mech. Dev.* 61:1199-209), Xwnt3a (Wolda *et al.* (1993) *Dev. Biol.* 155:46-5), Xfz8-pXT7 (Itoh *et al.* (1998) *Mech. Devel.* 74:145-157), Xdsh-pXT7 (Sokol, *et al.* (1995) *Mech. Devel.* 74:145-157). Protein expression from all

- 138 -

pCS2-Dkk constructs was confirmed by *in vitro* transcription and translation (TNT, Promega). For secondary axis induction, a single ventral blastomere of 4- or 8-cell embryos was injected with 10nl of a solution containing 2-4pg of Xwnt8 mRNA, 2.5-5pg of Xwnt3a mRNA or 10pg of Xwnt2B mRNA as described (Itoh *et al.* (1995) *supra.*). The effect of Dkk RNAs was tested by coinjecting Wnt mRNAs with 2.5ng of hDkk mRNAs. For studies of Frizzled and Dhshevelled, 5ng Fz8 and 1ng Xdsh mRNAs were injected as indicated. After injections, embryos were cultured in 3% FicolI 400 (Pharmacia), 0.5xMMR. Secondary axes were scored at stage 35 as complete, when they contained anterior neuroectodermal derivatives including pronounced cement gland and eyes, and as partial, when the secondary neural tube with melanocytes, but without head structures, was apparent.

Inhibition of Secondary Axis Induction by hDkk-1 and hDkk-4 in *Xenopus* Embryos

hDkk-1 or hDkk-2 mRNAs were coinjected with Xwnt8 mRNA into single ventral blastomeres of 4- or 8-cell embryos. Injected embryos were cultured for 2 days and secondary axes were scored based on external morphology. Xwnt8 injected embryos displayed complete axis duplication, which was inhibited by co-injection with mRNAs encoding hDkk-1 and hDkk-4. To determine whether hDkks interacted with specific Wnt ligands, several different Wnts were assayed in combination with hDkk-1 or hDkk-4 for secondary axis formation. hDkk-1 and hDkk-4 inhibited axis duplication in response to Xwnt3a and Xwnt2b in addition to Xwnt8. hDkk-1 consistently demonstrated stronger inhibition of Wnt signaling than hDkk-4. Thus, hDkk-1 and hDkk -4 do not show any clear selectivity for the Wnt ligands used in this study. This compares to the FRPs, which also show little specificity with respect to their ability to inhibit Wnts (Leyns *et al.* (1997) *supra*; Wang *et al.* (1997) *supra*; Salic *et al.* (1997) *supra*; Mayr *et al.* (1997) *supra*; Finch *et al.* (1997) *supra*).

To investigate the mechanism by which hDkk-1 and hDkk-4 inhibit Wnt signaling, Dkk mRNAs were coinjected with Xdsh, a downstream component of the Wnt signaling pathway (Itoh *et al.* (1998) *supra*). hDkks-1 and -4 did not block secondary axis formation by Xdsh, indicating that Dkks function upstream of, or parallel with, Xdsh signaling. Similar findings have been reported previously for xDkk-1 (Glinka *et*

al. (1998) *supra*). It was also determined whether hDkks could antagonize signaling by *Xenopus* Frizzled-8 (Xfz8), which can also induce a secondary axis through Wnt signaling (Itoh *et al.* (1998) *supra*). Neither hDkk-1 or hDkk-4 inhibited the axis-inducing activity of Xfz8 mRNA. This data, taken together with the fact that hDkk-1 and hDkk-4 are secreted, indicate that Dkks antagonize Wnt signaling at a point upstream of Wnt receptors.

Assay for Inhibition of Secondary Axis Induction by hDkk-2, hDkk-3 and hSoggy-1 in *Xenopus* Embryos

hDkk-2, hDkk-3 or Soggy mRNAs were coinjected with Xwnt8 mRNA into single ventral blastomeres of 4- or 8-cell embryos and secondary axes were scored after two days as described for hDkk-1 and hDkk-4. Injection of mRNAs encoding hDkk-2, hDkk-3 or hSoggy-1 had no effect on Xwnt8-induced axis duplication. The ability of hDkk-2, hDkk-3 and hSoggy-1 to interact with specific Wnt ligands was also determined as described previously. hDkk-2, hDkk-3 and hSoggy-1 were inactive against each of the three Wnts tested. The lack of activity of hDkk-2, hDkk-3 and hSoggy-1 suggests that these proteins antagonize other members of the Wnt superfamily not tested here, or that they perform functions distinct from Wnt inhibition.

Example 10: Preparation of Antibodies Specific for hDkk and hSoggy Proteins

This example describes the making of polyclonal antibodies specific for hDkk-1, hDkk-4, hDkk-1, hDkk-2, and hSoggy-1.

Peptides were synthesized using Fmoc solid phase methodology utilizing MAP resin technology which increases the antigenic response (Tarn (1988) *Proc. Natl. Acad. Sci. USA* 85:5409-5413. For each protein, the peptides used for immunization are listed below:

hDkk-3

peptide #44 FREVEELMEDTQHKL

peptide #46 GSFMEEVRQELEDLE

hDkk-4

peptide #91 HAEGTTGHPVQENQP

- 140 -

hDkk-1

peptide #93 GNKYQTIDNYQPYPG

hDkk-2

peptide #56 GHYSNHDLDGWQNLGR

5 hSoggy-1

peptide #58 LQAIRDGLRKGTHKD

Peptides were designed to meet at least the following criteria: (1) not included within the cysteine-rich domain; (2) not including an N-glycosylation site; and (3) hydrophilic (*e.g.*, solvent exposed).

Antibodies were generated in New Zealand white rabbits over a 10-week period. The immunogen includes KLH-peptide emulsified by mixing with an equal volume of Freund's Adjuvant, and injected into three subcutaneous dorsal sites, for a total of 0.1mg peptide per immunization. Animals were bled from the articular artery. The blood was allowed to clot and the serum collected by centrifugation. The serum is stored at -20°C.

For purification, peptide antigens were immobilized on an activated support. Antisera was passed through the sera column and then washed. Specific antibodies were eluted *via* a pH gradient, collected and stored in a borate buffer (0.125M total borate) at ~0.25mg/ml. The anti-peptide titers were determined using ELISA methodology with free peptide bound in solid phase (1µg/well). Detection was obtained using biotinylated anti-rabbit IgG, HRP-SA conjugate, and ABTS.

All antibodies performed well in ELISA assays. Anti-peptide #44, #46, and #58 are particularly useful for detection of hDkk-3 and hSoggy-1, respectively, as determined by western blotting of supernatants from hDkk-3- and hSoggy-1-transfected 293T cells.

The Dkk family comprises a novel family of secreted proteins which to date includes hDkk-1, hDkk-2, hDkk-3, hDkk-4, xDkk-1, mDkk-1 and cDkk-1. Structurally, Dkks 1-4 are related by several conserved features. Firstly, all four

proteins are secreted proteins. Secondly, Dkks 1-4 each possess two distinct cysteine rich domains. Each domain contains 10 conserved cysteine residues, and these domains are highly conserved between family members. The C-terminal cysteine rich domain, referred to as CRD-2, bears significant similarity to proteins of the colipase family and sequence conservation among the Dkks is greatest within CRD-2 (Aravind and Koonin, *supra*). This may reflect a need for Dkks to interact with lipids in order to regulate Wnt function, since Wnt proteins remain tightly associated with the cell surface.

Despite the similarities between Dkks 1-4, notable differences between these family members appear with regard to their mRNA expression patterns. In adult human tissues hDkk-1 and hDkk-4 showed highly restricted mRNA expression patterns while hDkk-2 and hDkk-3 are more widely expressed. Murine Dkk-3 mRNA was found to be restricted to the myocytes of the atria in the heart, neurons of the cortex and hippocampus in the brain and also to the retinal neurons and lens epithelium in the eye. Such specific expression patterns reflect localized action of the Dkks as regulators of Wnt activity and/or that of other signaling molecules. Different Wnt family members have been shown to have divergent patterns of mRNA expression in adult and embryonic mammalian tissues. For example, murine Wnts 4, 7a and 7b are expressed in brain and lung, whereas Wnt6 is highly expressed in testis (Gavin *et al.*, 1990). Wnts 5b and 13 are more broadly expressed (Gavin *et al.* (1990) *supra*; Katoh *et al.* (1996) *supra*). Although Wnts have been studied mostly in the context of their roles in embryonic development and tumorigenesis, the expression of many family members in normal adult human and mouse tissues, together with their regulators such as the Dkks, suggests that these signaling proteins play important roles in normal tissue homeostasis.

Marked differences in the post-translational processing of different human Dkk proteins was also observed. hDkk-3 is secreted from 293T cells as a heterogeneously glycosylated protein, whereas Dkk-1, 2 and 4 proteins show no evidence of glycosylation. This is consistent with sequence analysis that identifies 4 potential sites of N-glycosylation in the hDkk-3 protein but no sites in either hDkk-2 or hDkk-4. A single putative site in hDkk-1 does not appear to be utilized in 293T cells and may

- 142 -

well not be a significant site for N-linked carbohydrate addition in view of its C-terminal location and lack of conservation with xDkk-1. C-terminal proteolysis of hDkk4 in 293T cells was also characterized. Dkk proteins contain multiple potential sites of proteolytic processing. Secreted hDkk-4 was consistently detected as three major C-terminal fragments. N-terminal sequencing identified two of these as mature, full length hDkk4 and CRD-2, which was derived from the full length protein by a specific proteolytic event at lysines 132 and 133. Thus, the hDkk-4 CRD-2 is released from the full length protein upon expression in 293T cells. Similar processing of hDkk4 in COS cells has been observed.

Within the Dkk family, Dkks 1, 2 and 4 display closest similarity whereas Dkk-3 is more distantly related. For example, Dkk-3 contains a linker region connecting CRD-1 and CRD-2 which is shorter than in other Dkks. Dkk-3 also possesses extended N-and C-terminal regions compared to other Dkks. Within the Dkk-3 N-terminal unique region, a distinct domain has been identified that is also found in Soggy (the SGY domain). The SGY domains of human and mouse Soggy-1 and Dkk-3 proteins contain two conserved sites of N-linked glycosylation. Protein expression studies confirm that, like hDkk3, hSoggy is secreted as a glycoprotein. Murine Soggy-1 is highly expressed in adult testis and also displays a highly restricted mRNA expression in E15 -E16 mouse embryos, being localized mainly to the cartilage primordia within the developing vertebrae/developing dorsal root ganglia. Soggy mRNA was also detected at high levels in the developing eye, similar to Dkk-3. This developmentally regulated pattern of Soggy expression overlaps with that of Dkk-3 suggesting that Soggy may play a role in the regulation of Dkk-3 function.

25 Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

What is claimed:

1. An isolated nucleic acid molecule selected from the group consisting of:
 - a) a nucleic acid molecule comprising a nucleotide sequence which
5 is at least 60% homologous to a nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:20, SEQ ID NO:22, the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide
10 sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a complement thereof;
 - b) a nucleic acid molecule comprising a fragment of at least 1000 nucleotides of a nucleic acid comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID
15 NO:9, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:20, SEQ ID NO:22, the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a complement thereof;
 - c) a nucleic acid molecule which encodes a polypeptide comprising
20 an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21, the amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the amino acid
25 sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the amino acid sequence encoded by the DNA insert of the plasmid deposited with the ATCC as Accession Number _____;
 - d) a nucleic acid molecule which encodes a fragment of a
30 polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number

- 144 -

98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, wherein the fragment comprises at least 15 contiguous amino acid residues of the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with the ATCC as Accession Number _____; and

e) a nucleic acid molecule which encodes a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, wherein the nucleic acid molecule hybridizes to a nucleic acid molecule comprising SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:20, SEQ ID NO:22, the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the DNA insert of the plasmid deposited with Accession Number _____ under stringent conditions.

- 145 -

2. The isolated nucleic acid molecule of claim 1 which is selected from the group consisting of:

- 5 a) a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:20, SEQ ID NO:22, the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a complement thereof;
- 10 and
- b) a nucleic acid molecule which encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with the ATCC as Accession Number _____.
- 15

3. The nucleic acid molecule of claim 1 further comprising vector nucleic acid sequences.

20

4. The nucleic acid molecule of claim 1 further comprising nucleic acid sequences encoding a heterologous polypeptide.

25 5. A host cell which contains the nucleic acid molecule of claim 1.

6. The host cell of claim 5 which is a mammalian host cell.

7. A non-human mammalian host cell containing the nucleic acid molecule of claim 1.

30

8. An isolated polypeptide selected from the group consisting of:

a) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, wherein the fragment comprises at least 15

contiguous amino acids of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with the ATCC as Accession Number _____;

b) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule comprising SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:20, SEQ ID NO:22, the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the DNA insert of the plasmid deposited with the ATCC as Accession Number _____ under stringent conditions;

c) a polypeptide which is encoded by a nucleic acid molecule comprising a nucleotide sequence which is at least 60% homologous to a nucleic

acid comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:20, SEQ ID NO:22, the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the DNA insert of the plasmid deposited with ATCC as Accession Number _____; and

d) a polypeptide comprising an amino acid sequence which is at least 60% homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____.

9. The isolated polypeptide of claim 8 comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____.

10. The polypeptide of claim 8 further comprising heterologous amino acid sequences.

11. An antibody which selectively binds to a polypeptide of claim 8.

12. A method for producing a polypeptide selected from the group consisting of:

- 5 a) a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____;
- 10 b) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the
- 15 nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide
- 20 encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____; and
- 25 c) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, wherein the polypeptide is
- 30 encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule comprising SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID

NO:7, SEQ ID NO:9, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:20, SEQ ID NO:22, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the
5 nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____ under stringent conditions;

comprising culturing the host cell of claim 5 under conditions in which the nucleic acid molecule is expressed.

10

13. A method for detecting the presence of a polypeptide of claim 8 in a sample comprising:

- a) contacting the sample with a compound which selectively binds to the polypeptide; and
- 15 b) determining whether the compound binds to the polypeptide in the sample to thereby detect the presence of a polypeptide of claim 8 in the sample.

14. The method of claim 13, wherein the compound which binds to the
20 polypeptide is an antibody.

15. A kit comprising a compound which selectively binds to a polypeptide of claim 8 and instructions for use.

25 16. A method for detecting the presence of a nucleic acid molecule in claim 1 in a sample comprising:

- a) contacting the sample with a nucleic acid probe or primer which selectively hybridizes to the nucleic acid molecule; and
- 30 b) determining whether the nucleic acid probe or primer binds to a nucleic acid molecule in the sample to thereby detect the presence of a nucleic acid molecule of claim 1 in the sample.

- 150 -

17. The method of claim 16, wherein the sample comprises mRNA molecules and is contacted with a nucleic acid probe.

18. A kit comprising a compound which selectively hybridizes to a nucleic acid molecule of claim 1 and instructions for use.

19. A method for identifying a compound which binds to a polypeptide of claim 8 comprising:

- a) contacting the polypeptide, or a cell expressing the polypeptide with a test compound; and
- b) determining whether the polypeptide binds to the test compound.

20. The method of claim 19, wherein the binding of the test compound to the polypeptide is detected by a method selected from the group consisting of:

- a) detection of binding by direct detection of test compound/polypeptide binding;
- b) detection of binding using a competition binding assay; and
- c) detection of binding using an assay for Dkk activity.

21. A method of modulating the activity of a polypeptide of claim 8 comprising contacting the polypeptide or a cell expressing the polypeptide with a compound which binds to the polypeptide in a sufficient concentration to modulate the activity of the polypeptide.

22. A method for identifying a compound which modulates the activity of a polypeptide of claim 8 comprising:

- a) contacting a polypeptide of claim 8 with a test compound; and
- b) determining the effect of the test compound on the activity of the polypeptide to thereby identify a compound which modulates the activity of the polypeptide.

Figure 1A

GGCACGAGGGGGCGGCGGCTGCGGGCGCAGAGCGGAG										M	Q	R	L	G	A	T	L	L	C	
										ATG	CAG	CGG	CTT	GGC	GCC	ACC	CTG	CTG	TGC	67
L	L	L	A	A	A	V	P	T	A	P	A	P	A	P	T	A	T	S	A	30
CTG	CTG	CTG	GCG	GCG	GCG	GTC	CCC	ACG	GCC	CCC	GCG	CCC	GCT	CCG	ACG	GCG	ACC	TCG	GCT	127
P	V	K	P	G	P	A	L	S	Y	P	Q	E	E	A	T	L	N	E	M	50
CCA	GTC	AAG	CCC	GGC	CCG	GCT	CTC	AGC	TAC	CCG	CAG	GAG	GAG	GCC	ACC	CTC	AAT	GAG	ATG	187
F	R	E	V	E	E	L	M	E	D	T	Q	H	K	L	R	S	A	V	E	70
TTC	CGC	GAG	GTT	GAG	GAA	CTG	ATG	GAG	GAC	ACG	CAG	CAC	AAA	TTG	CGC	AGC	GCG	GTG	GAA	247
E	M	E	A	E	E	A	A	A	K	A	S	S	E	V	N	L	A	N	L	90
GAG	ATG	GAG	GCA	GAA	GAA	GCT	GCT	GCT	AAA	GCA	TCA	TCA	GAA	GTG	AAC	CTG	GCA	AAC	TTA	307
P	P	S	Y	H	N	E	T	N	T	D	T	K	V	G	N	N	T	I	H	110
CCT	CCC	AGC	TAT	CAC	AAT	GAG	ACC	AAC	ACA	GAC	ACG	AAG	GTT	GGA	AAT	AAT	ACC	ATC	CAT	367
V	H	R	E	I	H	K	I	T	N	N	Q	T	G	Q	M	V	F	S	E	130
GTG	CAC	CGA	GAA	ATT	CAC	AAG	ATA	ACC	AAC	AAC	CAG	ACT	GGA	CAA	ATG	GTC	TTT	TCA	GAG	427
T	V	I	T	S	V	G	D	E	E	G	R	R	S	H	E	C	I	I	D	150
ACA	GTT	ATC	ACA	TCT	GTG	GGA	GAC	GAA	GAA	GGC	AGA	AGG	AGC	CAC	GAG	TGC	ATC	ATC	GAC	487
E	D	C	G	P	S	M	Y	C	Q	F	A	S	F	Q	Y	T	C	Q	P	170
GAG	GAC	TGT	GGG	CCC	AGC	ATG	TAC	TGC	CAG	TTT	GCC	AGC	TTC	CAG	TAC	ACC	TGC	CAG	CCA	547
C	R	G	Q	R	M	L	C	T	R	D	S	E	C	C	G	D	Q	L	C	190
TGC	CGG	GGC	CAG	AGG	ATG	CTC	TGC	ACC	CGG	GAC	AGT	GAG	TGC	TGT	GGA	GAC	CAG	CTG	TGT	607
V	W	G	H	C	T	K	M	A	T	R	G	S	N	G	T	I	C	D	N	210
GTC	TGG	GGT	CAC	TGC	ACC	AAA	ATG	GCC	ACC	AGG	GGC	AGC	AAT	GGG	ACC	ATC	TGT	GAC	AAC	567
Q	R	D	C	Q	P	G	L	C	C	A	F	Q	R	G	L	L	F	P	V	230
CAG	AGG	GAC	TGC	CAG	CCG	GGG	CTG	TGC	TGT	GCC	TTC	CAG	AGA	GGC	CTG	CTG	TTC	CCT	GTG	727
C	T	P	L	P	V	E	G	E	L	C	H	D	P	A	S	R	L	L	D	250
TGC	ACA	CCC	CTG	CCC	GTG	GAG	GGC	GAG	CTT	TGC	CAT	GAC	CCC	GCC	AGC	CGG	CTT	CTG	GAC	787
L	I	T	W	E	L	E	P	D	G	A	L	D	R	C	P	C	A	S	G	270
CTC	ATC	ACC	TGG	GAG	CTA	GAG	CCT	GAT	GGA	GCC	TTG	GAC	CGA	TGC	CCT	TGT	GCC	AGT	GGC	347
L	L	C	Q	P	H	S	H	S	L	V	Y	V	C	K	P	T	F	V	G	290
CTC	CTC	TGC	CAG	CCC	CAC	AGC	CAC	AGC	CTG	GTG	TAT	GTG	TGC	AAG	CCG	ACC	TTC	GTG	GGG	327
S	R	D	Q	D	G	E	I	L	L	P	R	E	V	P	D	D	Y	E	V	310
AGC	CGT	GAC	CAA	GAT	GGG	GAG	ATC	CTG	CTG	CCC	AGA	GAG	GTC	CCC	GAT	GAG	TAT	GAA	GTT	567
G	S	F	M	E	E	V	R	Q	E	L	E	D	L	E	R	S	L	T	E	330
GGC	AGC	TTC	ATG	GAG	GAG	GTG	CGC	CAG	GAG	CTG	GAG	GAC	CTG	GAG	AGG	AGC	CTG	ACT	GAA	1027
E	M	A	L	G	E	P	A	A	A	A	A	A	L	L	G	G	E	E	I	350
GAG	ATG	GCG	CTG	GGG	GAG	CCT	GCG	GCT	GCC	GCC	GCT	GCA	CTG	CTG	GGA	GGG	GAA	GAG	ATT	1087
★																				351
TAG																				1090

2/15

Figure 1B

ATCTGGACCAGGCTGTGGGTAGATGTGCAATAGAAATAGCTAATTTATTTCCCCANGTGTGTGCTTTAAGCGTGGGCTG 1169
ACCAGGCTTCTTCCTACATCTTCTTCCCAGTAAGTTTCCCCTCTGGCTTGACAGCATGAGGTGTTGTGCATTTGTTTCAG 1248
CTCCCCCAGGCTGTTCTCCAGGCTTCACAGTCTGGTGCTTGGGAGAGTCAGGCAGGGTTAAACTGCAGGAGCAGTTTGC 1327
CACCCCTGTCCAGATTATTGGCTGCTTTGCCTCTACCAGTTGGCAGACAGCCGTTTGTCTACATGGCTTTGATAATTG 1406
TTTGAGGGGAGGAGATGGAAACAATGTGGAGTCTCCCTCTGATTGGTTTTGGGGAAATGTGGAGAAGAGTGCCCTGCTT 1485
TGCAAACATCAACCTGGCAAAAATGCAACAAATGAATTTTCCACGCAGTTCTTTCCATGGGCATAGGTAAGCTGTGCCT 1564
TCAGCTGTTGCAGATGAAATGTTCTGTTCAACCCTGCATTACATGTGTTTATTCATCCAGCAGTGTGCTCAGCTCCTAC 1643
CTCTGTGCCAGGGCAGCATTTTTCATATCCAAGATCAATTCCCTCTCTCAGCACAGCCTGGGGAGGGGGTCATTGTTCTC 1722
CTCGTCCATCAGGGATTTTCAGAGGCTCAGAGACTGCAAGCTGCTTGCCCAAGTCACACAGCTAGTGAAGACCAGAGCAG 1801
TTTCATCTGGTTGTGACTCTAAGCTCAGTGCTCTCTCCACTACCCACACCAGCCTTGGTGCCACCAAAAGTGCTCCCC 1880
AAAAGGAAGGAGAATGGGATTTTTCTTTTGAGGCATGCACATCTGGAATTAAGGTCAAACCTAATTCTCACATCCCTCTA 1959
AAAGTAACTACTGTTAGGAACAGCAGTGTCTCACAGTGTGGGGCAGCCGTCCTTCTAATGAAGACAATGATATTGAC 2038
ACTGTCCCTCTTTGGCAGTTGCATTAGTAACTTTGAAAGGTATATGACTGAGCGTAGCATAACAGGTTAACCTGCAGAAA 2117
CAGTACTTAGGTAATTGTAGGGCGAGGATTATAAATGAAATTTGCAAAATCACTTAGCAGCAACTGAAGACAATTATCA 2196
ACCACGTGGAGAAAATCAAACCGAGCAGGGCTGTGTGAAACATGGTTGTAATATGCGACTGCGAACACTGAACTCTACG 2275
CCACTCCACAAATGATGTTTTTCAGGTGTCATGGACTGTTGCCACCATGTATTCATCCAGAGTTCTTAAAGTTTAAAGTT 2354
GCACATGATTGTATAAGCATGCTTTCTTTGAGTTTTAAATTATGTATAAACATAAGTTGCATTTAGAAATCAAGCATAA 2433
ATCACTTCAACTGCTAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA 2479

3/15

Figure 2

GAATTCGGCACGAGAGACGACGTGCTGAGCTGCCAGCTTAGTGGAAGCTCTGCTCTGGGTGGAGAGCAGCCTCGCTTTG 79

GTGACGCACAGTGCTGGGACCCTCCAGGAGCCCCGGGATTGAAGG M V A A V L L G 8
 148

L S W L C S P L G A L V L D F N N I R S 28
 CTG AGC TGG CTC TGC TCT CCC CTG GGA GCT CTG GTC CTG GAC TTC AAC AAC ATC AGG AGC 208

S A D L H G A R K G S Q C L S D T D C N 48
 TCT GCT GAC CTG CAT GGG GCC CGG AAG GGC TCA CAG TGC CTG TCT GAC ACG GAC TGC AAT 268

T R K F C L Q P R D E K P F C A T C R G 68
 ACC AGA AAG TTC TGC CTC CAG CCC CGC GAT GAG AAG CCG TTC TGT GCT ACA TGT CGT GGG 328

L R R R C Q R D A M C C P G T L C V N D 98
 TTG CGG AGG AGG TGC CAG CGA GAT GCC ATG TGC TGC CCT GGG ACA CTC TGT GTG AAC GAT 388

V C T T M E D A T P I L E R Q L D E Q D 108
 GTT TGT ACT ACG ATG GAA GAT GCA ACC CCA ATA TTA GAA AGG CAG CTT GAT GAG CAA GAT 448

G T H A E G T T G H P V Q E N Q P K R K 128
 GGC ACA CAT GCA GAA GGA ACA ACT GGG CAC CCA GTC CAG GAA AAC CAA CCC AAA AGG AAG 508

P S I K K S Q G R K G Q E G E S C L R T 148
 CCA AGT ATT AAG AAA TCA CAA GGC AGG AAG GGA CAA GAG GGA GAA AGT TGT CTG AGA ACT 568

F D C G P G L C C A R H F W T K I C K P 168
 TTT GAC TGT GGC CCT GGA CTT TGC TGT GCT CGT CAT TTT TGG ACG AAA ATT TGT AAG CCA 628

V L L E G Q V C S R R G H K D T A Q A P 188
 GTC CTT TTG GAG GGA CAG GTC TGC TCC AGA AGA GGG CAT AAA GAC ACT GCT CAA GCT CCA 688

E I F Q R C D C G P G L L C R S Q L T S 208
 GAA ATC TTC CAG CGT TGC GAC TGT GGC CCT GGA CTA CTG TGT CGA AGC CAA TTG ACC AGC 748

N R Q H A R L R V C Q K I E K L * 225
 AAT CGG CAG CAT GCT CGA TTA AGA GTA TGC CAA AAA ATA GAA AAG CTA TAA 788

ATATTTCAAAATAAAGAAGAATCCACATTGCAAAAAAAAAAAAAAAAAAAAA 343

Figure 3

4/15

GTCGACCCACGCGTCCGCGGACGCGTGGGCGGCACGGTTTCGTGGGGACCCAGGCTTGCAAAGTGACGGTCATTTTCTC 79
 TTTCTTTCTCCCTCTTGAGTCCTTCTGAG ATG ATG GCT CTG GGC GCA GCG GGA GCT ACC CGG GTC 12
 144
 F V A M V A A A L G G H P L L G V S A T 32
 TTT GTC GCG ATG GTA GCG GCG GCT CTC GGC GGC CAC CCT CTG CTG GGA GTG AGC GCC ACC 204
 L N S V L N S N A I K N L P P P L G G A 52
 TTG AAC TCG GTT CTC AAT TCC AAC GCT ATC AAG AAC CTG CCC CCA CCG CTG GGC GGC GCT 264
 A G H P G S A V S A A P G I L Y P G G N 72
 GCG GGG CAC CCA GGC TCT GCA GTC AGC GCC GCG CCG GGA ATC CTG TAC CCG GGC GGG AAT 324
 K Y Q T I D N Y Q P Y P C A E D E E C G 92
 AAG TAC CAG ACC ATT GAC AAC TAC CAG CCG TAC CCG TGC GCA GAG GAC GAG GAG TGC GGC 384
 T D E Y C A S P T R G G D A G V Q I C L 112
 ACT GAT GAG TAC TGC GCT AGT CCC ACC CGC GGA GGG GAC GCA GGC GTG CAA ATC TGT CTC 444
 A C R K R R K R C M R H A M C C P G N Y 132
 GCC TGC AGG AAG CGC CGA AAA CGC TGC ATG CGT CAC GCT ATG TGC TGC CCC GGG AAT TAC 504
 C K N G I C V S S D Q N H F R G E I E E 152
 TGC AAA AAT GGA ATA TGT GTG TCT TCT GAT CAA AAT CAT TTC CGA GGA GAA ATT GAG GAA 564
 T I T E S F G N D H S T L D G Y S R R T 172
 ACC ATC ACT GAA AGC TTT GGT AAT GAT CAT AGC ACC TTG GAT GGG TAT TCC AGA AGA ACC 624
 T L S S K M Y H T K G Q E G S V C L R S 192
 ACC TTG TCT TCA AAA ATG TAT CAC ACC AAA GGA CAA GAA GGT TCT GTT TGT CTC CGG TCA 684
 S D C A S G L C C A R H F W S K I C K P 212
 TCA GAC TGT GCC TCA GGA TTG TGT TGT GCT AGA CAC TTC TGG TCC AAG ATC TGT AAA CCT 744
 V L K E G Q V C T K H R R K G S H G L E 232
 GTC CTG AAA GAA GGT CAA GTG TGT ACC AAG CAT AGG AGA AAA GGC TCT CAT GGA CTA GAA 804
 I F Q R C Y C G E G L S C R I Q K D H H 252
 ATA TTC CAG CGT TGT TAC TGT GGA GAA GGT CTG TCT TGC CGG ATA CAG AAA GAT CAC CAT 864
 Q A S N S S R L H T C Q R H * 267
 CAA GCC AGT AAT TCT TCT AGG CTT CAC ACT TGT CAG AGA CAC TAA 909
 ACCAGCTATCCAAATGCAGTGAACCTCCTTTTATATAATAGATGCTATGAAAACCTTTTATGACCTTCATCAACTCAATC 988
 CTAAGGATATACAAGTTCTGTGGTTTCAGTTAAGCATTCCAATAACACCTTCCAAAAACCTGGAGTGTAAGAGCTTTGT 1067
 TTCTTTATGGAACCTCCCCTGTGATTGCAGTAAATTACTGTATTGTAAATTCTCAGTGTGGCACTTACCTGTAAATGCAA 1146
 TGAAACTTTTAATTATTTTCTAAAGGTGCTGCACTGCCTATTTTCTCTTGTATGTAAATTTTGTACACATTGAT 1225
 TGTTATCTTGACTGACAAATATTCTATATTGAACTGAAGTAAATCATTTTCAGCTTATAGTTCTTAAAGCATAACCCCTT 1304
 TACCCCATTTAATTCTAGAGTCTAGAACGCAAGGATCTCTTGAATGACAAATGATAGGTACCTAAAAATGTAACATGAA 1383
 AATACTAGCTTATTTTCTGAAATGTACTATCTTAATGCTTAAATTATATTTCCCTTTAGGCTGTGATAGTTTTTGAAAT 1462
 AAAATTTAACATTTAATATCATGAAATGTTATAAGTAGACATAAAAAAAAAAAAAAAAAAAGGGCGGCCGC 1536

Figure 4A

5/15

GTCGACCCACGCGTCCGGCGGGAGCCCGCGGCGAGCGTAGCGCAAGTCCGCTCCCTAGGCATCGCTGCGCTGGCAGCGA 79
 TTCGCTGTCTCTTGTGAGTCAGGGGACAACGCTTCGGGGCAACTGTGAGTGCGCGTGTGGGGGACCTCGATTCTCTTCA 158
 GATCTCGAGGATTCGGTCCGGGGACGTCTCCTGATCCCCCTACTAAAGCGCCTGCTAACTTTGAAAAGGAGCACTGTGTC 237
 CTGCAAAGTTTGACACATAAAGGATAGGAAAAGAGAGGAGAGAAAAGCAACTGAGTTGAAGGAGAAGGAGCTGATGCGG 316
 GCCTCCTGATCAATTAAGAGGAGAGTTAAACCGCCGAGATCCCGGCGGGACCAAGGAGGTGCGGGGCAAGAAGGAACGG 395
 AAGCGGTGCGATCCACAGGGCTGGGTTTTCTTGACCTTGGGTACGCCTCCTTGGCGAGAAAGCGCCTCGCATTGTGAT 474
 TGCTTCCAGTTATTGCAGAACTTCTGTCTTGGTGGAGAAGCGGGTCTCGCTTGGGTTCGGCTAATTTCTGTCTGAGG 553
 CGTGAGACTGAGTTCATAGGGTCTTGGGTCCCCGAACCAGGAAGGGTTGAGGGAACACAATCTGCAAGCCCCGCGACC 632
 CAAGTGAGGGGCCCCGTGTTGGGGTCTCCTCCCTCCCTTTGCATTCCCACCCCTCCGGGCTTTGCGTCTTCTGGGGACCC 711
 CCTCGCCGGGAG ATG GCC GCG TTG ATG CGG AGC AAG GAT TCG TCC TGC TGC CTG CTC CTA CTG 774
 A A V L M V E S S Q I G S S R A K L N S 37
 GCC GCG GTG CTG ATG GTG GAG AGC TCA CAG ATC GGC AGT TCG CGG GCC AAA CTC AAC TCC 834
 I K S S L G G E T P G Q A A N R S A G M 57
 ATC AAG TCC TCT CTG GGC GGG GAG AC G CCT GGT CAG GCC GCG AAT CGA TCT GCG GGC ATG 894
 Y Q G L A F G G S K K G K N L G Q A Y P 77
 TAC CAA GGA CTG GCA TTC GGC GGC AGT AAG AAG GGC AAA AAC CTG GGG CAG GCC TAC CCT 954
 C S S D K E C E V G R Y C H S P H Q G S 97
 TGT AGC AGT GAT AAG GAG TGT GAA GTT GGG AGG TAT TGC CAC AGT CCC CAC CAA GGA TCA 1014
 S A C M V C R R K K R C H R D G M C C 117
 TCG GCC TGC ATG GTG TGT CGG AGA AAA AAG AAG CGC TGC CAC CGA GAT GGC ATG TGC TGC 1074
 P S T R C N N G I C I P V T E S I L T P 137
 CCC AGT ACC CGC TGC AAT AAT GGC ATC TGT ATC CCA GTT ACT GAA AGC ATC TTA ACC CCT 1134
 H I P A L D G T R H R D R N H G H Y S N 157
 CAC ATC CCG GCT CTG GAT GGT ACT CGG CAC AGA GAT CGA AAC CAC GGT CAT TAC TCA AAC 1194
 H D L G W Q N L G R P H T K M S H I K G 177
 CAT GAC TTG GGA TGG CAG AAT CTA GGA AGA CCA CAC ACT AAG ATG TCA CAT ATA AAA GGG 1254
 H E G D P C L R S S D C I E G F C C A R 197
 CAT GAA GGA GAC CCC TGC CTA CGA TCA TCA GAC TGC ATT GAA GGG TTT TGC TGT GCT CGT 1314
 H F W T K I C K P V L H Q G E V C T K Q 217
 CAT TTC TGG ACC AAA ATC TGC AAA CCA GTG CTC CAT CAG GGG GAA GTC TGT ACC AAA CAA 1374
 R K K G S H G L E I F Q R C D C A K G L 237
 CGC AAG AAG GGT TCT CAT GGG CTG GAA ATT TTC CAG CGT TGC GAC TGT GCG AAG GGC CTG 1434
 S C K V W K D A T Y S S K A R L H V C Q 257
 TCT TGC AAA GTA TGG AAA GAT GCC ACC TAC TCC TCC AAA GCC AGA CTC CAT GTG TGT CAG 1494
 K I * 260
 AAA ATT TGA 1503
 TCACCATTGAGGAACATCATCAATTGCAGACTGTGAAGTTGTGTATTTAATGCATTATAGCATGGTGGAAAATAAGGTT 1582
 CAGATGCAGAAGAATGGCTAAAATAAGAAACGTGATAAGAATATAGATGATCACAAAAAGGGAGAAAGAAAACATGAAC 1661
 TGAATAGATTAGAATGGGTGACAAATGCAGTGCAGCCAGTGTTCATTATGCAACTTGTCTATGTAAATAATGTACAC 1740

6/15

Figure 4B

ATTTGTGGAAATGCTATTATTAAGAGAACAAGCACACAGTGGAATTACTGATGAGTAGCATGTGACTTTCCAAGAGT 1819
TTAGGTTGTGCTGGAGGAGAGGTTTCCTTCAGATTGCTGATTGCTTATACAAATAACCTACATGCCAGATTTCTATTCA 1898
ACGTTAGAGTTTAACAAAATACTCCTAGAAATACTTGTTATACAATAGGTTCTAAAAATAAAATTGCTAAACAAGAAAT 1977
GAAAACATGGAGCATTGTTAATTTACAACAGAAAATTACCTTTTGATTTGTAACACTACTTCTGCTGTTCAATCAAGAG 2056
TCTTGGTAGATAAGAAAAAATCAGTCAATATTTCCAAATAATTGCAAAATAATGGCCAGTTGTTTAGGAAGGCCTTTA 2135
GGAAGACAAATAAATAACAAACAAACAGCCACAAATACTTTTTTTTCAAATTTTAGTTTTACCTGTAATTAATAAGAA 2214
CTGATACAAGACAAAAACAGTTCCTTCAGATTCTACGGAATGACAGTATATCTCTCTTTATCCTATGTGATTCTTGCTC 2293
TGAATGCATTATATTTTCCAAAGTATACCCATAAATTGTGACTAGTAAATACTTACACAGAGCAGAATTTTCACAGAT 2372
GGCAAAAAAATTTAAAGATGTCCAATATATGTGGGAAAAGAGCTAACAGAGAGATCATTATTTCTTAAAGATTGGCCAT 2451
AACCTGTATTTTGATAGAATTAGATTGGTAAATACATGTATTCATACATACTCTGTGGTAATAGAGACTTGAGCTGGAT 2530
CTGTACTGCACTGGAGTAAGCAAGAAAATTGGGAAAACCTTTTCGTTTGTTTCAGGTTTTGGCAACACATAGATCATATG 2609
TCTGAGGCACAAGTTGGCTGTTTCATCTTTGAAACCAGGGGATGCACAGTCTAAATGAATATCTGCATGGGATTTGCTAT 2688
CATAATNTTTCCTATGCNGNTGAATTCNGTGTGAGGTCTGTGTCCGTCTATCCTCAAATTATTTATTTTATAGTGCT 2767
GAGATCCTCAAATAATCTCAATTTTCGGAGGTTTCACAAAATGGACTCCTGAAGTAGACAGAGTAGTGAGGTTTCATTGC 2846
CCTCTATAAGCTTCTGACTAGCCAATGGCATCATCCAATTTCTTCCCAAACCTCTGCAGCATCTGCTTTATTGCCAAA 2925
GGGCTAGTTTCGGTTTTCTGCCAGCCATTGCGGTTAAAAAATATAAGTAGGATAACTTGTAACCTGCATATTGCTAA 3004
TCTATAGACACCACAGTTTCTAAATTCTTTGAAACCACTTTACTACTTTTTTTAACTTAACCTCAGTTCTAAATACTTT 3083
GTCTGGAGCACAAAACAATAAAAGGTTATCTTATAGTTGTGACTTTAACTTTTGTAGACCACAATTCAGTTTATAGTT 3162
TTCTTTTACTTAAATCCCATCTGCAGTCTCAAATTTAAGTTCTCCAGTAGAGATTGAGTTTGAGCCTGTATATCTATT 3241
AAAAATTTCAACTTCCCACATATATTTACTAAGATGATTAAGACTTACATTTTCTGCACAGGTCTGCAAAAACAAAAAT 3320
TATAAACTAGTCCATCCAAGAACCAAGTTTGTATAAACAGGTTGCTATAAGCTTGGTGAAATGAAAATGGAACATTTT 3399
AATCAAACATTTCTATATAACAATTATTATTTACAATTTGGTTTCTGCAATATTTTCTTATGTCCACCCTTTTAA 3479
AAATTATTATTTGAAGTAATTTATTTACAGGAAATGTTAATGAGATGTATTTTCTTATAGAGATATTTCTTACAGAAAG 3557
CTTTGTAGCAGAATATATTTGCAGCTATTGACTTTGTAATTTAGGAAAAATGTATAATAAGATAAAATCTATTAAATTT 3636
TTCTCCTCTAAAACTGAAAAAAAAAAAAAAAAAAAAAAAAAAGGGCGGCCGC 3687

7/15

Figure 5A

FGTCGACCCACGCGTCCGCTGTGGCAGCCCAGCTACCGGTCGTGACCAGATCCAGCTTGACGCTCAGCTTTGTTTCATTC 79
 GAATTGGGCGGCGGCCAGCGCGGAACAAAC M Q R L G G I L L C T L 12
 ATG CAG CGG CTC GGG GGT ATT TTG CTG TGT ACA CTG 145
 L A A A V P T A P A P S P T V T W T P A 32
 CTG GCG GCG GCG GTC CCC ACT GCT CCT GCT CCT TCC CCG ACG GTC ACT TGG ACT CCG GCG 205
 E P G P A L N Y P Q E E A T L N E M F R 52
 GAG CCG GGC CCA GCT CTC AAC TAC CCT CAG GAG GAA GCT ACG CTC AAT GAG ATG TTT CGA 265
 E V E E L M E D T Q H K L R S A V E E M 72
 GAG GTG GAG GAG CTG ATG GAA GAC ACT CAG CAC AAA CTG CGC AGT GCC GTG GAG GAG ATG 325
 E A E E A A A K T S S E V N L A S L P P 92
 GAG GCG GAA GAA GCA GCT GCT AAA ACG TCC TCT GAG GTG AAC CTG GCA AGC TTA CCT CCC 385
 N Y H N E T S T E T R V G N N T V H V H 112
 AAC TAT CAC AAT GAG ACC AGC ACG GAG ACC AGG GTG GGA AAT AAC ACA GTC CAT GTG CAC 445
 Q E V H K I T N N Q S G Q V V F S E T V 132
 CAG GAA GTT CAC AAG ATA ACC AAC AAC CAG AGT GGA CAG GTG GTC TTT TCT GAG ACA GTC 505
 I T S V G D E E G K R S H E C I I D E D 152
 ATT ACA TCT GTA GGG GAT GAA GAA GGC AAG AGG AGC CAT GAA TGT ATC ATT GAT GAA GAC 565
 C G P T R Y C Q F S S F K Y T C Q P C R 172
 TGT GGG CCC ACC AGG TAC TGC CAG TTC TCC AGC TTC AAG TAC ACC TGC CAG CCA TGC CGG 625
 D Q Q M L C T R D S E C C G D Q L C A W 192
 GAC CAG CAG ATG CTA TGC ACC CGA GAC AGT GAG TGC TGT GGA GAC CAG CTG TGT GCC TGG 685
 G H C T Q K A T K G G N G T I C D N Q R 212
 GGT CAC TGC ACC CAA AAG GCC ACC AAA GGT GGC AAT GGG ACC ATC TGT GAC AAC CAG AGG 745
 D C Q P G L C C A F Q R G L L F P V C T 232
 GAT TGC CAG CCT GGC CTG TGT TGT GCC TTC CAA AGA GGC CTG CTG TTC CCC GTG TGC ACA 805
 P L P V E G E L C H D P T S Q L L D L I 252
 CCC CTG CCC GTG GAG GGA GAG CTC TGC CAT GAC CCC ACC AGC CAG CTG CTG GAT CTC ATC 865
 T W E L E P E G A L D R C P C A S G L L 272
 ACC TGG GAA CTG GAG CCT GAA GGA GCT TTG GAC CGA TGC CCC TGC GCC AGT GGC CTC CTA 925
 C Q P H S H S L V Y M C K P A F V G S H 292
 TGC CAG CCA CAC AGC CAC AGT CTG GTG TAC ATG TGC AAG CCA GCC TTC GTG GGC AGC CAT 985
 D H S E E S Q L P R E A P D E Y E D V G 312
 GAC CAC AGT GAG GAG AGC CAG CTG CCC AGG GAG GCC CCG GAT GAG TAC GAA GAT GTT GGC 1045
 F I G E V R Q E L E D L E R S L A Q E M 332
 TTC ATA GGG GAA GTG CGC CAG GAG CTG GAA GAC CTG GAG CGG AGC CTA GCC CAG GAG ATG 1105

8/15

Figure 5B

A F E G P A P V E S L G G E E E I *	350
GCA TTT GAG GGG CCT GCC CCT GTG GAG TCA CTA GGC GGA GAG GAG GAG ATT TAG	1159
CCCCAGACCCAGCTGAGTCACTGGTAGATGTGCAATAGAAATGGCTAATTTATTTTCCCAGGAGTGTCCCCAAGTGTGG	1238
AATGGCCGCAGCTCCTTCCCAGTAGCTTTTCTCTGGCTTGACAAGGTACAGTGCAGTACATTTCTTCCAGCCGCCCTG	1317
CTTCTCTGACTTGGGAAAGACAGGCATGGCGGGTAAGGGCAGCGGTGAGTCGTCCCTCGCTGTTGCTAGAAACGCTGTC	1396
TTGTTCTTCATGGATGGAAGATTTGTTTGAAGGGAGAGGATGGGAAGGGGTGAAGTCTGCTCATGATGGATTTGGGGGA	1475
TACAGGGAGGAGGATGCCTGCCTTGACAGCTGGACTTGCCAAAATGTAACCTTTGCTTTTGTCTTGCGCCGCTCCCAT	1554
GGGCTGAGGCAGTGGCTACACAAGAGCTATGCTGCTCTGTGGCCTCCACATATTCATCCCTGTGTTTCAGCTCCTACC	1633
TCACTGTCAAGCAGCCCTTCATAGCCACGCCCCCTCTTGCTCACCACAGCCTAGGAGGGGACCAGAGGGGACTTCTCT	1712
CAGAGCCCCATGCTCTCTCTCTCAACCCCATACCAGCCTCTGTGCCAGCGACAGTCCTTCCAAATGGAGGGAGTGAAAT	1791
CCTTTGGTTTAATTATTTTCTCCTTCAAGGCACGCCTGCCACTAAGGTGAGGCTGACTTGTCATGTCCCTCTAACGTTTCG	1870
TAGCAGTGTGGTGGACACTGTCTTCCACCGACTGCTTCAATACCTCTGAAAGCCAGTGTCTCGGAGTGCAGTTTCGTGTAA	1949
ATTAATTTGCAGGAAGTATACTTGGCTAATTGTAGGGCTAGGATTGTGAATGAAATTTGCAAAGTCGCTTAGCAACAAT	2028
GGAAAGCCTTTCTCAGTCACACCGAGAAGTCACAACCAAGCCAGGTTGTGTAGAGTACAGCTGTGACATACAGACAGAA	2107
GAAGGCTGGGCTGGATGTGAGGCCTCAGATGACGGTTTCAGGTGCCAGGAATATTACCATTCTGTATCTATCCAGAGT	2186
TATTAATAATTGAAAGTTGCACACATTTGTATAAGCATGCCTTTCTCCTGAGTTTAAATTATATGTATACACAAACATG	2265
TGGCCCTCAAAGATCATGCACAAACCACTACTCTTTGCTAATTCTTGACTTTTCTCTTTGATTTTCAATAAATACAAA	2344
TCCCCTTCATGCAAAAAAAAAAAAAAAAAAGGGCGGCCGC	2381

Figure 6

9/15

```

1
hdck-1 -----
mdck-1 -----
xdck-1 -----
hdck-2 -----
hdck-3 -----
mdck-3 -----
cdck-3 -----
hdck-4 -----

60
hdck-1 -----
mdck-1 -----
xdck-1 -----
hdck-2 -----
hdck-3 -----
mdck-3 -----
cdck-3 -----
hdck-4 -----

61
hdck-1 -----
mdck-1 -----
xdck-1 -----
hdck-2 -----
hdck-3 -----
mdck-3 -----
cdck-3 -----
hdck-4 -----

120
hdck-1 -----
mdck-1 -----
xdck-1 -----
hdck-2 -----
hdck-3 -----
mdck-3 -----
cdck-3 -----
hdck-4 -----

180
hdck-1 -----
mdck-1 -----
xdck-1 -----
hdck-2 -----
hdck-3 -----
mdck-3 -----
cdck-3 -----
hdck-4 -----

240
hdck-1 -----
mdck-1 -----
xdck-1 -----
hdck-2 -----
hdck-3 -----
mdck-3 -----
cdck-3 -----
hdck-4 -----

241
hdck-1 -----
mdck-1 -----
xdck-1 -----
hdck-2 -----
hdck-3 -----
mdck-3 -----
cdck-3 -----
hdck-4 -----

300
hdck-1 -----
mdck-1 -----
xdck-1 -----
hdck-2 -----
hdck-3 -----
mdck-3 -----
cdck-3 -----
hdck-4 -----

301
hdck-1 -----
mdck-1 -----
xdck-1 -----
hdck-2 -----
hdck-3 -----
mdck-3 -----
cdck-3 -----
hdck-4 -----

360
hdck-1 -----
mdck-1 -----
xdck-1 -----
hdck-2 -----
hdck-3 -----
mdck-3 -----
cdck-3 -----
hdck-4 -----

361
hdck-1 -----
mdck-1 -----
xdck-1 -----
hdck-2 -----
hdck-3 -----
mdck-3 -----
cdck-3 -----
hdck-4 -----

424
hdck-1 -----
mdck-1 -----
xdck-1 -----
hdck-2 -----
hdck-3 -----
mdck-3 -----
cdck-3 -----
hdck-4 -----

```

10/15


Figure 7

CTCGAGGCCAAAATTCGGCACGAGGCCGGGCTGTGGTCTAGCATAAAGGCGGAGCCCAGAAGAAGGGGCGGGGT	M	1
ATG		77
G E A S P P A P A R R H L L V L L L L L		21
GGA GAA GCC TCC CCA CCT GCC CCC GCA AGG CGG CAT CTG CTG GTC CTG CTG CTG CTC CTC		137
S T L V I P S A A A P I H D A D A Q E S		41
TCT ACC CTG GTG ATC CCC TCC GCT GCA GCT CCT ATC CAT GAT GCT GAC GCC CAA GAG AGC		197
S L G L T G L Q S L L Q G F S R L F L K		61
TCC TTG GGT CTC ACA GGC CTC CAG AGC CTA CTC CAA GGC TTC AGC CGA CTT TTC CTG AAA		257
G N L L R G I D S L F S A P M D F R G L		81
GGT AAC CTG CTT CGG GGC ATA GAC AGC TTA TTC TCT GCC CCC ATG GAC TTC CGG GGC CTC		317
P G N Y H K E E N Q E H Q L G N N T L S		101
CCT GGG AAC TAC CAC AAA GAG GAG AAC CAG GAG CAC CAG CTG GGG AAC AAC ACC CTC TCC		377
S H L Q I D K M T D N K T G E V L I S E		121
AGC CAC CTC CAG ATC GAC AAG ATG ACC GAC AAC AAG ACA GGA GAG GTG CTG ATC TCC GAG		437
N V V A S I Q P A E G S F E G D L K V P		141
AAT GTG GTG GCA TCC ATT CAA CCA GCG GAG GGG AGC TTC GAG GGT GAT TTG AAG GTA CCC		497
R M E E K E A L V P I Q K A T D S F H T		161
AGG ATG GAG GAG AAG GAG GCC CTG GTA CCC ATC CAG AAG GCC ACG GAC AGC TTC CAC ACA		557
E L H P R V A F W I I K L P R R R S H Q		181
GAA CTC CAT CCC CGG GTG GCC TTC TGG ATC ATT AAG CTG CCA CGG CGG AGG TCC CAC CAG		617
D A L E G G H W L S E K R H R L Q A I R		201
GAT GCC CTG GAG GGC GGC CAC TGG CTC AGC GAG AAG CGA CAC CGC CTG CAG GCC ATC CGG		677
D G L R K G T H K D V L E E G T E S S S		221
GAT GGA CTC CGC AAG GGG ACC CAC AAG GAC GTC CTA GAA GAG GGG ACC GAG AGC TCC TCC		737
H S R L S P R K T H L L Y I L R P S R Q		241
CAC TCC AGG CTG TCC CCC CGA AAG ACC CAC TTA CTG TAC ATC CTC AGG CCC TCT CGG CAG		797
L *		243
CTG TAG		803
GGGTGGGGACCGGGGAGCACCTGCCTGTAGCCCCCATCAGACCCTGCCCCAAGCACCATATGGAAATAAAGTTCTTTCT		882
TACATCTAAAAAAAAAAAAAAAAAAAAAAAAAATTGGCGGCCGC		928

Figure 8

														M	C	R	L	R	
GAATTCGGGCACGAGGCAGAAGGCGCGAATGAAGGCAAAGCCTCCCACCCACCTGCA														ATG	TGT	CGA	CTG	AGG	
V	L	L	L	L	L	P	L	A	F	V	S	S	S	A	L	P	I	H	D
GTC	TTG	CTG	CTG	CTG	CTC	CCC	TTG	GCC	TTC	GTG	TCC	TCC	TCT	GCT	CTC	CCC	ATC	CAT	GAT
V	D	S	Q	Q	N	T	S	G	F	L	G	L	Q	R	L	L	Q	S	F
GTC	GAC	TCT	CAG	CAG	AAC	ACC	TCC	GGG	TTC	CTG	GGC	CTT	CAG	AGG	CTT	CTC	CAA	AGC	TTT
S	R	L	F	L	K	N	D	L	L	R	D	L	D	N	F	F	S	S	P
AGT	CGA	CTG	TTC	CTA	AAA	AAT	GAC	CTG	CTA	CGA	GAC	CTG	GAC	AAC	TTC	TTC	TCC	TCC	CCC
M	D	F	R	D	L	P	R	N	F	H	Q	E	E	N	Q	E	H	R	M
ATG	GAC	TTC	CGA	GAC	CTT	CCT	AGG	AAC	TTC	CAT	CAG	GAA	GAG	AAC	CAG	GAG	CAC	AGA	ATG
G	N	H	T	L	S	S	H	L	Q	I	D	K	V	T	D	N	Q	T	G
GGC	AAC	CAT	ACC	CTC	TCC	AGC	CAC	CTA	CAG	ATA	GAC	AAG	GTG	ACT	GAC	AAC	CAG	ACA	GGG
E	V	H	I	S	E	K	V	E	A	S	I	E	P	E	R	N	P	E	G
GAG	GTG	CAC	ATC	TCG	GAG	AAA	GTC	GAG	GCC	TCC	ATT	GAG	CCA	GAA	CGG	AAC	CCG	GAA	GGG
D	W	K	V	P	K	V	E	A	K	E	P	P	V	P	V	Q	K	V	T
GAC	TGG	AAG	GTT	CCC	AAA	GTA	GAA	GCA	AAA	GAG	CCC	CCG	GTG	CCT	GTG	CAG	AAG	GTC	ACC
D	S	L	H	P	E	P	R	Q	V	A	F	W	I	M	K	M	P	R	P
GAC	AGC	TTG	CAC	CCA	GAG	CCC	CGG	CAG	GTG	GCT	TTC	TGG	ATC	ATG	AAG	ATG	CCA	AGG	CGG
R	T	Q	P	D	V	Q	D	G	G	R	W	L	I	E	K	R	H	R	M
AGG	ACC	CAG	CCC	GAT	GTC	CAG	GAT	GGA	GGC	CGC	TGG	CTC	ATA	GAA	AAG	CGA	CAT	CGC	ATG
Q	A	I	R	D	G	L	R	G	G	A	R	E	D	S	L	E	D	G	V
CAG	GCC	ATC	CGG	GAT	GGG	CTC	CGT	GGA	GGC	GCC	CGT	GAG	GAC	AGC	CTG	GAG	GAT	GGG	GTC
H	I	P	Q	H	A	K	L	P	V	R	K	T	H	F	L	Y	I	L	R
CAT	ATC	CCC	CAA	CAC	GCC	AAG	CTG	CCT	GTC	AGA	AAG	ACA	CAC	TTT	CTC	TAC	ATC	CTC	AGG
P	S	Q	Q	L	*														
CCA	TCC	CAA	CAG	CTG	TAA	GTGGGGACCAGATGTCCCACACCCTACCCCAACACCATATGGAATAAAGGTTTTC													
TTACATCTAAAAAAAAAAAAAAAAAAAAAAAAA																			

12/15

Figure 9**A**

13/15

Figure 10

```

1
hsoggy  MGEASPPAPA RRHL.LVLLL LLSTLVIPSA AAPIHDAQAQ ESSLG.....LTGLQSL 60
msoggy  ----- MCRL.RVLLL LLPLAFVSSS ALPIHDVDSQ QNTSG.....FLGLQRL
hdck-3  ----- MQLGATLLC LLLAAVPTA PAPAPTATSA PVKPGPALSY PQEATLNEM
mdck-3  ----- MQLGGILLC TLLAAVPTA PAPSPTVTWT PAEPGPALNY PQEATLNEM
          *          *          *

61
hsoggy  LQGF SRLF.. LKGNLLRGID SL..... FSAPMDFRGL PGNYHKEENQ EHQLGNNTLS 120
msoggy  LQSF SRLF.. LKNDLLRDLD NF..... FSSPMDFRDL PRNFHQEENQ EHRMGNHTLS
hdck-3  FREVEELMED TQHKLRSAVE EMEAEAAAK ASSEVNLANL PPSYHNETNT DTKVGNNTIH
mdck-3  FREVEELMED TQHKLRSAVE EMEAEAAAK TSSEVNLASL PPNYHNETST ETRVGNNTVH
          *          *          *          *          *          *          *

121
hsoggy  SHLQIDKMTD NKTGEVLISE NVVASIQPAE GSFEGDLKVP RMEEKALVP IQKATDSFHT 180
msoggy  SHLQIDKVTD NQTGEVHISE KVEASIEP.E RNPEGDWKVP KVEAKEPPVP VQKVTDLSLHP
hdck-3  VHREIHKITN NQTGQMVFSE TVITSVGDEE GR.....RSHECII...DED CGP
mdck-3  VHQEVHKITN NQSGQVVFSE TVITSVGDEE GK.....RSHECII...DED CGP
          *          *          *          *          *          *          *

181
hsoggy  ELHPR.VAFW IIKLPRRRSH ....QDALEG GHWLSEKRHR LQAIRDGLRK 240
msoggy  E..PRQVAFW IMKMPPRRRTQ ....PDVQDG GRWLEKRHR MQAIRDGLRG
hdck-3  SMYQCFASFQ YTCQPCRGQR MLCTRDSECC GDQLCVWGHG TKMATRGSNG TIG
mdck-3  TRYQCFSSFK YTCQPCRDQQ MLCTRDSECC GDQLCAWGHG TQKATKGGNG TIG
          *          *          *          *          *          *          *

241
hsoggy  [REDACTED] 300
msoggy  [REDACTED]
hdck-3  [REDACTED]
mdck-3  [REDACTED]
          *          *          *          *          *          *          *

301
hdck-3  CPHSHSLVZVCKPTFVGSR DQDGEILLPR EVPDEYEVGS FMEEVRQELE DLERSLTEEM 360
mdck-3  CPHSHSLVZVCKPAFVGSH DHSEESQLPR EAPDEYEDVG FIGEVQELE DLERSLAQEM

361          379
hdck-3  ALGEPAAAAA ALLGGEEI-
mdck-3  AFEGPAPVES ..LGEEEEI

```

Figure 11





Figure 12



- 1 -

SEQUENCE LISTING

<110> MILLENNIUM PHARMACEUTICALS, INC.

<120> HUMAN DICKKOPF-RELATED PROTEIN AND NUCLEIC ACID
MOLECULES AND USES THEREFOR

<130> MNI-108CP2PC

<140> 09/263,022

<141> 1999-03-05

<160> 38

<170> PatentIn Ver. 2.0

<210> 1

<211> 2479

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (38)..(1087)

<220>

<223> 'n' at position 1146 may be any nucleotide

<400> 1

```

ggcacgaggg ggcgggcggt gcggggcgag agcggag atg cag cgg ctt ggg gcc 55
                                   Met Gln Arg Leu Gly Ala
                                   1           5

acc ctg ctg tgc ctg ctg ctg gcg gcg gcg gtc ccc acg gcc ccc gcg 103
Thr Leu Leu Cys Leu Leu Leu Ala Ala Ala Val Pro Thr Ala Pro Ala
           10           15           20

ccc gct ccg acg gcg acc tcg gct cca gtc aag ccc ggc ccg gct ctc 151
Pro Ala Pro Thr Ala Thr Ser Ala Pro Val Lys Pro Gly Pro Ala Leu
           25           30           35

agc tac ccg cag gag gag gcc acc ctc aat gag atg ttc cgc gag gtt 199
Ser Tyr Pro Gln Glu Glu Ala Thr Leu Asn Glu Met Phe Arg Glu Val
           40           45           50

gag gaa ctg atg gag gac acg cag cac aaa ttg cgc agc gcg gtg gaa 247
Glu Glu Leu Met Glu Asp Thr Gln His Lys Leu Arg Ser Ala Val Glu
           55           60           65           70

gag atg gag gca gaa gaa gct gct gct aaa gca tca tca gaa gtg aac 295
Glu Met Glu Ala Glu Glu Ala Ala Ala Lys Ala Ser Ser Glu Val Asn
           75           80           85

ctg gca aac tta cct ccc agc tat cac aat gag acc aac aca gac acg 343
Leu Ala Asn Leu Pro Pro Ser Tyr His Asn Glu Thr Asn Thr Asp Thr
           90           95           100

aac gtt gga aat aat acc atc cat gtg cac cga gaa att cac aag ata 391

```

- 2 -

Asn	Val	Gly	Asn	Asn	Thr	Ile	His	Val	His	Arg	Glu	Ile	His	Lys	Ile		
		105					110					115					
acc	aac	aac	cag	act	gga	caa	atg	gtc	ttt	tca	gag	aca	gtt	atc	aca	439	
Thr	Asn	Asn	Gln	Thr	Gly	Gln	Met	Val	Phe	Ser	Glu	Thr	Val	Ile	Thr		
		120				125					130						
tct	gtg	gga	gac	gaa	gaa	ggc	aga	agg	agc	cac	gag	tgc	atc	atc	gac	487	
Ser	Val	Gly	Asp	Glu	Glu	Gly	Arg	Arg	Ser	His	Glu	Cys	Ile	Ile	Asp		
135					140					145					150		
gag	gac	tgt	ggg	ccc	agc	atg	tac	tgc	cag	ttt	gcc	agc	ttc	cag	tac	535	
Glu	Asp	Cys	Gly	Pro	Ser	Met	Tyr	Cys	Gln	Phe	Ala	Ser	Phe	Gln	Tyr		
				155					160					165			
acc	tgc	cag	cca	tgc	cgg	ggc	cag	agg	atg	ctc	tgc	acc	cgg	gac	agt	583	
Thr	Cys	Gln	Pro	Cys	Arg	Gly	Gln	Arg	Met	Leu	Cys	Thr	Arg	Asp	Ser		
			170					175					180				
gag	tgc	tgt	gga	gac	cag	ctg	tgt	gtc	tgg	ggg	cac	tgc	acc	aaa	atg	631	
Glu	Cys	Cys	Gly	Asp	Gln	Leu	Cys	Val	Trp	Gly	His	Cys	Thr	Lys	Met		
		185					190					195					
gcc	acc	agg	ggc	agc	aat	ggg	acc	atc	tgt	gac	aac	cag	agg	gac	tgc	679	
Ala	Thr	Arg	Gly	Ser	Asn	Gly	Thr	Ile	Cys	Asp	Asn	Gln	Arg	Asp	Cys		
	200					205					210						
cag	ccg	ggg	ctg	tgc	tgt	gcc	ttc	cag	aga	ggc	ctg	ctg	ttc	cct	gtg	727	
Gln	Pro	Gly	Leu	Cys	Cys	Ala	Phe	Gln	Arg	Gly	Leu	Leu	Phe	Pro	Val		
215					220					225					230		
tgc	aca	ccc	ctg	ccc	gtg	gag	ggc	gag	ctt	tgc	cat	gac	ccc	gcc	agc	775	
Cys	Thr	Pro	Leu	Pro	Val	Glu	Gly	Glu	Leu	Cys	His	Asp	Pro	Ala	Ser		
				235					240					245			
cgg	ctt	ctg	gac	ctc	atc	acc	tgg	gag	cta	gag	cct	gat	gga	gcc	ttg	823	
Arg	Leu	Leu	Asp	Leu	Ile	Thr	Trp	Glu	Leu	Glu	Pro	Asp	Gly	Ala	Leu		
			250					255					260				
gac	cga	tgc	cct	tgt	gcc	agt	ggc	ctc	ctc	tgc	cag	ccc	cac	agc	cac	871	
Asp	Arg	Cys	Pro	Cys	Ala	Ser	Gly	Leu	Leu	Cys	Gln	Pro	His	Ser	His		
		265					270					275					
agc	ctg	gtg	tat	gtg	tgc	aag	ccg	acc	ttc	gtg	ggg	agc	cgt	gac	caa	919	
Ser	Leu	Val	Tyr	Val	Cys	Lys	Pro	Thr	Phe	Val	Gly	Ser	Arg	Asp	Gln		
	280					285					290						
gat	ggg	gag	atc	ctg	ctg	ccc	aga	gag	gtc	ccc	gat	gag	tat	gaa	gtt	967	
Asp	Gly	Glu	Ile	Leu	Leu	Pro	Arg	Glu	Val	Pro	Asp	Glu	Tyr	Glu	Val		
	295				300					305					310		
ggc	agc	ttc	atg	gag	gag	gtg	cgc	cag	gag	ctg	gag	gac	ctg	gag	agg	1015	
Gly	Ser	Phe	Met	Glu	Glu	Val	Arg	Gln	Glu	Leu	Glu	Asp	Leu	Glu	Arg		
				315					320					325			
agc	ctg	act	gaa	gag	atg	gcg	ctg	agg	gag	cct	gcg	gct	gcc	gcc	gct	1063	
Ser	Leu	Thr	Glu	Glu	Met	Ala	Leu	Arg	Glu	Pro	Ala	Ala	Ala	Ala	Ala		
			330					335					340				

- 3 -

gca ctg ctg gga agg gaa gag att tagatctgga ccaggctgtg ggtagatgtg 1117
Ala Leu Leu Gly Arg Glu Glu Ile
345 350

caatagaaat	agctaattta	tttccccang	tgtgtgcttt	aagcgtgggc	tgaccaggct	1177
tcttcctaca	tcttcttccc	agtaagtttc	ccctctggct	tgacagcatg	aggtgttgtg	1237
catttgttca	gctccccag	gctgttctcc	aggcttcaca	gtctgggtgt	tgggagagtc	1297
aggcagggtt	aaactgcagg	agcagtttgc	caccctgtc	cagattattg	gotgctttgc	1357
ctctaccagt	tggcagacag	ccgtttgttc	tacatggctt	tgataattgt	ttgaggggag	1417
gagatggaaa	caatgtggag	tctccctctg	attggttttg	gggaaatgtg	gagaagagtg	1477
ccctgctttg	caaacatcaa	cctggcaaaa	atgcaacaaa	tgaattttcc	acgcagttct	1537
ttccatgggc	ataggtaagc	tgtgccttca	gctgttgcat	atgaaatgtt	ctgttcaccc	1597
tgcatcacat	gtgtttattc	atccagcagt	gttgcctcag	tcctacctct	gtgccagggc	1657
agcattttca	tatccaagat	caattccctc	tctcagcaca	gcctggggag	ggggtcattg	1717
ttctcctcgt	ccatcaggga	tttcagaggc	tcagagactg	caagctgctt	gcccagtgca	1777
cacagctagt	gaagaccaga	gcagtttcat	ctggttgtga	ctctaagctc	agtgtctctct	1837
ccactacccc	acaccagcct	tggtgccacc	aaaagtgtc	ccaaaagga	aggagaatgg	1897
gatttttctt	ttgaggcatg	cacatctgga	attaagggtc	aactaattct	cacatccctc	1957
taaaagtaaa	ctactgttag	gaacagcagt	gttctcacag	tgtggggcag	ccgtccttct	2017
aatgaagaca	atgatattga	cactgtccct	ctttggcagt	tgcattagta	actttgaaag	2077
gtatatgact	gagcgtagca	tacagggtta	cctgcagaaa	cagtacttag	gtaattgtag	2137
ggcgaggatt	ataaatgaaa	tttgcaaaat	cacttagcag	caactgaaga	caattatcaa	2197
ccacgtggag	aaaatcaaac	cgagcagggc	tgtgtgaaac	atggttgtaa	tatgcgactg	2257
cgaacactga	actctacgcc	actccacaaa	tgatgttttc	aggtgtcatg	gactgttgcc	2317
accatgtatt	catccagagt	tcttaaagtt	taaagttgca	catgattgta	taagcatgct	2377
ttctttgagt	tttaaattat	gtataaacat	aagttgcatt	tagaaatcaa	gcataaatca	2437
cttcaactgc	taaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aa		2479

 $\langle 210 \rangle \quad 2$

<211> 350

<212> PRT

<213> Homo sapiens

<400> 2

Met Gln Arg Leu Gly Ala Thr Leu Leu Cys Leu Leu Leu Ala Ala Ala
1 5 10 15

Val	Pro	Thr	Ala	Pro	Ala	Pro	Ala	Pro	Thr	Ala	Thr	Ser	Ala	Pro	Val
			20					25					30		
Lys	Pro	Gly	Pro	Ala	Leu	Ser	Tyr	Pro	Gln	Glu	Glu	Ala	Thr	Leu	Asn
		35					40					45			
Glu	Met	Phe	Arg	Glu	Val	Glu	Glu	Leu	Met	Glu	Asp	Thr	Gln	His	Lys
	50					55					60				
Leu	Arg	Ser	Ala	Val	Glu	Glu	Met	Glu	Ala	Glu	Glu	Ala	Ala	Ala	Lys
65				70						75					80
Ala	Ser	Ser	Glu	Val	Asn	Leu	Ala	Asn	Leu	Pro	Pro	Ser	Tyr	His	Asn
				85					90					95	
Glu	Thr	Asn	Thr	Asp	Thr	Asn	Val	Gly	Asn	Asn	Thr	Ile	His	Val	His
			100					105					110		
Arg	Glu	Ile	His	Lys	Ile	Thr	Asn	Asn	Gln	Thr	Gly	Gln	Met	Val	Phe
		115					120					125			
Ser	Glu	Thr	Val	Ile	Thr	Ser	Val	Gly	Asp	Glu	Glu	Gly	Arg	Arg	Ser
	130					135					140				
His	Glu	Cys	Ile	Ile	Asp	Glu	Asp	Cys	Gly	Pro	Ser	Met	Tyr	Cys	Gln
145					150					155					160
Phe	Ala	Ser	Phe	Gln	Tyr	Thr	Cys	Gln	Pro	Cys	Arg	Gly	Gln	Arg	Met
				165					170					175	
Leu	Cys	Thr	Arg	Asp	Ser	Glu	Cys	Cys	Gly	Asp	Gln	Leu	Cys	Val	Trp
			180					185					190		
Gly	His	Cys	Thr	Lys	Met	Ala	Thr	Arg	Gly	Ser	Asn	Gly	Thr	Ile	Cys
		195					200					205			
Asp	Asn	Gln	Arg	Asp	Cys	Gln	Pro	Gly	Leu	Cys	Cys	Ala	Phe	Gln	Arg
	210					215					220				
Gly	Leu	Leu	Phe	Pro	Val	Cys	Thr	Pro	Leu	Pro	Val	Glu	Gly	Glu	Leu
225					230					235					240
Cys	His	Asp	Pro	Ala	Ser	Arg	Leu	Leu	Asp	Leu	Ile	Thr	Trp	Glu	Leu
				245					250					255	
Glu	Pro	Asp	Gly	Ala	Leu	Asp	Arg	Cys	Pro	Cys	Ala	Ser	Gly	Leu	Leu
			260					265					270		
Cys	Gln	Pro	His	Ser	His	Ser	Leu	Val	Tyr	Val	Cys	Lys	Pro	Thr	Phe
		275					280					285			
Val	Gly	Ser	Arg	Asp	Gln	Asp	Gly	Glu	Ile	Leu	Leu	Pro	Arg	Glu	Val
	290					295					300				
Pro	Asp	Glu	Tyr	Glu	Val	Gly	Ser	Phe	Met	Glu	Glu	Val	Arg	Gln	Glu
305					310					315					320
Leu	Glu	Asp	Leu	Glu	Arg	Ser	Leu	Thr	Glu	Glu	Met	Ala	Leu	Arg	Glu

- 5 -

325										330				335			
Pro	Ala	Ala	Ala	Ala	Ala	Ala	Ala	Leu	Leu	Gly	Arg	Glu	Glu	Ile			
			340						345					350			
<210> 3																	
<211> 1050																	
<212> DNA																	
<213> Homo sapiens																	
<220>																	
<221> CDS																	
<222> (1)..(1050)																	
<400> 3																	
atg	cag	cgg	ctt	ggg	gcc	acc	ctg	ctg	tgc	ctg	ctg	ctg	gcg	gcg	gcg	48	
Met	Gln	Arg	Leu	Gly	Ala	Thr	Leu	Leu	Cys	Leu	Leu	Leu	Ala	Ala	Ala		
1				5					10				15				
gtc	ccc	acg	gcc	ccc	gcg	ccc	gct	ccg	acg	gcg	acc	tcg	gct	cca	gtc	96	
Val	Pro	Thr	Ala	Pro	Ala	Pro	Ala	Pro	Thr	Ala	Thr	Ser	Ala	Pro	Val		
			20					25					30				
aag	ccc	ggc	ccg	gct	ctc	agc	tac	ccg	cag	gag	gag	gcc	acc	ctc	aat	144	
Lys	Pro	Gly	Pro	Ala	Leu	Ser	Tyr	Pro	Gln	Glu	Glu	Ala	Thr	Leu	Asn		
		35					40					45					
gag	atg	ttc	cgc	gag	gtt	gag	gaa	ctg	atg	gag	gac	acg	cag	cac	aaa	192	
Glu	Met	Phe	Arg	Glu	Val	Glu	Glu	Leu	Met	Glu	Asp	Thr	Gln	His	Lys		
	50					55					60						
ttg	cgc	agc	gcg	gtg	gaa	gag	atg	gag	gca	gaa	gaa	gct	gct	gct	aaa	240	
Leu	Arg	Ser	Ala	Val	Glu	Glu	Met	Glu	Ala	Glu	Glu	Ala	Ala	Ala	Lys		
65				70						75					80		
gca	tca	tca	gaa	gtg	aac	ctg	gca	aac	tta	cct	ccc	agc	tat	cac	aat	288	
Ala	Ser	Ser	Glu	Val	Asn	Leu	Ala	Asn	Leu	Pro	Pro	Ser	Tyr	His	Asn		
				85					90					95			
gag	acc	aac	aca	gac	acg	aac	gtt	gga	aat	aat	acc	atc	cat	gtg	cac	336	
Glu	Thr	Asn	Thr	Asp	Thr	Asn	Val	Gly	Asn	Asn	Thr	Ile	His	Val	His		
			100					105					110				
cga	gaa	att	cac	aag	ata	acc	aac	aac	cag	act	gga	caa	atg	gtc	ttt	384	
Arg	Glu	Ile	His	Lys	Ile	Thr	Asn	Asn	Gln	Thr	Gly	Gln	Met	Val	Phe		
		115					120					125					
tca	gag	aca	gtt	atc	aca	tct	gtg	gga	gac	gaa	gaa	ggc	aga	agg	agc	432	
Ser	Glu	Thr	Val	Ile	Thr	Ser	Val	Gly	Asp	Glu	Glu	Gly	Arg	Arg	Ser		
	130					135					140						
cac	gag	tgc	atc	atc	gac	gag	gac	tgt	ggg	ccc	agc	atg	tac	tgc	cag	480	
His	Glu	Cys	Ile	Ile	Asp	Glu	Asp	Cys	Gly	Pro	Ser	Met	Tyr	Cys	Gln		
145					150				155						160		
ttt	gcc	agc	ttc	cag	tac	acc	tgc	cag	cca	tgc	cgg	ggc	cag	agg	atg	528	
Phe	Ala	Ser	Phe	Gln	Tyr	Thr	Cys	Gln	Pro	Cys	Arg	Gly	Gln	Arg	Met		
				165				170						175			

- 6 -

```

ctc tgc acc cgg gac agt gag tgc tgt gga gac cag ctg tgt gtc tgg 576
Leu Cys Thr Arg Asp Ser Glu Cys Cys Gly Asp Gln Leu Cys Val Trp
180 185 190

ggt cac tgc acc aaa atg gcc acc agg ggc agc aat ggg acc atc tgt 624
Gly His Cys Thr Lys Met Ala Thr Arg Gly Ser Asn Gly Thr Ile Cys
195 200 205

gac aac cag agg gac tgc cag ccg ggg ctg tgc tgt gcc ttc cag aga 672
Asp Asn Gln Arg Asp Cys Gln Pro Gly Leu Cys Cys Ala Phe Gln Arg
210 215 220

ggc ctg ctg ttc cct gtg tgc aca ccc ctg ccc gtg gag ggc gag ctt 720
Gly Leu Leu Phe Pro Val Cys Thr Pro Leu Pro Val Glu Gly Glu Leu
225 230 235 240

tgc cat gac ccc gcc agc cgg ctt ctg gac ctc atc acc tgg gag cta 768
Cys His Asp Pro Ala Ser Arg Leu Leu Asp Leu Ile Thr Trp Glu Leu
245 250 255

gag cct gat gga gcc ttg gac cga tgc cct tgt gcc agt ggc ctc ctc 816
Glu Pro Asp Gly Ala Leu Asp Arg Cys Pro Cys Ala Ser Gly Leu Leu
260 265 270

tgc cag ccc cac agc cac agc ctg gtg tat gtg tgc aag ccg acc ttc 864
Cys Gln Pro His Ser His Ser Leu Val Tyr Val Cys Lys Pro Thr Phe
275 280 285

gtg ggg agc cgt gac caa gat ggg gag atc ctg ctg ccc aga gag gtc 912
Val Gly Ser Arg Asp Gln Asp Gly Glu Ile Leu Leu Pro Arg Glu Val
290 295 300

ccc gat gag tat gaa gtt ggc agc ttc atg gag gag gtg cgc cag gag 960
Pro Asp Glu Tyr Glu Val Gly Ser Phe Met Glu Glu Val Arg Gln Glu
305 310 315 320

ctg gag gac ctg gag agg agc ctg act gaa gag atg gcg ctg agg gag 1008
Leu Glu Asp Leu Glu Arg Ser Leu Thr Glu Glu Met Ala Leu Arg Glu
325 330 335

cct gcg gct gcc gcc gct gca ctg ctg gga agg gaa gag att 1050
Pro Ala Ala Ala Ala Ala Ala Leu Leu Gly Arg Glu Glu Ile
340 345 350

```

```

<210> 4
<211> 848
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> CDS
<222> (125)..(796)

```

```

<400> 4
gaattcggca cgagagacga cgtgctgagc tgccagctta gtggaagctc tgctctgggt 60
ggagagcagc ctgcgttttg tgacgcacag tgctgggacc ctccaggagc cccgggattg 120

```

- 7 -

```

aagg atg gtg gcg gcc gtc ctg ctg ggg ctg agc tgg ctc tgc tct ccc 169
  Met Val Ala Ala Val Leu Leu Gly Leu Ser Trp Leu Cys Ser Pro
    1             5             10             15

ctg gga gct ctg gtc ctg gac ttc aac aac atc agg agc tct gct gac 217
Leu Gly Ala Leu Val Leu Asp Phe Asn Asn Ile Arg Ser Ser Ala Asp
              20             25             30

ctg cat ggg gcc cgg aag ggc tca cag tgc ctg tct gac acg gac tgc 265
Leu His Gly Ala Arg Lys Gly Ser Gln Cys Leu Ser Asp Thr Asp Cys
              35             40             45

aat acc aga aag ttc tgc ctc cag ccc cgc gat gag aag ccg ttc tgt 313
Asn Thr Arg Lys Phe Cys Leu Gln Pro Arg Asp Glu Lys Pro Phe Cys
              50             55             60

gct aca tgt cgt ggg ttg cgg agg agg tgc cag cga gat gcc atg tgc 361
Ala Thr Cys Arg Gly Leu Arg Arg Arg Cys Gln Arg Asp Ala Met Cys
              65             70             75

tgc cct ggg aca ctc tgt gtg aac gat gtt tgt act acg atg gaa gat 409
Cys Pro Gly Thr Leu Cys Val Asn Asp Val Cys Thr Thr Met Glu Asp
              80             85             90             95

gca acc cca ata tta gaa agg cag ctt gat gag caa gat ggc aca cat 457
Ala Thr Pro Ile Leu Glu Arg Gln Leu Asp Glu Gln Asp Gly Thr His
              100            105            110

gca gaa gga aca act ggg cac cca gtc cag gaa aac caa ccc aaa agg 505
Ala Glu Gly Thr Thr Gly His Pro Val Gln Glu Asn Gln Pro Lys Arg
              115            120            125

aag cca agt att aag aaa tca caa ggc agg aag gga caa gag gga gaa 553
Lys Pro Ser Ile Lys Lys Ser Gln Gly Arg Lys Gly Gln Glu Gly Glu
              130            135            140

agt tgt ctg aga act ttt gac tgt ggc cct gga ctt tgc tgt gct cgt 601
Ser Cys Leu Arg Thr Phe Asp Cys Gly Pro Gly Leu Cys Cys Ala Arg
              145            150            155

cat ttt tgg acg aaa att tgt aag cca gtc ctt ttg gag gga cag gtc 649
His Phe Trp Thr Lys Ile Cys Lys Pro Val Leu Leu Glu Gly Gln Val
              160            165            170            175

tgc tcc aga aga ggg cat aaa gac act gct caa gct cca gaa atc ttc 697
Cys Ser Arg Arg Gly His Lys Asp Thr Ala Gln Ala Pro Glu Ile Phe
              180            185            190

cag cgt tgc gac tgt ggc cct gga cta ctg tgt cga agc caa ttg acc 745
Gln Arg Cys Asp Cys Gly Pro Gly Leu Leu Cys Arg Ser Gln Leu Thr
              195            200            205

agc aat cgg cag cat gct cga tta aga gta tgc caa aaa ata gaa aag 793
Ser Asn Arg Gln His Ala Arg Leu Arg Val Cys Gln Lys Ile Glu Lys
              210            215            220

cta taaatatttc aaaataaaga agaatccaca ttgcaaaaaa aaaaaaaaaa aa 848
Leu

```

- 8 -

<210> 5
 <211> 224
 <212> PRT
 <213> Homo sapiens

<400> 5

```

Met Val Ala Ala Val Leu Leu Gly Leu Ser Trp Leu Cys Ser Pro Leu
  1              5              10              15

Gly Ala Leu Val Leu Asp Phe Asn Asn Ile Arg Ser Ser Ala Asp Leu
          20              25              30

His Gly Ala Arg Lys Gly Ser Gln Cys Leu Ser Asp Thr Asp Cys Asn
      35              40              45

Thr Arg Lys Phe Cys Leu Gln Pro Arg Asp Glu Lys Pro Phe Cys Ala
    50              55              60

Thr Cys Arg Gly Leu Arg Arg Arg Cys Gln Arg Asp Ala Met Cys Cys
    65              70              75              80

Pro Gly Thr Leu Cys Val Asn Asp Val Cys Thr Thr Met Glu Asp Ala
          85              90              95

Thr Pro Ile Leu Glu Arg Gln Leu Asp Glu Gln Asp Gly Thr His Ala
      100              105              110

Glu Gly Thr Thr Gly His Pro Val Gln Glu Asn Gln Pro Lys Arg Lys
      115              120              125

Pro Ser Ile Lys Lys Ser Gln Gly Arg Lys Gly Gln Glu Gly Glu Ser
    130              135              140

Cys Leu Arg Thr Phe Asp Cys Gly Pro Gly Leu Cys Cys Ala Arg His
    145              150              155              160

Phe Trp Thr Lys Ile Cys Lys Pro Val Leu Leu Glu Gly Gln Val Cys
      165              170              175

Ser Arg Arg Gly His Lys Asp Thr Ala Gln Ala Pro Glu Ile Phe Gln
      180              185              190

Arg Cys Asp Cys Gly Pro Gly Leu Leu Cys Arg Ser Gln Leu Thr Ser
    195              200              205

Asn Arg Gln His Ala Arg Leu Arg Val Cys Gln Lys Ile Glu Lys Leu
    210              215              220

```

<210> 6
 <211> 672
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(672)

<400> 6
 atg gtg gcg gcc gtc ctg ctg ggg ctg agc tgg ctc tgc tct ccc ctg 48
 Met Val Ala Ala Val Leu Leu Gly Leu Ser Trp Leu Cys Ser Pro Leu
 1 5 10 15

gga gct ctg gtc ctg gac ttc aac aac atc agg agc tct gct gac ctg 96
 Gly Ala Leu Val Leu Asp Phe Asn Asn Ile Arg Ser Ser Ala Asp Leu
 20 25 30

cat ggg gcc cgg aag ggc tca cag tgc ctg tct gac acg gac tgc aat 144
 His Gly Ala Arg Lys Gly Ser Gln Cys Leu Ser Asp Thr Asp Cys Asn
 35 40 45

acc aga aag ttc tgc ctc cag ccc cgc gat gag aag ccg ttc tgt gct 192
 Thr Arg Lys Phe Cys Leu Gln Pro Arg Asp Glu Lys Pro Phe Cys Ala
 50 55 60

aca tgt cgt ggg ttg cgg agg agg tgc cag cga gat gcc atg tgc tgc 240
 Thr Cys Arg Gly Leu Arg Arg Arg Cys Gln Arg Asp Ala Met Cys Cys
 65 70 75 80

cct ggg aca ctc tgt gtg aac gat gtt tgt act acg atg gaa gat gca 288
 Pro Gly Thr Leu Cys Val Asn Asp Val Cys Thr Thr Met Glu Asp Ala
 85 90 95

acc cca ata tta gaa agg cag ctt gat gag caa gat ggc aca cat gca 336
 Thr Pro Ile Leu Glu Arg Gln Leu Asp Glu Gln Asp Gly Thr His Ala
 100 105 110

gaa gga aca act ggg cac cca gtc cag gaa aac caa ccc aaa agg aag 384
 Glu Gly Thr Thr Gly His Pro Val Gln Glu Asn Gln Pro Lys Arg Lys
 115 120 125

cca agt att aag aaa tca caa ggc agg aag gga caa gag gga gaa agt 432
 Pro Ser Ile Lys Lys Ser Gln Gly Arg Lys Gly Gln Glu Gly Glu Ser
 130 135 140

tgt ctg aga act ttt gac tgt ggc cct gga ctt tgc tgt gct cgt cat 480
 Cys Leu Arg Thr Phe Asp Cys Gly Pro Gly Leu Cys Cys Ala Arg His
 145 150 155 160

ttt tgg acg aaa att tgt aag cca gtc ctt ttg gag gga cag gtc tgc 528
 Phe Trp Thr Lys Ile Cys Lys Pro Val Leu Leu Glu Gly Gln Val Cys
 165 170 175

tcc aga aga ggg cat aaa gac act gct caa gct cca gaa atc ttc cag 576
 Ser Arg Arg Gly His Lys Asp Thr Ala Gln Ala Pro Glu Ile Phe Gln
 180 185 190

cgt tgc gac tgt ggc cct gga cta ctg tgt cga agc caa ttg acc agc 624
 Arg Cys Asp Cys Gly Pro Gly Leu Leu Cys Arg Ser Gln Leu Thr Ser
 195 200 205

aat cgg cag cat gct cga tta aga gta tgc caa aaa ata gaa aag cta 672
 Asn Arg Gln His Ala Arg Leu Arg Val Cys Gln Lys Ile Glu Lys Leu
 210 215 220

```
<210> 7
<211> 1536
<212> DNA
<213> Homo sapiens
```

```
<220>
<221> CDS
<222> (109) .. (906)
```

<400> 7																
gtcgacccac gcgctccgcgg acgcgtgggc ggcacggttt cgtgggggacc caggccttgca 60																
aagtgacggt cattttctct ttctttctcc ctcttgagtc cttctgag atg atg gct 117																
Met Met Ala 1																
ctg ggc gca gcg gga gct acc cgg gtc ttt gtc gcg atg gta gcg gcg 165																
Leu Gly Ala Ala Gly Ala Thr Arg Val Phe Val Ala Met Val Ala Ala																
5 10 15																
gct ctc ggc ggc cac cct ctg ctg gga gtg agc gcc acc ttg aac tcg 213																
Ala Leu Gly Gly His Pro Leu Leu Gly Val Ser Ala Thr Leu Asn Ser																
20 25 30 35																
gtt ctc aat tcc aac gct atc aag aac ctg ccc cca ccg ctg ggc ggc 261																
Val Leu Asn Ser Asn Ala Ile Lys Asn Leu Pro Pro Pro Leu Gly Gly																
40 45 50																
gct gcg ggg cac cca ggc tct gca gtc agc gcc gcg ccg gga atc ctg 309																
Ala Ala Gly His Pro Gly Ser Ala Val Ser Ala Ala Pro Gly Ile Leu																
55 60 65																
tac ccg ggc ggg aat aag tac cag acc att gac aac tac cag ccg tac 357																
Tyr Pro Gly Gly Asn Lys Tyr Gln Thr Ile Asp Asn Tyr Gln Pro Tyr																
70 75 80																
ccg tgc gca gag gac gag gag tgc ggc act gat gag tac tgc gct agt 405																
Pro Cys Ala Glu Asp Glu Glu Cys Gly Thr Asp Glu Tyr Cys Ala Ser																
85 90 95																
ccc acc cgc gga ggg gac gca ggc gtg caa atc tgt ctc gcc tgc agg 453																
Pro Thr Arg Gly Gly Asp Ala Gly Val Gln Ile Cys Leu Ala Cys Arg																
100 105 110 115																
aag cgc cga aaa cgc tgc atg cgt cac gct atg tgc tgc ccc ggg aat 501																
Lys Arg Arg Lys Arg Cys Met Arg His Ala Met Cys Cys Pro Gly Asn																
120 125 130																
tac tgc aaa aat gga ata tgt gtg tct tct gat caa aat cat ttc cga 549																
Tyr Cys Lys Asn Gly Ile Cys Val Ser Ser Asp Gln Asn His Phe Arg																
135 140 145																
gga gaa att gag gaa acc atc act gaa agc ttt ggt aat gat cat agc 597																
Gly Glu Ile Glu Glu Thr Ile Thr Glu Ser Phe Gly Asn Asp His Ser																
150 155 160																
acc ttg gat ggg tat tcc aga aga acc acc ttg tct tca aaa atg tat 645																
Thr Leu Asp Gly Tyr Ser Arg Arg Thr Thr Leu Ser Ser Lys Met Tyr																
165 170 175																

- 11 -

cac acc aaa gga caa gaa ggt tct gtt tgt ctc cgg tca tca gac tgt 693
 His Thr Lys Gly Gln Glu Gly Ser Val Cys Leu Arg Ser Ser Asp Cys
 180 185 190 195

gcc tca gga ttg tgt tgt gct aga cac ttc tgg tcc aag atc tgt aaa 741
 Ala Ser Gly Leu Cys Cys Ala Arg His Phe Trp Ser Lys Ile Cys Lys
 200 205 210

cct gtc ctg aaa gaa ggt caa gtg tgt acc aag cat agg aga aaa ggc 789
 Pro Val Leu Lys Glu Gly Gln Val Cys Thr Lys His Arg Arg Lys Gly
 215 220 225

tct cat gga cta gaa ata ttc cag cgt tgt tac tgt gga gaa ggt ctg 837
 Ser His Gly Leu Glu Ile Phe Gln Arg Cys Tyr Cys Gly Glu Gly Leu
 230 235 240

tct tgc cgg ata cag aaa gat cac cat caa gcc agt aat tct tct agg 885
 Ser Cys Arg Ile Gln Lys Asp His His Gln Ala Ser Asn Ser Ser Arg
 245 250 255

ctt cac act tgt cag aga cac taaaccagct atccaaatgc agtgaactcc 936
 Leu His Thr Cys Gln Arg His
 260 265

ttttatataa tagatgctat gaaaaccttt tatgaccttc atcaactcaa tcctaaggat 996
 atacaagttc tgtggtttca gttaagcatt ccaataacac cttccaaaaa cctggagtg 1056
 aagagctttg tttctttatg gaactccct gtgattgcag taaattactg tattgtaaat 1116
 tctcagtgtg gcacttacct gttaaagcaa tgaaactttt aattattttt cttaaagggtgc 1176
 tgcactgcct atttttcctc ttgttatgta aatttttgta cacattgatt gttatcttga 1236
 ctgacaaata ttctatattg aactgaagta aatcatttca gcttatagtt cttaaaagca 1296
 taacccttta ccccatTTaa ttctagagtc tagaacgcaa ggatctcttg gaatgacaaa 1356
 tgataggtac ctaaaatgta acatgaaaat actagcttat tttctgaaat gtactatctt 1416
 aatgcttaaa ttatatttcc ctttaggctg tgatagtttt tgaaataaaa tttaacattt 1476
 aatatcatga aatgttataa gtagacataa aaaaaaaaaa aaaaaaaaaa gggcgggccgc 1536

<210> 8
 <211> 266
 <212> PRT
 <213> Homo sapiens

<400> 8
 Met Met Ala Leu Gly Ala Ala Gly Ala Thr Arg Val Phe Val Ala Met
 1 5 10 15
 Val Ala Ala Ala Leu Gly Gly His Pro Leu Leu Gly Val Ser Ala Thr
 20 25 30
 Leu Asn Ser Val Leu Asn Ser Asn Ala Ile Lys Asn Leu Pro Pro Pro

```
<220>
<221> CDS
<222> (1) .. (798)
```

<400> 9
atg atg gct ctg ggc gca gcg gga gct acc cgg gtc ttt gtc gcg atg 48
Met Met Ala Leu Gly Ala Ala Gly Ala Thr Arg Val Phe Val Ala Met
1 5 10 15

- 13 -

gta gcg gcg gct ctc ggc ggc cac cct ctg ctg gga gtg agc gcc acc	96
Val Ala Ala Ala Leu Gly Gly His Pro Leu Leu Gly Val Ser Ala Thr	
20 25 30	
ttg aac tcg gtt ctc aat tcc aac gct atc aag aac ctg ccc cca ccg	144
Leu Asn Ser Val Leu Asn Ser Asn Ala Ile Lys Asn Leu Pro Pro Pro	
35 40 45	
ctg ggc ggc gct gcg ggg cac cca ggc tct gca gtc agc gcc gcg ccg	192
Leu Gly Gly Ala Ala Gly His Pro Gly Ser Ala Val Ser Ala Ala Pro	
50 55 60	
gga atc ctg tac ccg ggc ggg aat aag tac cag acc att gac aac tac	240
Gly Ile Leu Tyr Pro Gly Gly Asn Lys Tyr Gln Thr Ile Asp Asn Tyr	
65 70 75 80	
cag ccg tac ccg tgc gca gag gac gag gag tgc ggc act gat gag tac	288
Gln Pro Tyr Pro Cys Ala Glu Asp Glu Glu Cys Gly Thr Asp Glu Tyr	
85 90 95	
tgc gct agt ccc acc cgc gga ggg gac gca ggc gtg caa atc tgt ctc	336
Cys Ala Ser Pro Thr Arg Gly Gly Asp Ala Gly Val Gln Ile Cys Leu	
100 105 110	
gcc tgc agg aag cgc cga aaa cgc tgc atg cgt cac gct atg tgc tgc	384
Ala Cys Arg Lys Arg Arg Lys Arg Cys Met Arg His Ala Met Cys Cys	
115 120 125	
ccc ggg aat tac tgc aaa aat gga ata tgt gtg tct tct gat caa aat	432
Pro Gly Asn Tyr Cys Lys Asn Gly Ile Cys Val Ser Ser Asp Gln Asn	
130 135 140	
cat ttc cga gga gaa att gag gaa acc atc act gaa agc ttt ggt aat	480
His Phe Arg Gly Glu Ile Glu Glu Thr Ile Thr Glu Ser Phe Gly Asn	
145 150 155 160	
gat cat agc acc ttg gat ggg tat tcc aga aga acc acc ttg tct tca	528
Asp His Ser Thr Leu Asp Gly Tyr Ser Arg Arg Thr Thr Leu Ser Ser	
165 170 175	
aaa atg tat cac acc aaa gga caa gaa ggt tct gtt tgt ctc cgg tca	576
Lys Met Tyr His Thr Lys Gly Gln Glu Gly Ser Val Cys Leu Arg Ser	
180 185 190	
tca gac tgt gcc tca gga ttg tgt tgt gct aga cac ttc tgg tcc aag	624
Ser Asp Cys Ala Ser Gly Leu Cys Cys Ala Arg His Phe Trp Ser Lys	
195 200 205	
atc tgt aaa cct gtc ctg aaa gaa ggt caa gtg tgt acc aag cat agg	672
Ile Cys Lys Pro Val Leu Lys Glu Gly Gln Val Cys Thr Lys His Arg	
210 215 220	
aga aaa ggc tct cat gga cta gaa ata ttc cag cgt tgt tac tgt gga	720
Arg Lys Gly Ser His Gly Leu Glu Ile Phe Gln Arg Cys Tyr Cys Gly	
225 230 235 240	
gaa ggt ctg tct tgc cgg ata cag aaa gat cac cat caa gcc agt aat	768
Glu Gly Leu Ser Cys Arg Ile Gln Lys Asp His His Gln Ala Ser Asn	
245 250 255	

- 14 -

tct tct agg ctt cac act tgt cag aga cac 798
 Ser Ser Arg Leu His Thr Cys Gln Arg His
 260 265

<210> 10
 <211> 702
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(537)

<400> 10
 gaa ttc ggc acg agg gtt ggg agg tat tgc cac agt ccc cac caa gga 48
 Glu Phe Gly Thr Arg Val Gly Arg Tyr Cys His Ser Pro His Gln Gly
 1 5 10 15
 tca tgc gcc tgc atg gtg tgt cgg aga aaa aag aag cgc tgc cac cga 96
 Ser Ser Ala Cys Met Val Cys Arg Arg Lys Lys Lys Arg Cys His Arg
 20 25 30
 gat ggc atg tgc tgc ccc agt acc cgc tgc aat aat ggc atc tgt atc 144
 Asp Gly Met Cys Cys Pro Ser Thr Arg Cys Asn Asn Gly Ile Cys Ile
 35 40 45
 cca gtt act gaa agc atc tta acc cct cac atc ccg gct ctg gat ggt 192
 Pro Val Thr Glu Ser Ile Leu Thr Pro His Ile Pro Ala Leu Asp Gly
 50 55 60
 act cgg cac aga gat cga aac cac ggt cat tac tca aac cat gac ttg 240
 Thr Arg His Arg Asp Arg Asn His Gly His Tyr Ser Asn His Asp Leu
 65 70 75 80
 gga tgg cag aat cta gga aga cca cac act aag atg tca cat ata aaa 288
 Gly Trp Gln Asn Leu Gly Arg Pro His Thr Lys Met Ser His Ile Lys
 85 90 95
 ggg cat gaa gga gac ccc tgc cta cga tca tca gac tgc att gaa ggg 336
 Gly His Glu Gly Asp Pro Cys Leu Arg Ser Ser Asp Cys Ile Glu Gly
 100 105 110
 ttt tgc tgt gct cgt cat ttc tgg acc aaa atc tgc aaa cca gtg ctc 384
 Phe Cys Cys Ala Arg His Phe Trp Thr Lys Ile Cys Lys Pro Val Leu
 115 120 125
 cat cag ggg gaa gtc tgt acc aaa caa cgc aag aag ggt tct cat ggg 432
 His Gln Gly Glu Val Cys Thr Lys Gln Arg Lys Lys Gly Ser His Gly
 130 135 140
 ctg gaa att ttc cag cgt tgc gac tgt gcg aag ggc ctg tct tgc aaa 480
 Leu Glu Ile Phe Gln Arg Cys Asp Cys Ala Lys Gly Leu Ser Cys Lys
 145 150 155 160
 gta tgg aaa gat gcc acc tac tcc tcc aaa gcc aga ctc cat gtg tgt 528
 Val Trp Lys Asp Ala Thr Tyr Ser Ser Lys Ala Arg Leu His Val Cys
 165 170 175

- 15 -

cag aaa att tgatcaccat tgaggaacat catcaattgc agactgtgaa 577
 Gln Lys Ile
 gttgtgtatt taatgcatta tagcatgggtg gaaaataagg ttcagatgca gaagaatggc 637
 taaaataaga aacgtgataa gaatatagat gatcacaaaa aaaaaaaaaa aaaagatgcg 697
 gccgc 702

<210> 11
 <211> 179
 <212> PRT
 <213> Homo sapiens

<400> 11
 Glu Phe Gly Thr Arg Val Gly Arg Tyr Cys His Ser Pro His Gln Gly
 1 5 10 15
 Ser Ser Ala Cys Met Val Cys Arg Arg Lys Lys Lys Arg Cys His Arg
 20 25 30
 Asp Gly Met Cys Cys Pro Ser Thr Arg Cys Asn Asn Gly Ile Cys Ile
 35 40 45
 Pro Val Thr Glu Ser Ile Leu Thr Pro His Ile Pro Ala Leu Asp Gly
 50 55 60
 Thr Arg His Arg Asp Arg Asn His Gly His Tyr Ser Asn His Asp Leu
 65 70 75 80
 Gly Trp Gln Asn Leu Gly Arg Pro His Thr Lys Met Ser His Ile Lys
 85 90 95
 Gly His Glu Gly Asp Pro Cys Leu Arg Ser Ser Asp Cys Ile Glu Gly
 100 105 110
 Phe Cys Cys Ala Arg His Phe Trp Thr Lys Ile Cys Lys Pro Val Leu
 115 120 125
 His Gln Gly Glu Val Cys Thr Lys Gln Arg Lys Lys Gly Ser His Gly
 130 135 140
 Leu Glu Ile Phe Gln Arg Cys Asp Cys Ala Lys Gly Leu Ser Cys Lys
 145 150 155 160
 Val Trp Lys Asp Ala Thr Tyr Ser Ser Lys Ala Arg Leu His Val Cys
 165 170 175
 Gln Lys Ile

<210> 12
 <211> 537
 <212> DNA
 <213> Homo sapiens

- 16 -

<220>

<221> CDS

<222> (1)..(537)

<400> 12

gaa ttc ggc acg agg gtt ggg agg tat tgc cac agt ccc cac caa gga	48
Glu Phe Gly Thr Arg Val Gly Arg Tyr Cys His Ser Pro His Gln Gly	
1 5 10 15	
tca tcg gcc tgc atg gtg tgt cgg aga aaa aag aag cgc tgc cac cga	96
Ser Ser Ala Cys Met Val Cys Arg Arg Lys Lys Lys Arg Cys His Arg	
20 25 30	
gat ggc atg tgc tgc ccc agt acc cgc tgc aat aat ggc atc tgt atc	144
Asp Gly Met Cys Cys Pro Ser Thr Arg Cys Asn Asn Gly Ile Cys Ile	
35 40 45	
cca gtt act gaa agc atc tta acc cct cac atc ccg gct ctg gat ggt	192
Pro Val Thr Glu Ser Ile Leu Thr Pro His Ile Pro Ala Leu Asp Gly	
50 55 60	
act cgg cac aga gat cga aac cac ggt cat tac tca aac cat gac ttg	240
Thr Arg His Arg Asp Arg Asn His Gly His Tyr Ser Asn His Asp Leu	
65 70 75 80	
gga tgg cag aat cta gga aga cca cac act aag atg tca cat ata aaa	288
Gly Trp Gln Asn Leu Gly Arg Pro His Thr Lys Met Ser His Ile Lys	
85 90 95	
ggg cat gaa gga gac ccc tgc cta cga tca tca gac tgc att gaa ggg	336
Gly His Glu Gly Asp Pro Cys Leu Arg Ser Ser Asp Cys Ile Glu Gly	
100 105 110	
ttt tgc tgt gct cgt cat ttc tgg acc aaa atc tgc aaa cca gtg ctc	384
Phe Cys Cys Ala Arg His Phe Trp Thr Lys Ile Cys Lys Pro Val Leu	
115 120 125	
cat cag ggg gaa gtc tgt acc aaa caa cgc aag aag ggt tct cat ggg	432
His Gln Gly Glu Val Cys Thr Lys Gln Arg Lys Lys Gly Ser His Gly	
130 135 140	
ctg gaa att ttc cag cgt tgc gac tgt gcg aag ggc ctg tct tgc aaa	480
Leu Glu Ile Phe Gln Arg Cys Asp Cys Ala Lys Gly Leu Ser Cys Lys	
145 150 155 160	
gta tgg aaa gat gcc acc tac tcc tcc aaa gcc aga ctc cat gtg tgt	528
Val Trp Lys Asp Ala Thr Tyr Ser Ser Lys Ala Arg Leu His Val Cys	
165 170 175	
cag aaa att	537
Gln Lys Ile	

<210> 13

<211> 928

<212> DNA

<213> Homo sapiens

<220>

- 17 -

<221> CDS

<222> (75)..(800)

<400> 13

```

ctcgaggcca aaattcggca cgaggccggg ctgtggtcta gcataaaggc ggagcccaga 60

agaaggggacg ggggt atg gga gaa gcc tcc cca cct gcc ccc gca agg cgg 110
                Met Gly Glu Ala Ser Pro Pro Ala Pro Ala Arg Arg
                  1                      5                      10

cat ctg ctg gtc ctg ctg ctg ctc ctc tct acc ctg gtg atc ccc tcc 158
His Leu Leu Val Leu Leu Leu Leu Ser Thr Leu Val Ile Pro Ser
      15                      20                      25

gct gca gct cct atc cat gat gct gac gcc caa gag agc tcc ttg ggt 206
Ala Ala Ala Pro Ile His Asp Ala Asp Ala Gln Glu Ser Ser Leu Gly
      30                      35                      40

ctc aca ggc ctc cag agc cta ctc caa ggc ttc agc cga ctt ttc ctg 254
Leu Thr Gly Leu Gln Ser Leu Leu Gln Gly Phe Ser Arg Leu Phe Leu
      45                      50                      55                      60

aaa ggt aac ctg ctt cgg ggc ata gac agc tta ttc tct gcc ccc atg 302
Lys Gly Asn Leu Leu Arg Gly Ile Asp Ser Leu Phe Ser Ala Pro Met
                65                      70                      75

gac ttc cgg ggc ctc cct ggg aac tac cac aaa gag gag aac cag gag 350
Asp Phe Arg Gly Leu Pro Gly Asn Tyr His Lys Glu Glu Asn Gln Glu
                80                      85                      90

cac cag ctg ggg aac aac acc ctc tcc agc cac ctc cag atc gac aag 398
His Gln Leu Gly Asn Asn Thr Leu Ser Ser His Leu Gln Ile Asp Lys
      95                      100                      105

atg acc gac aac aag aca gga gag gtg ctg atc tcc gag aat gtg gtg 446
Met Thr Asp Asn Lys Thr Gly Glu Val Leu Ile Ser Glu Asn Val Val
      110                      115                      120

gca tcc att caa cca gcg gag ggg agc ttc gag ggt gat ttg aag gta 494
Ala Ser Ile Gln Pro Ala Glu Gly Ser Phe Glu Gly Asp Leu Lys Val
      125                      130                      135                      140

ccc agg atg gag gag aag gag gcc ctg gta ccc atc cag aag gcc acg 542
Pro Arg Met Glu Glu Lys Glu Ala Leu Val Pro Ile Gln Lys Ala Thr
                145                      150                      155

gac agc ttc cac aca gaa ctc cat ccc cgg gtg gcc ttc tgg atc att 590
Asp Ser Phe His Thr Glu Leu His Pro Arg Val Ala Phe Trp Ile Ile
                160                      165                      170

aag ctg cca cgg cgg agg tcc cac cag gat gcc ctg gag ggc ggc cac 638
Lys Leu Pro Arg Arg Arg Ser His Gln Asp Ala Leu Glu Gly Gly His
      175                      180                      185

tgg ctc agc gag aag cga cac cgc ctg cag gcc atc cgg gat gga ctc 686
Trp Leu Ser Glu Lys Arg His Arg Leu Gln Ala Ile Arg Asp Gly Leu
      190                      195                      200

cgc aag ggg acc cac aag gac gtc cta gaa gag ggg acc gag agc tcc 734

```

- 18 -

Arg Lys Gly Thr His Lys Asp Val Leu Glu Glu Gly Thr Glu Ser Ser
 205 210 215 220
 tcc cac tcc agg ctg tcc ccc cga aag acc cac tta ctg tac atc ctc 782
 Ser His Ser Arg Leu Ser Pro Arg Lys Thr His Leu Leu Tyr Ile Leu
 225 230 235
 agg ccc tct cgg cag ctg taggggtggg gaccggggag cacctgcctg 830
 Arg Pro Ser Arg Gln Leu
 240
 tagcccccat cagaccctgc cccaagcacc atatggaaat aaagttcttt cttacatcta 890
 aaaaaaaaaa aaaaaaaaaa aaaaaaattg gcggccgc 928

<210> 14
 <211> 242
 <212> PRT
 <213> Homo sapiens

<400> 14
 Met Gly Glu Ala Ser Pro Pro Ala Pro Ala Arg Arg His Leu Leu Val
 1 5 10 15
 Leu Leu Leu Leu Leu Ser Thr Leu Val Ile Pro Ser Ala Ala Ala Pro
 20 25 30
 Ile His Asp Ala Asp Ala Gln Glu Ser Ser Leu Gly Leu Thr Gly Leu
 35 40 45
 Gln Ser Leu Leu Gln Gly Phe Ser Arg Leu Phe Leu Lys Gly Asn Leu
 50 55 60
 Leu Arg Gly Ile Asp Ser Leu Phe Ser Ala Pro Met Asp Phe Arg Gly
 65 70 75 80
 Leu Pro Gly Asn Tyr His Lys Glu Glu Asn Gln Glu His Gln Leu Gly
 85 90 95
 Asn Asn Thr Leu Ser Ser His Leu Gln Ile Asp Lys Met Thr Asp Asn
 100 105 110
 Lys Thr Gly Glu Val Leu Ile Ser Glu Asn Val Val Ala Ser Ile Gln
 115 120 125
 Pro Ala Glu Gly Ser Phe Glu Gly Asp Leu Lys Val Pro Arg Met Glu
 130 135 140
 Glu Lys Glu Ala Leu Val Pro Ile Gln Lys Ala Thr Asp Ser Phe His
 145 150 155 160
 Thr Glu Leu His Pro Arg Val Ala Phe Trp Ile Ile Lys Leu Pro Arg
 165 170 175
 Arg Arg Ser His Gln Asp Ala Leu Glu Gly Gly His Trp Leu Ser Glu
 180 185 190
 Lys Arg His Arg Leu Gln Ala Ile Arg Asp Gly Leu Arg Lys Gly Thr

- 19 -

195	200	205	
His Lys Asp Val Leu Glu Glu Gly Thr Glu Ser Ser Ser His Ser Arg			
210	215	220	
Leu Ser Pro Arg Lys Thr His Leu Leu Tyr Ile Leu Arg Pro Ser Arg			
225	230	235	240
Gln Leu			
<210> 15			
<211> 726			
<212> DNA			
<213> Homo sapiens			
<220>			
<221> CDS			
<222> (1)..(726)			
<400> 15			
atg gga gaa gcc tcc cca cct gcc ccc gca agg cgg cat ctg ctg gtc	48		
Met Gly Glu Ala Ser Pro Pro Ala Pro Ala Arg Arg His Leu Leu Val			
1 5 10 15			
ctg ctg ctg ctc ctc tct acc ctg gtg atc ccc tcc gct gca gct cct	96		
Leu Leu Leu Leu Ser Thr Leu Val Ile Pro Ser Ala Ala Ala Pro			
20 25 30			
atc cat gat gct gac gcc caa gag agc tcc ttg ggt ctc aca ggc ctc	144		
Ile His Asp Ala Asp Ala Gln Glu Ser Ser Leu Gly Leu Thr Gly Leu			
35 40 45			
cag agc cta ctc caa ggc ttc agc cga ctt ttc ctg aaa ggt aac ctg	192		
Gln Ser Leu Leu Gln Gly Phe Ser Arg Leu Phe Leu Lys Gly Asn Leu			
50 55 60			
ctt cgg ggc ata gac agc tta ttc tct gcc ccc atg gac ttc cgg ggc	240		
Leu Arg Gly Ile Asp Ser Leu Phe Ser Ala Pro Met Asp Phe Arg Gly			
65 70 75 80			
ctc cct ggg aac tac cac aaa gag gag aac cag gag cac cag ctg ggg	288		
Leu Pro Gly Asn Tyr His Lys Glu Glu Asn Gln Glu His Gln Leu Gly			
85 90 95			
aac aac acc ctc tcc agc cac ctc cag atc gac aag atg acc gac aac	336		
Asn Asn Thr Leu Ser Ser His Leu Gln Ile Asp Lys Met Thr Asp Asn			
100 105 110			
aag aca gga gag gtg ctg atc tcc gag aat gtg gtg gca tcc att caa	384		
Lys Thr Gly Glu Val Leu Ile Ser Glu Asn Val Val Ala Ser Ile Gln			
115 120 125			
cca gcg gag ggg agc ttc gag ggt gat ttg aag gta ccc agg atg gag	432		
Pro Ala Glu Gly Ser Phe Glu Gly Asp Leu Lys Val Pro Arg Met Glu			
130 135 140			
gag aag gag gcc ctg gta ccc atc cag aag gcc acg gac agc ttc cac	480		

- 20 -

Glu	Lys	Glu	Ala	Leu	Val	Pro	Ile	Gln	Lys	Ala	Thr	Asp	Ser	Phe	His	
145					150					155					160	
aca	gaa	ctc	cat	ccc	cgg	gtg	gcc	ttc	tgg	atc	att	aag	ctg	cca	cgg	528
Thr	Glu	Leu	His	Pro	Arg	Val	Ala	Phe	Trp	Ile	Ile	Lys	Leu	Pro	Arg	
				165					170					175		
cgg	agg	tcc	cac	cag	gat	gcc	ctg	gag	ggc	ggc	cac	tgg	ctc	agc	gag	576
Arg	Arg	Ser	His	Gln	Asp	Ala	Leu	Glu	Gly	Gly	His	Trp	Leu	Ser	Glu	
			180					185					190			
aag	cga	cac	cgc	ctg	cag	gcc	atc	cgg	gat	gga	ctc	cgc	aag	ggg	acc	624
Lys	Arg	His	Arg	Leu	Gln	Ala	Ile	Arg	Asp	Gly	Leu	Arg	Lys	Gly	Thr	
		195					200					205				
cac	aag	gac	gtc	cta	gaa	gag	ggg	acc	gag	agc	tcc	tcc	cac	tcc	agg	672
His	Lys	Asp	Val	Leu	Glu	Glu	Gly	Thr	Glu	Ser	Ser	Ser	His	Ser	Arg	
	210					215					220					
ctg	tcc	ccc	cga	aag	acc	cac	tta	ctg	tac	atc	ctc	agg	ccc	tct	cgg	720
Leu	Ser	Pro	Arg	Lys	Thr	His	Leu	Leu	Tyr	Ile	Leu	Arg	Pro	Ser	Arg	
225					230					235					240	
cag	ctg															726
Gln	Leu															

<210> 16

<211> 2380

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (109)..(1155)

<400> 26

gtcgacccac gcgtccgctg tggcagccca gctaccggtc gtgaccagat ccagcttgca 60

gctcagcttt gttcattcga attgggcggc ggccagcgcg gaacaaac atg cag cgg 117
Met Gln Arg
1

ctc	ggg	ggt	att	ttg	ctg	tgt	aca	ctg	ctg	gcg	gcg	gcg	gtc	ccc	act	165
Leu	Gly	Gly	Ile	Leu	Leu	Cys	Thr	Leu	Leu	Ala	Ala	Ala	Val	Pro	Thr	
	5					10				15						

got	cct	gct	cct	tcc	cgg	acg	gtc	act	tgg	act	ccg	gcg	gag	ccg	ggc	213
Ala	Pro	Ala	Pro	Ser	Pro	Thr	Val	Thr	Trp	Thr	Pro	Ala	Glu	Pro	Gly	
20					25				30						35	

cca	gct	ctc	aac	tac	cct	cag	gag	gaa	gct	acg	ctc	aat	gag	atg	ttt	261
Pro	Ala	Leu	Asn	Tyr	Pro	Gln	Glu	Glu	Ala	Thr	Leu	Asn	Glu	Met	Phe	
			40						45					50		

cga	gag	gtg	gag	gag	ctg	atg	gaa	gac	act	cag	cac	aaa	ctg	cgc	agt	309
Arg	Glu	Val	Glu	Glu	Leu	Met	Glu	Asp	Thr	Gln	His	Lys	Leu	Arg	Ser	
			55					60						65		

- 21 -

gcc	gtg	gag	gag	atg	gag	gcg	gaa	gaa	gca	gct	gct	aaa	acg	tcc	tct	357
Ala	Val	Glu	Glu	Met	Glu	Ala	Glu	Glu	Ala	Ala	Ala	Lys	Thr	Ser	Ser	
		70					75					80				
gag	gtg	aac	ctg	gca	agc	tta	cct	ccc	aac	tat	cac	aat	gag	acc	agc	405
Glu	Val	Asn	Leu	Ala	Ser	Leu	Pro	Pro	Asn	Tyr	His	Asn	Glu	Thr	Ser	
	85					90					95					
acg	gag	acc	agg	gtg	gga	aat	aac	aca	gtc	cat	gtg	cac	cag	gaa	gtt	453
Thr	Glu	Thr	Arg	Val	Gly	Asn	Asn	Thr	Val	His	Val	His	Gln	Glu	Val	
100					105					110					115	
cac	aag	ata	acc	aac	aac	cag	agt	gga	cag	gtg	gtc	ttt	tct	gag	aca	501
His	Lys	Ile	Thr	Asn	Asn	Gln	Ser	Gly	Gln	Val	Val	Phe	Ser	Glu	Thr	
				120					125					130		
gtc	att	aca	tct	gta	ggg	gat	gaa	gaa	ggc	aag	agg	agc	cat	gaa	tgt	549
Val	Ile	Thr	Ser	Val	Gly	Asp	Glu	Glu	Gly	Lys	Arg	Ser	His	Glu	Cys	
			135				140						145			
atc	att	gat	gaa	gac	tgt	ggg	ccc	acc	agg	tac	tgc	cag	ttc	tcc	agc	597
Ile	Ile	Asp	Glu	Asp	Cys	Gly	Pro	Thr	Arg	Tyr	Cys	Gln	Phe	Ser	Ser	
		150					155					160				
ttc	aag	tac	acc	tgc	cag	cca	tgc	cgg	gac	cag	cag	atg	cta	tgc	acc	645
Phe	Lys	Tyr	Thr	Cys	Gln	Pro	Cys	Arg	Asp	Gln	Gln	Met	Leu	Cys	Thr	
	165					170					175					
cga	gac	agt	gag	tgc	tgt	gga	gac	cag	ctg	tgt	gcc	tgg	ggt	cac	tgc	693
Arg	Asp	Ser	Glu	Cys	Cys	Gly	Asp	Gln	Leu	Cys	Ala	Trp	Gly	His	Cys	
180					185					190					195	
acc	caa	aag	gcc	acc	aaa	ggt	ggc	aat	ggg	acc	atc	tgt	gac	aac	cag	741
Thr	Gln	Lys	Ala	Thr	Lys	Gly	Gly	Asn	Gly	Thr	Ile	Cys	Asp	Asn	Gln	
				200				205						210		
agg	gat	tgc	cag	cct	ggc	ctg	tgt	tgt	gcc	ttc	caa	aga	ggc	ctg	ctg	789
Arg	Asp	Cys	Gln	Pro	Gly	Leu	Cys	Cys	Ala	Phe	Gln	Arg	Gly	Leu	Leu	
			215					220					225			
ttc	ccc	gtg	tgc	aca	ccc	ctg	ccc	gtg	gag	gga	gag	ctc	tgc	cat	gac	837
Phe	Pro	Val	Cys	Thr	Pro	Leu	Pro	Val	Glu	Gly	Glu	Leu	Cys	His	Asp	
		230					235					240				
ccc	acc	agc	cag	ctg	ctg	gat	ctc	atc	acc	tgg	gaa	ctg	gag	cct	gaa	885
Pro	Thr	Ser	Gln	Leu	Leu	Asp	Leu	Ile	Thr	Trp	Glu	Leu	Glu	Pro	Glu	
		245				250					255					
gga	gct	ttg	gac	cga	tgc	ccc	tgc	gcc	agt	ggc	ctc	cta	tgc	cag	cca	933
Gly	Ala	Leu	Asp	Arg	Cys	Pro	Cys	Ala	Ser	Gly	Leu	Leu	Cys	Gln	Pro	
260					265					270					275	
cac	agc	cac	agt	ctg	gtg	tac	atg	tgc	aag	cca	gcc	ttc	gtg	ggc	agc	981
His	Ser	His	Ser	Leu	Val	Tyr	Met	Cys	Lys	Pro	Ala	Phe	Val	Gly	Ser	
				280				285						290		
cat	gac	cac	agt	gag	gag	agc	cag	ctg	ccc	agg	gag	gcc	ccg	gat	gag	1029
His	Asp	His	Ser	Glu	Glu	Ser	Gln	Leu	Pro	Arg	Glu	Ala	Pro	Asp	Glu	
			295					300					305			

- 22 -

```

tac gaa gat gtt ggc ttc ata ggg gaa gtg cgc cag gag ctg gaa gac 1077
Tyr Glu Asp Val Gly Phe Ile Gly Glu Val Arg Gln Glu Leu Glu Asp
      310                      315                      320

ctg gag cgg agc cta gcc cag gag atg gca ttt gag ggg cct gcc cct 1125
Leu Glu Arg Ser Leu Ala Gln Glu Met Ala Phe Glu Gly Pro Ala Pro
      325                      330                      335

gtg gag tca cta ggc gga gag gag gag att taggcccaga cccagctgag 1175
Val Glu Ser Leu Gly Gly Glu Glu Glu Ile
340                      345

tcaactggtag atgtgcaata gaaatggcta atttatatttc ccaggagtgt ccccaagtgt 1235

ggaatggccg cagctccttc ccagtagctt ttctctctggc ttgacaaggt acagtgcagt 1295

acattttcttc cagccgcctt gcttctctga cttgggaaag acaggcatgg cgggtaaggg 1355

cagcgggtgag tcgtccctcg ctgttgctag aaacgctgtc ttgttcttca tggatggaag 1415

atgtgtttga agggagagga tgggaagggg tgaagtctgc tcatgatgga tttgggggat 1475

acagggagga ggatgcctgc cttgcagacg tggacttggc aaaatgtaac ctttgctttt 1535

gtcttgccgc gctcccatgg gctgaggcag tggctacaca agagctatgc tgctctgtgg 1595

cctcccatat attcatcctt gtgtttcagc tcctacctca ctgtcagcac agcccttcat 1655

agccacgccc cctcttgctc accacagcct aggaggggac cagaggggac ttctctcaga 1715

gccccatgct ctctctctca accccatacc agcctctgtg ccagcgacag tccttccaaa 1775

tggaggggagt gaaatccttt ggtttaatta ttttctcctt caaggcacgc ctgccactaa 1835

ggtcaggctg acttgcatgt ccctctaacg ttcgtagcag tgtggtggac actgtcttcc 1895

accgactgct tcaatacctc tgaaagccag tgctcggagt gcagttcgtg taaattaatt 1955

tgcaggaagt atacttggtt aattgtaggg ctaggattgt gaatgaaatt tgcaaagtcg 2015

cttagcaaca atggaaagcc tttctcagtc acaccgagaa gtcacaacca agccaggttg 2075

tgtagagtac agctgtgaca tacagacaga agaaggctgg gctggatgtc aggcctcaga 2135

tgacggtttc aggtgccagg aactattacc attctgtatc tatccagagt tattaaaatt 2195

gaaagttgca cacatttgta taagcatgcc tttctcctga gtttttaaatt atatgtatac 2255

acaaacatgt ggccctcaaa gatcatgcac aaaccactac tctttgctaa ttcttggaact 2315

tttctctttg attttcaata aatacaaatc cccttcatgc aaaaaaaaaa aaaaagggcg 2375

gccgc 2380

```

<210> 17
 <211> 349
 <212> PRT

- 23 -

<213> Homo sapiens

<400> 17

```

Met  Gln  Arg  Leu  Gly  Gly  Ile  Leu  Leu  Cys  Thr  Leu  Leu  Ala  Ala  Ala
 1          5          10          15

Val  Pro  Thr  Ala  Pro  Ala  Pro  Ser  Pro  Thr  Val  Thr  Trp  Thr  Pro  Ala
          20          25          30

Glu  Pro  Gly  Pro  Ala  Leu  Asn  Tyr  Pro  Gln  Glu  Glu  Ala  Thr  Leu  Asn
          35          40          45

Glu  Met  Phe  Arg  Glu  Val  Glu  Glu  Leu  Met  Glu  Asp  Thr  Gln  His  Lys
 50          55          60

Leu  Arg  Ser  Ala  Val  Glu  Glu  Met  Glu  Ala  Glu  Glu  Ala  Ala  Ala  Lys
 65          70          75          80

Thr  Ser  Ser  Glu  Val  Asn  Leu  Ala  Ser  Leu  Pro  Pro  Asn  Tyr  His  Asn
          85          90          95

Glu  Thr  Ser  Thr  Glu  Thr  Arg  Val  Gly  Asn  Asn  Thr  Val  His  Val  His
          100          105          110

Gln  Glu  Val  His  Lys  Ile  Thr  Asn  Asn  Gln  Ser  Gly  Gln  Val  Val  Phe
          115          120          125

Ser  Glu  Thr  Val  Ile  Thr  Ser  Val  Gly  Asp  Glu  Glu  Gly  Lys  Arg  Ser
 130          135          140

His  Glu  Cys  Ile  Ile  Asp  Glu  Asp  Cys  Gly  Pro  Thr  Arg  Tyr  Cys  Gln
 145          150          155          160

Phe  Ser  Ser  Phe  Lys  Tyr  Thr  Cys  Gln  Pro  Cys  Arg  Asp  Gln  Gln  Met
          165          170          175

Leu  Cys  Thr  Arg  Asp  Ser  Glu  Cys  Cys  Gly  Asp  Gln  Leu  Cys  Ala  Trp
          180          185          190

Gly  His  Cys  Thr  Gln  Lys  Ala  Thr  Lys  Gly  Gly  Asn  Gly  Thr  Ile  Cys
          195          200          205

Asp  Asn  Gln  Arg  Asp  Cys  Gln  Pro  Gly  Leu  Cys  Cys  Ala  Phe  Gln  Arg
 210          215          220

Gly  Leu  Leu  Phe  Pro  Val  Cys  Thr  Pro  Leu  Pro  Val  Glu  Gly  Glu  Leu
 225          230          235          240

Cys  His  Asp  Pro  Thr  Ser  Gln  Leu  Leu  Asp  Leu  Ile  Thr  Trp  Glu  Leu
          245          250          255

Glu  Pro  Glu  Gly  Ala  Leu  Asp  Arg  Cys  Pro  Cys  Ala  Ser  Gly  Leu  Leu
          260          265          270

Cys  Gln  Pro  His  Ser  His  Ser  Leu  Val  Tyr  Met  Cys  Lys  Pro  Ala  Phe
          275          280          285

Val  Gly  Ser  His  Asp  His  Ser  Glu  Glu  Ser  Gln  Leu  Pro  Arg  Glu  Ala
 290          295          300

```

- 24 -

Pro Asp Glu Tyr Glu Asp Val Gly Phe Ile Gly Glu Val Arg Gln Glu
305 310 315 320

Leu Glu Asp Leu Glu Arg Ser Leu Ala Gln Glu Met Ala Phe Glu Gly
325 330 335

Pro Ala Pro Val Glu Ser Leu Gly Gly Glu Glu Glu Ile
340 345

<210> 18

<211> 1047

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(1047)

<400> 18

atg cag cgg ctc ggg ggt att ttg ctg tgt aca ctg ctg gcg gcg gcg 48
Met Gln Arg Leu Gly Gly Ile Leu Leu Cys Thr Leu Leu Ala Ala Ala
1 5 10 15

gtc ccc act gct cct gct cct tcc ccg acg gtc act tgg act ccg gcg 96
Val Pro Thr Ala Pro Ala Pro Ser Pro Thr Val Thr Trp Thr Pro Ala
20 25 30

gag ccg ggc cca gct ctc aac tac cct cag gag gaa gct acg ctc aat 144
Glu Pro Gly Pro Ala Leu Asn Tyr Pro Gln Glu Glu Ala Thr Leu Asn
35 40 45

gag atg ttt cga gag gtg gag gag ctg atg gaa gac act cag cac aaa 192
Glu Met Phe Arg Glu Val Glu Glu Leu Met Glu Asp Thr Gln His Lys
50 55 60

ctg cgc agt gcc gtg gag gag atg gag gcg gaa gaa gca gct gct aaa 240
Leu Arg Ser Ala Val Glu Glu Met Glu Ala Glu Glu Ala Ala Ala Lys
65 70 75 80

acg tcc tct gag gtg aac ctg gca agc tta cct ccc aac tat cac aat 288
Thr Ser Ser Glu Val Asn Leu Ala Ser Leu Pro Pro Asn Tyr His Asn
85 90 95

gag acc agc acg gag acc agg gtg gga aat aac aca gtc cat gtg cac 336
Glu Thr Ser Thr Glu Thr Arg Val Gly Asn Asn Thr Val His Val His
100 105 110

cag gaa gtt cac aag ata acc aac aac cag agt gga cag gtg gtc ttt 384
Gln Glu Val His Lys Ile Thr Asn Asn Gln Ser Gly Gln Val Val Phe
115 120 125

tct gag aca gtc att aca tct gta ggg gat gaa gaa ggc aag agg agc 432
Ser Glu Thr Val Ile Thr Ser Val Gly Asp Glu Glu Gly Lys Arg Ser
130 135 140

cat gaa tgt atc att gat gaa gac tgt ggg ccc acc agg tac tgc cag 480
His Glu Cys Ile Ile Asp Glu Asp Cys Gly Pro Thr Arg Tyr Cys Gln

- 25 -

145	150	155	160	
ttc tcc agc ttc aag tac acc tgc cag cca tgc cgg gac cag cag atg				528
Phe Ser Ser Phe Lys Tyr Thr Cys Gln Pro Cys Arg Asp Gln Gln Met				
	165	170	175	
cta tgc acc cga gac agt gag tgc tgt gga gac cag ctg tgt gcc tgg				576
Leu Cys Thr Arg Asp Ser Glu Cys Cys Gly Asp Gln Leu Cys Ala Trp				
	180	185	190	
ggt cac tgc acc caa aag gcc acc aaa ggt ggc aat ggg acc atc tgt				624
Gly His Cys Thr Gln Lys Ala Thr Lys Gly Gly Asn Gly Thr Ile Cys				
	195	200	205	
gac aac cag agg gat tgc cag cct ggc ctg tgt tgt gcc ttc caa aga				672
Asp Asn Gln Arg Asp Cys Gln Pro Gly Leu Cys Cys Ala Phe Gln Arg				
	210	215	220	
ggc ctg ctg ttc ccc gtg tgc aca ccc ctg ccc gtg gag gga gag ctc				720
Gly Leu Leu Phe Pro Val Cys Thr Pro Leu Pro Val Glu Gly Glu Leu				
	225	230	235	240
tgc cat gac ccc acc agc cag ctg ctg gat ctc atc acc tgg gaa ctg				768
Cys His Asp Pro Thr Ser Gln Leu Leu Asp Leu Ile Thr Trp Glu Leu				
	245	250	255	
gag cct gaa gga gct ttg gac cga tgc ccc tgc gcc agt ggc ctc cta				816
Glu Pro Glu Gly Ala Leu Asp Arg Cys Pro Cys Ala Ser Gly Leu Leu				
	260	265	270	
tgc cag cca cac agc cac agt ctg gtg tac atg tgc aag cca gcc ttc				864
Cys Gln Pro His Ser His Ser Leu Val Tyr Met Cys Lys Pro Ala Phe				
	275	280	285	
gtg ggc agc cat gac cac agt gag gag agc cag ctg ccc agg gag gcc				912
Val Gly Ser His Asp His Ser Glu Glu Ser Gln Leu Pro Arg Glu Ala				
	290	295	300	
ccg gat gag tac gaa gat gtt ggc ttc ata ggg gaa gtg cgc cag gag				960
Pro Asp Glu Tyr Glu Asp Val Gly Phe Ile Gly Glu Val Arg Gln Glu				
	305	310	315	320
ctg gaa gac ctg gag cgg agc cta gcc cag gag atg gca ttt gag ggg				1008
Leu Glu Asp Leu Glu Arg Ser Leu Ala Gln Glu Met Ala Phe Glu Gly				
	325	330	335	
cct gcc cct gtg gag tca cta ggc gga gag gag gag att				1047
Pro Ala Pro Val Glu Ser Leu Gly Gly Glu Glu Glu Ile				
	340	345		

<210> 19

<211> 8

<212> PRT

<213> synthtic construct

<400> 19

Asp Tyr Lys Asp Asp Asp Asp Lys

1

5

```
<220>  
<221> CDS  
<222> (712)..(1500)
```

<400>	20
gtcgcacccac gcgctccggcg ggagcccgcg gcgagcgtag cgcaagtccg ctccctaggc	60
atcgctgcgc tggcagcgat tcgctgtctc ttgtgagtca ggggacaacg cttcggggga	120
actgtgagtg cgcgtgtggg ggacctcgat tctcttcaga tctcgaggat tcgggtccggg	180
gacgtctcct gatccccctac taaagcgcct gctaactttg aaaaggagca ctgtgtcctg	240
caaagtttga cacataaagg ataggaaaag agaggagaga aaagcaactg agttgaagga	300
gaaggagctg atgcgggcct cctgatcaat taagaggaga gttaaaccgc cgagatcccg	360
gcggggacca ggaggtgcgg ggcaagaagg aacggaagcg gtgcgatcca cagggctggg	420
ttttcttgca ccttgggtca cgcctccttg gcgagaaaag ccctcgcatt tgattgcttc	480
cagttattgc agaacttcct gtcttggtgg agaagcgggt ctcgcttggg ttccgctaata	540
ttctgtcctg aggcgtaga ctgagttcat agggtccttg gtccccgaac cagggaagggt	600
tgagggaaca caatctgcaa gccccgcga cccaagttag gggccccgtg ttggggctct	660
ccctcccttt gcattcccac ccctccgggc tttgcgtctt cctggggacc c cct cgc	717
	Pro Arg 1
cgg gag atg gcc gcg ttg atg cgg agc aag gat tcg tcc tgc tgc ctg	765
Arg Glu Met Ala Ala Leu Met Arg Ser Lys Asp Ser Ser Cys Cys Leu	5 10 15
ctc cta ctg gcc gcg gtg ctg atg gtg gag agc tca cag atc ggc agt	813
Leu Leu Leu Ala Ala Val Leu Met Val Glu Ser Ser Gln Ile Gly Ser	20 25 30
tcg cgg gcc aaa ctc aac tcc atc aag tcc tct ctg ggc ggg gag acg	861
Ser Arg Ala Lys Leu Asn Ser Ile Lys Ser Ser Leu Gly Gly Glu Thr	35 40 45 50
cct ggt cag gcc gcc aat cga tct gcg ggc atg tac caa gga ctg gca	909
Pro Gly Gln Ala Ala Asn Arg Ser Ala Gly Met Tyr Gln Gly Leu Ala	55 60 65
ttc ggc ggc agt aag aag ggc aaa aac ctg ggg cag gcc tac cct tgt	957
Phe Gly Gly Ser Lys Lys Gly Lys Asn Leu Gly Gln Ala Tyr Pro Cys	70 75 80
agc agt gat aag gagtgt gaa gtt ggg agg tat tgc cac agt ccc cac	1005

Ser	Ser	Asp	Lys	Glu	Cys	Glu	Val	Gly	Arg	Tyr	Cys	His	Ser	Pro	His	
		85					90					95				
caa	gga	tca	tcg	gcc	tgc	atg	gtg	tgt	cgg	aga	aaa	aag	aag	cgc	tgc	1053
Gln	Gly	Ser	Ser	Ala	Cys	Met	Val	Cys	Arg	Arg	Lys	Lys	Lys	Arg	Cys	
	100					105					110					
cac	cga	gat	ggc	atg	tgc	tgc	ccc	agt	acc	cgc	tgc	aat	aat	ggc	atc	1101
His	Arg	Asp	Gly	Met	Cys	Cys	Pro	Ser	Thr	Arg	Cys	Asn	Asn	Gly	Ile	
	115				120					125					130	
tgt	atc	cca	gtt	act	gaa	agc	atc	tta	acc	cct	cac	atc	ccg	gct	ctg	1149
Cys	Ile	Pro	Val	Thr	Glu	Ser	Ile	Leu	Thr	Pro	His	Ile	Pro	Ala	Leu	
				135					140					145		
gat	ggt	act	cgg	cac	aga	gat	cga	aac	cac	ggt	cat	tac	tca	aac	cat	1197
Asp	Gly	Thr	Arg	His	Arg	Asp	Arg	Asn	His	Gly	His	Tyr	Ser	Asn	His	
			150					155					160			
gac	ttg	gga	tgg	cag	aat	cta	gga	aga	cca	cac	act	aag	atg	tca	cat	1245
Asp	Leu	Gly	Trp	Gln	Asn	Leu	Gly	Arg	Pro	His	Thr	Lys	Met	Ser	His	
		165					170					175				
ata	aaa	ggg	cat	gaa	gga	gac	ccc	tgc	cta	cga	tca	tca	gac	tgc	att	1293
Ile	Lys	Gly	His	Glu	Gly	Asp	Pro	Cys	Leu	Arg	Ser	Ser	Asp	Cys	Ile	
	180					185					190					
gaa	ggg	ttt	tgc	tgt	gct	cgt	cat	ttc	tgg	acc	aaa	atc	tgc	aaa	cca	1341
Glu	Gly	Phe	Cys	Cys	Ala	Arg	His	Phe	Trp	Thr	Lys	Ile	Cys	Lys	Pro	
	195				200					205					210	
gtg	ctc	cat	cag	ggg	gaa	gtc	tgt	acc	aaa	caa	cgc	aag	aag	ggt	tct	1389
Val	Leu	His	Gln	Gly	Glu	Val	Cys	Thr	Lys	Gln	Arg	Lys	Lys	Gly	Ser	
				215				220						225		
cat	ggg	ctg	gaa	att	ttc	cag	cgt	tgc	gac	tgt	gcg	aag	ggc	ctg	tct	1437
His	Gly	Leu	Glu	Ile	Phe	Gln	Arg	Cys	Asp	Cys	Ala	Lys	Gly	Leu	Ser	
			230				235						240			
tgc	aaa	gta	tgg	aaa	gat	gcc	acc	tac	tcc	tcc	aaa	gcc	aga	ctc	cat	1485
Cys	Lys	Val	Trp	Lys	Asp	Ala	Thr	Tyr	Ser	Ser	Lys	Ala	Arg	Leu	His	
		245					250					255				
gtg	tgt	cag	aaa	att	tgatcaccat	tgaggaacat	catcaattgc	agactgtgaa								1540
Val	Cys	Gln	Lys	Ile												
	260															
gtt	gtg	tatt	taatgcatta	tagcatgggtg	gaaaataaagg	ttcagatgca	gaagaatggc									1600
taaa	ataaga	aacgtgataa	gaatatagat	gatcacaaaa	aggggagaaag	aaaacatgaa										1660
ctga	atagat	tagaatgggt	gacaaatgca	gtgcagccag	tgttttccatt	atgcaacttg										1720
tctat	gtaaa	taatgtacac	atttgtggaa	aatgctatta	ttaagagaaac	aagcacacag										1780
tgga	aattac	tgatgagtag	catgtgactt	tccaagagtt	taggttgtgc	tggaggagag										1840
gttt	ccttca	gattgct														

gttagagttt aacaaaatac tcctagaata acttggtata caatagggtc taaaaataaa 1960
 attgctaaac aagaaatgaa aacatggagc attgttaatt tacaacagaa aattaccttt 2020
 tgatttgtaa cactacttct gctgttcaat caagagtctt ggtagataag aaaaaaatca 2080
 gtcaatattt ccaaataatt gcaaaataat ggccagttgt ttaggaaggc ctttaggaag 2140
 acaataaat aacaaacaaa cagccacaaa tacttttttt tcaaaatttt agttttacct 2200
 gtaattaata agaactgata caagacaaaa acagttcctt cagattctac ggaatgacag 2260
 tatactcttc tttatcctat gtgattcctg ctctgaatgc attatatttt ccaaagtata 2320
 ccataaatt gtgactagta aaatacttac acagagcaga attttcacag atggcaaaaa 2380
 aatttaaaga tgtccaatat atgtgggaaa agagctaaca gagagatcat tatttcttaa 2440
 agattggcca taacctgtat tttgatagaa ttagattggg aaatacatgt attcatacat 2500
 actctgtggt aatagagact tgagctggat ctgtactgca ctggagtaag caagaaaatt 2560
 gggaaaactt tttcgtttgt tcaggttttg gcaacacata gatcatatgt ctgaggcaca 2620
 agttggctgt tcatctttga aaccagggga tgcacagtct aaatgaatat ctgcatggga 2680
 tttgtatcat aatatttact atgcagatga attcagtgtg aggtcctgtg tccgtactat 2740
 cctcaaatta tttattttat agtgctgaga tcctcaaata atctcaattt caggaggttt 2800
 caaaaatgg actcctgaag tagacagagt agtgagggtt cattgccctc tataagcttc 2860
 tgactagcca atggcatcat ccaattttct tcccaaacct ctgcagcatc tgctttattg 2920
 ccaaagggtt agtttcggtt ttctgcagcc attgcggtta aaaaatataa gtaggataac 2980
 ttgtaaaacc tgcataattgc taatctatag acaccacagt ttctaaattc tttgaaacca 3040
 ctttactact ttttttaaac ttaactcagt tctaaatact ttgtctggag cacaaaacaa 3100
 taaaagggtta tcttatagtt gtgactttta actttttagt accacaattc acttttttagt 3160
 tttcttttac ttaaattcca tctgcagtct caaatttaag ttctcccagt agagattgag 3220
 tttgagcctg tatacttatt aaaaatttca acttcccaca tatatttact aagatgatta 3280
 agacttacat tttctgcaca ggtctgcaaa acaaaaaatt ataaactagt ccatccaaga 3340
 accaaagtth gtataaacag gttgctataa gcttggtgaa atgaaaatgg aacatttcaa 3400
 tcaaacattt cctatataac aattattata tttacaattt ggtttctgca atatttttct 3460
 tatgtccacc cttttaaaaa ttattatttg aagtaattta ttacaggaa atgttaatga 3520
 gatgtatttt cttatagaga tatttcttac agaaagcttt gtagcagaat atatttgcag 3580
 ctattgactt tgtaatttag gaaaaatgta taataagata aaatctatta aatttttctc 3640
 ctctaaaaac tgaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaagggc ggccgc 3696

<210> 21
 <211> 263
 <212> PRT
 <213> Homo sapiens

<400> 21

```

Pro Arg Arg Glu Met Ala Ala Leu Met Arg Ser Lys Asp Ser Ser Cys
 1           5           10           15

Cys Leu Leu Leu Leu Ala Ala Val Leu Met Val Glu Ser Ser Gln Ile
      20           25           30

Gly Ser Ser Arg Ala Lys Leu Asn Ser Ile Lys Ser Ser Leu Gly Gly
      35           40           45

Glu Thr Pro Gly Gln Ala Ala Asn Arg Ser Ala Gly Met Tyr Gln Gly
 50           55           60

Leu Ala Phe Gly Gly Ser Lys Lys Gly Lys Asn Leu Gly Gln Ala Tyr
 65           70           75           80

Pro Cys Ser Ser Asp Lys Glu Cys Glu Val Gly Arg Tyr Cys His Ser
      85           90           95

Pro His Gln Gly Ser Ser Ala Cys Met Val Cys Arg Arg Lys Lys Lys
      100          105          110

Arg Cys His Arg Asp Gly Met Cys Cys Pro Ser Thr Arg Cys Asn Asn
      115          120          125

Gly Ile Cys Ile Pro Val Thr Glu Ser Ile Leu Thr Pro His Ile Pro
      130          135          140

Ala Leu Asp Gly Thr Arg His Arg Asp Arg Asn His Gly His Tyr Ser
      145          150          155          160

Asn His Asp Leu Gly Trp Gln Asn Leu Gly Arg Pro His Thr Lys Met
      165          170          175

Ser His Ile Lys Gly His Glu Gly Asp Pro Cys Leu Arg Ser Ser Asp
      180          185          190

Cys Ile Glu Gly Phe Cys Cys Ala Arg His Phe Trp Thr Lys Ile Cys
      195          200          205

Lys Pro Val Leu His Gln Gly Glu Val Cys Thr Lys Gln Arg Lys Lys
      210          215          220

Gly Ser His Gly Leu Glu Ile Phe Gln Arg Cys Asp Cys Ala Lys Gly
      225          230          235          240

Leu Ser Cys Lys Val Trp Lys Asp Ala Thr Tyr Ser Ser Lys Ala Arg
      245          250          255

Leu His Val Cys Gln Lys Ile
      260

```

- 30 -

<210> 22
 <211> 789
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(789)

<400> 22

```

cct cgc cgg gag atg gcc gcg ttg atg cgg agc aag gat tcg tcc tgc      48
Pro Arg Arg Glu Met Ala Ala Leu Met Arg Ser Lys Asp Ser Ser Cys
   1                               5                               10                               15

tgc ctg ctc cta ctg gcc gcg gtg ctg atg gtg gag agc tca cag atc      96
Cys Leu Leu Leu Leu Ala Ala Val Leu Met Val Glu Ser Ser Gln Ile
                20                               25                               30

ggc agt tcg cgg gcc aaa ctc aac tcc atc aag tcc tct ctg ggc ggc      144
Gly Ser Ser Arg Ala Lys Leu Asn Ser Ile Lys Ser Ser Leu Gly Gly
                35                               40                               45

gag acg cct ggt cag gcc gcc aat cga tct gcg ggc atg tac caa gga      192
Glu Thr Pro Gly Gln Ala Ala Asn Arg Ser Ala Gly Met Tyr Gln Gly
                50                               55                               60

ctg gca ttc ggc ggc agt aag aag ggc aaa aac ctg ggg cag gcc tac      240
Leu Ala Phe Gly Gly Ser Lys Lys Gly Lys Asn Leu Gly Gln Ala Tyr
   65                               70                               75                               80

cct tgt agc agt gat aag gag tgt gaa gtt ggg agg tat tgc cac agt      288
Pro Cys Ser Ser Asp Lys Glu Cys Glu Val Gly Arg Tyr Cys His Ser
                85                               90                               95

ccc cac caa gga tca tcg gcc tgc atg gtg tgt cgg aga aaa aag aag      336
Pro His Gln Gly Ser Ser Ala Cys Met Val Cys Arg Arg Lys Lys Lys
                100                               105                               110

cgc tgc cac cga gat ggc atg tgc tgc ccc agt acc cgc tgc aat aat      384
Arg Cys His Arg Asp Gly Met Cys Cys Pro Ser Thr Arg Cys Asn Asn
                115                               120                               125

ggc atc tgt atc cca gtt act gaa agc atc tta acc cct cac atc ccg      432
Gly Ile Cys Ile Pro Val Thr Glu Ser Ile Leu Thr Pro His Ile Pro
                130                               135                               140

gct ctg gat ggt act cgg cac aga gat cga aac cac ggt cat tac tca      480
Ala Leu Asp Gly Thr Arg His Arg Asp Arg Asn His Gly His Tyr Ser
   145                               150                               155                               160

aac cat gac ttg gga tgg cag aat cta gga aga cca cac act aag atg      528
Asn His Asp Leu Gly Trp Gln Asn Leu Gly Arg Pro His Thr Lys Met
                165                               170                               175

tca cat ata aaa ggc cat gaa gga gac ccc tgc cta cga tca tca gac      576
Ser His Ile Lys Gly His Glu Gly Asp Pro Cys Leu Arg Ser Ser Asp
                180                               185                               190

```

- 31 -

```

tgc att gaa ggg ttt tgc tgt gct cgt cat ttc tgg acc aaa atc tgc 624
Cys Ile Glu Gly Phe Cys Cys Ala Arg His Phe Trp Thr Lys Ile Cys
      195                200                205

aaa cca gtg ctc cat cag ggg gaa gtc tgt acc aaa caa cgc aag aag 672
Lys Pro Val Leu His Gln Gly Glu Val Cys Thr Lys Gln Arg Lys Lys
      210                215                220

ggt tct cat ggg ctg gaa att ttc cag cgt tgc gac tgt gcg aag ggc 720
Gly Ser His Gly Leu Glu Ile Phe Gln Arg Cys Asp Cys Ala Lys Gly
      225                230                235                240

ctg tct tgc aaa gta tgg aaa gat gcc acc tac tcc tcc aaa gcc aga 768
Leu Ser Cys Lys Val Trp Lys Asp Ala Thr Tyr Ser Ser Lys Ala Arg
      245                250                255

ctc cat gtg tgt cag aaa att
Leu His Val Cys Gln Lys Ile
      260

```

<210> 23

<211> 54

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: consensus sequence

<220>

<223> Xaa's at positions
2, 3, 5, 6, 8-12, 14-26, 28, 29, 31-36, 38-42, 45-48, and
50-53 may be any amino acid

<220>

<223> Xaa's at postions 22-26 may be absent

<400> 23

```

Cys Xaa Xaa Asp Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa Xaa
  1              5              10              15

```

```

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa Cys Xaa Xaa
      20              25              30

```

```

Xaa Xaa Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa Cys Cys Xaa Xaa Xaa Xaa
      35              40              45

```

```

Cys Xaa Xaa Xaa Xaa Cys
      50

```

<210> 24

<211> 123

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: consensus

- 32 -

sequence

<220>

<223> Xaa's at positions

2,5,8,9,11,14-23,25,27-30,32,33,35-53,60,62,63,65,
68,70,71,73-96,98,100,101,104, and 106-122 may be
any amino acid

<220>

<223> Xaa's at positions 22,23,51-58,89-96 and 116-122
may be absent

<400> 24

Cys	Xaa	Xaa	Xaa	Xaa	Asp	Cys	Xaa	Xaa	Gly	Xaa	Cys	Cys	Xaa	Xaa	Xaa	1	5	10	15
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Cys	Xaa	Pro	Xaa	Xaa	Xaa	Xaa	Gly	Xaa	20	25	30	
Xaa	Cys	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	35	40	45	
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Cys	Xaa	Cys	Xaa	Xaa	Pro	50	55	60	
Xaa	Xaa	Xaa	Xaa	Gly	Xaa	Cys	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	65	70	75	80
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	85	90	95	
Cys	Xaa	Cys	Xaa	Xaa	Gly	Leu	Xaa	Cys	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	100	105	110	
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Cys						115	120		

<210> 25

<211> 87

<212> PRT

<213> Homo sapiens

<400> 25

Ile	Asn	Leu	Glu	Asn	Gly	Glu	Leu	Cys	Met	Asn	Ser	Ala	Gln	Cys	Lys	1	5	10	15
Ser	Asn	Cys	Cys	Gln	His	Ser	Ser	Ala	Leu	Gly	Leu	Ala	Arg	Cys	Thr	20	25	30	
Ser	Met	Ala	Ser	Glu	Asn	Ser	Glu	Cys	Ser	Val	Lys	Thr	Leu	Tyr	Gly	35	40	45	
Ile	Tyr	Tyr	Lys	Cys	Pro	Cys	Glu	Arg	Gly	Leu	Thr	Cys	Glu	Gly	Asp	50	55	60	
Lys	Thr	Ile	Val	Gly	Ser	Ile	Thr	Asn	Thr	Asn	Phe	Gly	Ile	Cys	His	65	70	75	80

- 33 -

Asp Ala Gly Arg Ser Lys Gln
85

<210> 26
<211> 835
<212> DNA
<213> Mus musculus

<220>
<221> CDS
<222> (57)..(746)

<400> 25
gaattcggca cgaggcagaa ggcgcgaaatg aaggcaaagc ctcccaccca cctgca atg 59
Met
1

tgt cga ctg agg gtc ttg ctg ctg ctg ctc ccc ttg gcc ttc gtg tcc 107
Cys Arg Leu Arg Val Leu Leu Leu Leu Leu Pro Leu Ala Phe Val Ser
5 10 15

tcc tct gct ctc ccc atc cat gat gtc gac tct cag cag aac acc tcc 155
Ser Ser Ala Leu Pro Ile His Asp Val Asp Ser Gln Gln Asn Thr Ser
20 25 30

ggg ttc ctg ggc ctt cag agg ctt ctc caa agc ttt agt cga ctg ttc 203
Gly Phe Leu Gly Leu Gln Arg Leu Leu Gln Ser Phe Ser Arg Leu Phe
35 40 45

cta aaa aat gac ctg cta cga gac ctg gac aac ttc ttc tcc tcc ccc 251
Leu Lys Asn Asp Leu Leu Arg Asp Leu Asp Asn Phe Phe Ser Ser Pro
50 55 60 65

atg gac ttc cga gac ctt cct agg aac ttc cat cag gaa gag aac cag 299
Met Asp Phe Arg Asp Leu Pro Arg Asn Phe His Gln Glu Glu Asn Gln
70 75 80

gag cac aga atg ggc aac cat acc ctc tcc agc cac cta cag ata gac 347
Glu His Arg Met Gly Asn His Thr Leu Ser Ser His Leu Gln Ile Asp
85 90 95

aag gtg act gac aac cag aca ggg gag gtg cac atc tcg gag aaa gtc 395
Lys Val Thr Asp Asn Gln Thr Gly Glu Val His Ile Ser Glu Lys Val
100 105 110

gag gcc tcc att gag cca gaa cgg aac ccg gaa ggg gac tgg aag gtt 443
Glu Ala Ser Ile Glu Pro Glu Arg Asn Pro Glu Gly Asp Trp Lys Val
115 120 125

ccc aaa gta gaa gca aaa gag ccc ccg gtg cct gtg cag aag gtc acc 491
Pro Lys Val Glu Ala Lys Glu Pro Pro Val Pro Val Gln Lys Val Thr
130 135 140 145

gac agc ttg cac cca gag ccc cgg cag gtg gct ttc tgg atc atg aag 539
Asp Ser Leu His Pro Glu Pro Arg Gln Val Ala Phe Trp Ile Met Lys
150 155 160

atg cca agg cgg agg acc cag ccc gat gtc cag gat gga ggc cgc tgg 587

- 34 -

Met Pro Arg Arg Arg Thr Gln Pro Asp Val Gln Asp Gly Gly Arg Trp
165 170 175

ctc ata gaa aag cga cat cgc atg cag gcc atc cgg gat ggg ctc cgt 635
Leu Ile Glu Lys Arg His Arg Met Gln Ala Ile Arg Asp Gly Leu Arg
180 185 190

gga ggc gcc cgt gag gac agc ctg gag gat ggg gtc cat atc ccc caa 683
Gly Gly Ala Arg Glu Asp Ser Leu Glu Asp Gly Val His Ile Pro Gln
195 200 205

cac gcc aag ctg cct gtc aga aag aca cac ttt ctc tac atc ctc agg 731
His Ala Lys Leu Pro Val Arg Lys Thr His Phe Leu Tyr Ile Leu Arg
210 215 220 225

cca tcc caa cag ctg taagtgggga ccagatgtcc cacaccctac cccaacacca 786
Pro Ser Gln Gln Leu
230

tatggaaata aaggttttct tacatctaaa aaaaaaaaaa aaaaaaaaaa 835

<210> 27
<211> 230
<212> PRT
<213> Mus musculus

<400> 27
Met Cys Arg Leu Arg Val Leu Leu Leu Leu Leu Pro Leu Ala Phe Val
1 5 10 15

Ser Ser Ser Ala Leu Pro Ile His Asp Val Asp Ser Gln Gln Asn Thr
20 25 30

Ser Gly Phe Leu Gly Leu Gln Arg Leu Leu Gln Ser Phe Ser Arg Leu
35 40 45

Phe Leu Lys Asn Asp Leu Leu Arg Asp Leu Asp Asn Phe Phe Ser Ser
50 55 60

Pro Met Asp Phe Arg Asp Leu Pro Arg Asn Phe His Gln Glu Glu Asn
65 70 75 80

Gln Glu His Arg Met Gly Asn His Thr Leu Ser Ser His Leu Gln Ile
85 90 95

Asp Lys Val Thr Asp Asn Gln Thr Gly Glu Val His Ile Ser Glu Lys
100 105 110

Val Glu Ala Ser Ile Glu Pro Glu Arg Asn Pro Glu Gly Asp Trp Lys
115 120 125

Val Pro Lys Val Glu Ala Lys Glu Pro Pro Val Pro Val Gln Lys Val
130 135 140

Thr Asp Ser Leu His Pro Glu Pro Arg Gln Val Ala Phe Trp Ile Met
145 150 155 160

Lys Met Pro Arg Arg Arg Thr Gln Pro Asp Val Gln Asp Gly Gly Arg

- 35 -

	165		170		175
Trp Leu Ile	Glu Lys Arg His Arg Met Gln Ala Ile Arg Asp Gly Leu				
	180		185		190
Arg Gly Gly	Ala Arg Glu Asp Ser Leu Glu Asp Gly Val His Ile Pro				
	195		200		205
Gln His Ala	Lys Leu Pro Val Arg Lys Thr His Phe Leu Tyr Ile Leu				
	210		215		220
Arg Pro Ser	Gln Gln Leu				
	225		230		

<210> 28
 <211> 690
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (1)..(690)

<400> 28
 atg tgt cga ctg agg gtc ttg ctg ctg ctg ctc ccc ttg gcc ttc gtg 48
 Met Cys Arg Leu Arg Val Leu Leu Leu Leu Leu Pro Leu Ala Phe Val
 1 5 10 15

tcc tcc tct gct ctc ccc atc cat gat gtc gac tct cag cag aac acc 96
 Ser Ser Ser Ala Leu Pro Ile His Asp Val Asp Ser Gln Gln Asn Thr
 20 25 30

tcc ggg ttc ctg ggc ctt cag agg ctt ctc caa agc ttt agt cga ctg 144
 Ser Gly Phe Leu Gly Leu Gln Arg Leu Leu Gln Ser Phe Ser Arg Leu
 35 40 45

ttc cta aaa aat gac ctg cta cga gac ctg gac aac ttc ttc tcc tcc 192
 Phe Leu Lys Asn Asp Leu Leu Arg Asp Leu Asp Asn Phe Phe Ser Ser
 50 55 60

ccc atg gac ttc cga gac ctt cct agg aac ttc cat cag gaa gag aac 240
 Pro Met Asp Phe Arg Asp Leu Pro Arg Asn Phe His Gln Glu Glu Asn
 65 70 75 80

cag gag cac aga atg ggc aac cat acc ctc tcc agc cac cta cag ata 288
 Gln Glu His Arg Met Gly Asn His Thr Leu Ser Ser His Leu Gln Ile
 85 90 95

gac aag gtg act gac aac cag aca ggg gag gtg cac atc tcg gag aaa 336
 Asp Lys Val Thr Asp Asn Gln Thr Gly Glu Val His Ile Ser Glu Lys
 100 105 110

gtc gag gcc tcc att gag cca gaa cgg aac ccg gaa ggg gac tgg aag 384
 Val Glu Ala Ser Ile Glu Pro Glu Arg Asn Pro Glu Gly Asp Trp Lys
 115 120 125

gtt ccc aaa gta gaa gca aaa gag ccc ccg gtg cct gtg cag aag gtc 432
 Val Pro Lys Val Glu Ala Lys Glu Pro Pro Val Pro Val Gln Lys Val

- 36 -

130	135	140	
acc gac agc ttg cac cca gag ccc cgg cag gtg gct ttc tgg atc atg			480
Thr Asp Ser Leu His Pro Glu Pro Arg Gln Val Ala Phe Trp Ile Met			
145	150	155	160
aag atg cca agg cgg agg acc cag ccc gat gtc cag gat gga ggc cgc			528
Lys Met Pro Arg Arg Arg Thr Gln Pro Asp Val Gln Asp Gly Gly Arg			
	165	170	175
tgg ctc ata gaa aag cga cat cgc atg cag gcc atc cgg gat ggg ctc			576
Trp Leu Ile Glu Lys Arg His Arg Met Gln Ala Ile Arg Asp Gly Leu			
	180	185	190
cgt gga ggc gcc cgt gag gac agc ctg gag gat ggg gtc cat atc ccc			624
Arg Gly Gly Ala Arg Glu Asp Ser Leu Glu Asp Gly Val His Ile Pro			
	195	200	205
caa cac gcc aag ctg cct gtc aga aag aca cac ttt ctc tac atc ctc			672
Gln His Ala Lys Leu Pro Val Arg Lys Thr His Phe Leu Tyr Ile Leu			
	210	215	220
agg cca tcc caa cag ctg			690
Arg Pro Ser Gln Gln Leu			
225	230		

<210> 29

<211> 51

<212> PRT

<213> Artificial Sequence

<220>

<223> Xaa at positions

3-5, 7, 9-15, 18, 20-22, 24-27, 29, 31, 33, 34, 36-39, 42, 44,
45, and 47-50 may be any amino acid

<220>

<223> Description of Artificial Sequence: consensus sequence

<400> 29

Leu	Pro	Xaa	Xaa	Xaa	His	Xaa	Glu	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Gly
1				5					10						15

Asn	Xaa	Thr	Xaa	Xaa	Xaa	His	Xaa	Xaa	Xaa	Xaa	Lys	Xaa	Thr	Xaa	Asn
			20					25					30		

Xaa	Xaa	Gly	Xaa	Xaa	Xaa	Xaa	Ser	Glu	Xaa	Val	Xaa	Xaa	Ser	Xaa	Xaa
		35					40					45			

Xaa	Xaa	Glu
		50

<210> 30

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 30

cagtgagtgc tgtggagacc

20

<210> 31

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 31

tcttcagtca ggctcctctc

20

<210> 32

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 32

acctgcaatg tgtcgactga g

21

<210> 33

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 33

cacttacagc tggtgggatg

20

<210> 34

<211> 10

<212> PRT

<213> Homo sapiens

<220>

<223> Xaa at position 1 may be any amino acid

<400> 34

Xaa Val Leu Asp Phe Asn Asn Ile Arg Ser
1 5 10

<210> 35

<211> 10

- 38 -

<212> PRT

<213> Homo sapiens

<400> 35

Ser Gln Gly Arg Lys Gly Gln Glu Gly Ser
 1 5 10

<210> 36

<211> 272

<212> PRT

<213> Mus musculus

<400> 36

Met Met Val Val Cys Ala Pro Ala Ala Val Arg Phe Leu Ala Val Phe
 1 5 10 15

Thr Met Met Ala Leu Cys Ser Leu Pro Leu Leu Gly Ala Ser Ala Thr
 20 25 30

Leu Asn Ser Val Leu Ile Asn Ser Asn Ala Ile Lys Asn Leu Pro Pro
 35 40 45

Pro Leu Gly Gly Ala Gly Gly Gln Pro Gly Ser Ala Val Ser Val Ala
 50 55 60

Pro Gly Val Leu Tyr Glu Gly Gly Asn Lys Tyr Gln Thr Leu Asp Asn
 65 70 75 80

Tyr Gln Pro Tyr Pro Cys Ala Glu Asp Glu Glu Cys Gly Ser Asp Glu
 85 90 95

Tyr Cys Ser Ser Pro Ser Arg Gly Ala Ala Gly Val Gly Gly Val Gln
 100 105 110

Ile Cys Leu Ala Cys Arg Lys Arg Arg Lys Arg Cys Met Thr His Ala
 115 120 125

Met Cys Cys Pro Gly Asn Tyr Cys Lys Asn Gly Ile Cys Met Pro Ser
 130 135 140

Asp His Ser His Phe Pro Arg Gly Glu Ile Glu Glu Ser Ile Ile Glu
 145 150 155 160

Asn Leu Gly Asn Asp His Asn Ala Ala Ala Gly Asp Gly Tyr Pro Arg
 165 170 175

Arg Thr Thr Leu Thr Ser Lys Ile Tyr His Thr Lys Gly Gln Glu Gly
 180 185 190

Ser Val Cys Leu Arg Ser Ser Asp Cys Ala Ala Gly Leu Cys Cys Ala
 195 200 205

Arg His Phe Trp Ser Lys Ile Cys Lys Pro Val Leu Lys Glu Gly Gln
 210 215 220

Val Cys Thr Lys His Lys Arg Lys Gly Ser His Gly Leu Glu Ile Phe
 225 230 235 240

- 39 -

Gln Arg Cys Tyr Cys Gly Glu Gly Leu Ala Cys Arg Ile Gln Lys Asp
245 250 255

His His Gln Ala Ser Asn Ser Ser Arg Leu His Thr Cys Gln Arg His
260 265 270

<210> 37

<211> 259

<212> PRT

<213> Xenopus laevis

<400> 37

Met Gly Ser Asn Met Phe Pro Val Pro Leu Ile Val Phe Trp Gly Phe
1 5 10 15

Ile Leu Asp Gly Ala Leu Gly Phe Val Met Met Thr Asn Ser Asn Ser
20 25 30

Ile Lys Asn Val Pro Ala Ala Pro Ala Gly Gln Pro Ile Gly Tyr Tyr
35 40 45

Pro Val Ser Val Ser Pro Asp Ser Leu Tyr Asp Ile Ala Asn Lys Tyr
50 55 60

Gln Pro Leu Asp Ala Tyr Pro Leu Tyr Ser Cys Thr Glu Asp Asp Asp
65 70 75 80

Cys Ala Leu Asp Glu Phe Cys His Ser Ser Arg Asn Gly Asn Ser Leu
85 90 95

Val Cys Leu Ala Cys Arg Lys Arg Arg Lys Arg Cys Leu Arg Asp Ala
100 105 110

Met	Cys	Cys	Thr	Gly	Asn	Tyr	Cys	Ser	Asn	Gly	Ile	Cys	Val	Pro	Val
		115					120					125			

Glu Gln Asp Gln Glu Arg Phe Gln His Gln Gly Tyr Leu Glu Glu Thr
130 135 140

Ile Leu Glu Asn Tyr Asn Asn Ala Asp His Ala Thr Met Asp Thr His
145 150 155 160

Ser Lys Leu Thr Thr Ser Pro Ser Gly Met Gln Pro Phe Lys Gly Arg
165 170 175

Asp Gly Asp Val Cys Leu Arg Ser Thr Asp Cys Ala Pro Gly Leu Cys
180 185 190

Cys Ala Arg His Phe Trp Ser Lys Ile Cys Lys Pro Val Leu Asp Glu
195 200 205

Gly Gln Val Cys Thr Lys His Arg Arg Lys Gly Ser His Gly Leu Glu
210 215 220

```

Ile Phe Gln Arg Cys His Cys Gly Ala Gly Leu Ser Cys Arg Leu Gln
225                                230                        235                240

```

Lys Gly Glu Phe Thr Thr Val Pro Lys Thr Ser Arg Leu His Thr Cys

245

250

255

Gln Arg His

<210> 38

<211> 350

<212> PRT

<213> Gallus gallus

<400> 38

Met Arg Arg Gly Glu Gly Pro Ala Pro Arg Arg Arg Trp Leu Leu Leu
 1 5 10 15

Leu Ala Val Leu Ala Ala Leu Cys Cys Ala Ala Ala Gly Ser Gly Gly
 20 25 30

Arg Arg Arg Ala Ala Ser Leu Gly Glu Met Leu Arg Glu Val Glu Ala
 35 40 45

Leu Met Glu Asp Thr Gln His Lys Leu Arg Asn Ala Val Gln Glu Met
 50 55 60

Glu Ala Glu Glu Glu Gly Ala Lys Lys Leu Ser Glu Val Asn Phe Glu
 65 70 75 80

Asn Leu Pro Pro Thr Tyr His Asn Glu Ser Asn Thr Glu Thr Arg Ile
 85 90 95

Gly Asn Lys Thr Val Gln Thr His Gln Glu Ile Asp Lys Val Thr Asp
 100 105 110

Asn Arg Thr Gly Ser Thr Ile Phe Ser Glu Thr Ile Ile Thr Ser Ile
 115 120 125

Lys Gly Gly Glu Asn Lys Arg Asn His Glu Cys Ile Ile Asp Glu Asp
 130 135 140

Cys Glu Thr Gly Lys Tyr Cys Gln Phe Ser Thr Phe Glu Tyr Lys Cys
 145 150 155 160

Gln Pro Cys Lys Thr Gln His Thr His Cys Ser Arg Asp Val Glu Cys
 165 170 175

Cys Gly Asp Gln Leu Cys Val Trp Gly Glu Cys Arg Lys Ala Thr Ser
 180 185 190

Arg Gly Glu Asn Gly Thr Ile Cys Glu Asn Gln His Asp Cys Asn Pro
 195 200 205

Gly Thr Cys Cys Ala Phe Gln Lys Glu Leu Leu Phe Pro Val Cys Thr
 210 215 220

Pro Leu Pro Glu Glu Gly Glu Pro Cys His Asp Pro Ser Asn Arg Leu
 225 230 235 240

Leu Asn Leu Ile Thr Trp Glu Leu Glu Pro Asp Gly Val Leu Glu Arg
 245 250 255

Cys	Pro	Cys	Ala	Ser	Gly	Leu	Ile	Cys	Gln	Pro	Gln	Ser	Ser	His	Ser	
			260					265					270			
Thr	Thr	Ser	Val	Cys	Glu	Leu	Ser	Ser	Asn	Glu	Thr	Arg	Lys	Asn	Glu	
		275					280					285				
Lys	Glu	Asp	Pro	Leu	Asn	Met	Asp	Glu	Met	Pro	Phe	Ile	Ser	Leu	Ile	
	290					295					300					
Pro	Arg	Asp	Ile	Leu	Ser	Asp	Tyr	Glu	Glu	Ser	Ser	Val	Ile	Gln	Glu	
305					310					315					320	
Val	Arg	Lys	Glu	Leu	Glu	Ser	Leu	Glu	Asp	Gln	Ala	Gly	Val	Lys	Ser	
				325					330					335		
Glu	His	Asp	Pro	Ala	His	Asp	Leu	Phe	Leu	Gly	Asp	Glu	Ile			
			340					345					350			

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 September 2000 (08.09.2000)

PCT

(10) International Publication Number
WO 00/52047 A3

(51) International Patent Classification⁷: **C07K 14/47**,
C12N 15/12, C07K 16/18, G01N 33/50, 33/53, C12Q
1/68, A61K 38/17

(74) Agents: MANDRAGOURAS, Amy, E.; Lahive & Cock-
field, LLP, 28 State Street, Boston, MA 02109 et al. (US).

(21) International Application Number: PCT/US00/05452

(81) Designated States (*national*): AU, CA, JP.

(22) International Filing Date: 3 March 2000 (03.03.2000)

(84) Designated States (*regional*): European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE).

(25) Filing Language: English

(26) Publication Language: English

Published:

(30) Priority Data:
09/263,022 5 March 1999 (05.03.1999) US

— With international search report.

(71) Applicant: MILLENNIUM PHARMACEUTICALS,
INC. [US/US]; 75 Sidney Street, Cambridge, MA 02139
(US).

(88) Date of publication of the international search report:
1 February 2001

(72) Inventor: MCCARTHY, Sean; 62 Commonwealth Av-
enue, Apartment #4, Boston, MA 02116 (US).

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*



WO 00/52047 A3

(54) Title: HUMAN DICKKOPF-RELATED PROTEIN AND NUCLEIC ACID MOLECULES AND USES THEREFOR

(57) Abstract: Novel Dkk and Dkk-related polypeptides, proteins, and nucleic acid molecules are disclosed. In addition to isolated, full-length Dkk and Dkk-related proteins, the invention further provides isolated fusion proteins, antigenic peptides and antibodies. The invention also provides Dkk and Dkk-related nucleic acid molecules, recombinant expression vectors containing a nucleic acid molecule of the invention, host cells into which the expression vectors have been introduced and non-human transgenic animals in which a Dkk and Dkk-related gene has been introduced or disrupted. Diagnostic, screening and therapeutic methods utilizing compositions of the invention are also provided.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/05452

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07K14/47 C12N15/12 C07K16/18 G01N33/50 G01N33/53
C12Q1/68 A61K38/17

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 98 46755 A (MCCARTHY SEAN A ;MILLENNIUM BIOTHERAPEUTICS INC (US)) 22 October 1998 (1998-10-22) Note: 100.0% nt seq identity of SEQ ID NO:1 (3) with SEQ ID NO:1 (3) in 2479 bp (1050 bp) overlap, 100.0% aa seq identity of SEQ ID NO:2 with SEQ ID NO:2 in 350 aa overlap. the whole document example 1 page 85, line 31 -page 86, line 7 page 89-95 claims 1-22</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	1-22



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

° Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

5 July 2000

Date of mailing of the international search report

18.10.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

van de Kamp, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/05452

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 98 27932 A (HUMAN GENOME SCIENCES INC ;RUBEN STEVEN M (US); SOPPET DANIEL R (U) 2 July 1998 (1998-07-02) Note: 99.7% nt seq identity of SEQ ID NO:1 with SEQ ID NO:1 (3) in 2445 bp (1050 bp) overlap, 99.1% aa seq identity of SEQ ID NO:2 with SEQ ID NO:2 in 350 aa overlap. the whole document page 2, line 22 -page 3, line 12 page 4, line 8-12 examples 1-4 page 45-49 claims 1-4,8-20 figures 1,2</p> <p>---</p>	1-22
A	<p>GLINKA ET AL: "Dickkopf-1 is a member of a new family of secreted proteins and functions in head induction" NATURE, vol. 391, no. 6665, 22 January 1998 (1998-01-22), pages 357-362, XP002140766 the whole document page 357, right-hand column, line 4-13; figure 1B</p> <p>---</p>	1-22
A	<p>FINCH P W ET AL.: "Purification and molecular cloning of a secreted, Frizzled-related antagonist of Wnt action" PROC. NATL. ACAD. SCI. USA, vol. 94, no. 13, June 1997 (1997-06), pages 6770-6775, XP002140787 abstract; figure 2</p> <p>---</p>	1-22
P,X	<p>KRUPNIK V E ET AL.: "Functional and structural diversity of the human Dickkopf gene family" GENE, vol. 238, no. 2, 1 October 1999 (1999-10-01), pages 301-313, XP002140768 Note: 99.9% (99.7%) nt seq identity of hdkk-3 with SEQ ID NO:1 (3) in 2479 bp (1050 bp) overlap, 99.1% aa seq identity of hDkk-3 with SEQ ID NO:2 in 350 aa overlap. the whole document page 304, left-hand column, line 21-34; figure 1</p> <p>---</p> <p>-/--</p>	1-22

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/05452

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 99 22000 A (DEUTSCHES KREBSFORSCH ;GLINKA ANDREI (DE); NIEHRS CHRISTOF (DE)) 6 May 1999 (1999-05-06) Note: 67.6% nt seq identity of SEQ ID NO:7 with SEQ ID NO:3 in 873 bp overlap, 65.4% aa seq identity of translated SEQ ID NO:7 with SEQ ID NO:2 in 254 aa overlap. the whole document claim 4A; figure 2 ---	1-22
E	WO 00 18194 A (ARNOLD & RICHTER KG ;BRAUCKMANN WILFRIED (DE); STEGMAIER KLAUS DIE) 30 March 2000 (2000-03-30) Note: 99.7% nt seq identity of SEQ ID NO:2 with SEQ ID NO:1 (3) in 1053 bp (1050 bp) overlap, 99.1% aa seq identity of SEQ ID NO:9 with SEQ ID NO:2 in 350 aa overlap. the whole document figures 2,9; examples 2,10,11 claims 1-20 -----	1-22

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 00/05452

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-22 all partially

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1-22 (all partially)

An isolated nucleic acid molecule: a) comprising a nucleotide (nt) sequence which is at least 60% homologous to a nt sequence of SEQ ID NO:1 or 3, or of the DNA insert of the plasmid with ATCC Accession No. 98452, or a complement thereof; or b) comprising at least 1000 nts from said nt sequences; or c) encoding a polypeptide comprising an amino acid (aa) sequence at least about 60% homologous to the aa sequence of SEQ ID NO:2, or to the aa sequence encoded by the insert of said plasmid; or d) encoding a fragment of at least 15 contiguous aa residues of said aa sequence; or e) encoding a naturally occurring allelic variant of said polypeptide, wherein the nucleic acid molecule hybridizes to a nucleic acid molecule comprising SEQ ID NO:1 or 3 or the DNA insert of said plasmid, under stringent conditions. An isolated nucleic acid molecule: f) comprising the nt sequence of SEQ ID NO:1 or 3, or of the DNA insert of the plasmid with ATCC Accession No. 98452; or g) encoding a polypeptide comprising the aa sequence of SEQ ID NO:2, or encoded by the DNA insert of said plasmid. Vectors, host cells. Isolated polypeptides analogous to said isolated nucleic acid molecules (a,c,d,e,g). Fusion proteins. Antibodies. Methods for producing polypeptides. Methods for detecting the presence of said polypeptides. Methods for detecting the presence of said nucleic acid molecules. Methods for identifying compounds which bind to or modulate said polypeptides. Methods for modulating the activity of said polypeptides.

2. Claims: 1-22 (all partially)

As invention 1, but concerning SEQ ID NO:4-6, and the plasmid with ATCC Accession No. _____.

3. Claims: 1-22 (all partially)

As invention 1, but concerning SEQ ID NO:7-9, and the plasmid with ATCC Accession No. 98633.

4. Claims: 1-22 (all partially)

As invention 1, but concerning SEQ ID NO:13-15, and the plasmid with ATCC Accession No. _____.

5. Claims: 1-22 (all partially)

As invention 1, but concerning SEQ ID NO:20-22, and the plasmid with ATCC Accession No. _____.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/05452

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9846755	A	22-10-1998	AU EP	7137398 A 0975755 A	11-11-1998 02-02-2000

WO 9827932	A	02-07-1998	AU EP	5613498 A 0954575 A	17-07-1998 10-11-1999

WO 9922000	A	06-05-1999	DE EP	19747418 C 1027440 A	15-07-1999 16-08-2000

WO 0018194	A	30-03-2000	DE	19844293 A	30-03-2000

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 September 2000 (08.09.2000)

PCT

(10) International Publication Number
WO 00/52047 A3

(51) International Patent Classification⁷: C07K 14/47,
C12N 15/12, C07K 16/18, G01N 33/50, 33/53, C12Q
1/68, A61K 38/17

(81) Designated States (*national*): AU, CA, JP.

(21) International Application Number: PCT/US00/05452

(84) Designated States (*regional*): European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE).

(22) International Filing Date: 3 March 2000 (03.03.2000)

Published:

(25) Filing Language: English

— with international search report

(26) Publication Language: English

(88) Date of publication of the international search report:
1 February 2001

(30) Priority Data:
09/263,022 5 March 1999 (05.03.1999) US

(48) Date of publication of this corrected version:
14 March 2002

(71) Applicant: MILLENNIUM PHARMACEUTICALS,
INC. [US/US]; 75 Sidney Street, Cambridge, MA 02139
(US).

(15) Information about Correction:
see PCT Gazette No. 11/2002 of 14 March 2002, Section
II

(72) Inventor: MCCARTHY, Sean; 62 Commonwealth Av-
enue, Apartment #4, Boston, MA 02116 (US).

(74) Agents: MANDRAGOURAS, Amy, E.; Lahive & Cock-
field, LLP, 28 State Street, Boston, MA 02109 et al. (US).

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: HUMAN DICKKOPF-RELATED PROTEIN AND NUCLEIC ACID MOLECULES AND USES THEREFOR

(57) Abstract: Novel Dkk and Dkk-related polypeptides, proteins, and nucleic acid molecules are disclosed. In addition to isolated, full-length Dkk and Dkk-related proteins, the invention further provides isolated fusion proteins, antigenic peptides and antibodies. The invention also provides Dkk and Dkk-related nucleic acid molecules, recombinant expression vectors containing a nucleic acid molecule of the invention, host cells into which the expression vectors have been introduced and non-human transgenic animals in which a Dkk and Dkk-related gene has been introduced or disrupted. Diagnostic, screening and therapeutic methods utilizing compositions of the invention are also provided.

WO 00/52047 A3



HUMAN DICKKOPF-RELATED PROTEIN AND NUCLEIC ACID MOLECULES AND USES THEREFOR

Background of the Invention

5 Secreted proteins play an integral role in the formation, differentiation, and maintenance of cells in multicellular organisms. For instance, secretory proteins are known in the art to be involved in signaling between cells which are not in direct contact. Such secreted signaling molecules are particularly important in the development of vertebrate tissue during embryogenesis as well as in the maintenance of
10 the differentiated state of adult tissues. For example, inductive interactions that occur between neighboring cell layers and tissues in the developing embryo are largely dependent on the existence and regulation of secreted signaling molecules. In inductive interactions, biochemical signals secreted by one cell population influence the developmental fate of a second cell population, typically by altering the fate of the
15 second cell population. For example, the Wnt proteins are now recognized as one of the major families of developmentally important signaling molecules in organisms ranging from *Drosophila* to mice.

 The Wnt gene family encode a large class of secreted proteins related to the *Int1/Wnt1* proto-oncogene and *Drosophila wingless* ("Wg"), a *Drosophila Wnt1*
20 homologue, (Cadigan *et al.* (1997) *Genes & Development* 11:3286-3305). Wnts are expressed in a variety of tissues and organs and are required for many developmental processes, including segmentation in *Drosophila*, endoderm development in *Caenorhabditis elegans*, establishment of limb polarity, neural crest differentiation, kidney morphogenesis, sex determination, and brain development in mammals
25 (reviewed in Parr and McMahon (1994) *Curr. Opinion Genetics & Devel.* 4:523-528; Cadigan and Nusse, *supra*).

 Recent studies in diverse organisms have led to identification of several components of the Wnt signal transduction pathway in responding cells (Cadigan and Nusse, *supra*). Wnt signals are transduced by the Frizzled ("Fz") family of seven
30 transmembrane domain receptors (Bhanot *et al.* (1996) *Nature* 382:225-230). The resulting signal leads to the activation of the cytoplasmic protein Dishevelled (Dsh) and

- 2 -

stabilization of Armadillo/ β -catenin (Perrimon (1994) *Cell* 76:781-784). Negative regulators of the Wnt pathway include glycogen synthase kinase 3 (GSK3)/shaggy (Perrimon, *supra*), the tumor suppressor gene product adenomatous polyposis coli (APC) (Gumbiner (1997) *Curr. Biol.* 7:R443-436) and a novel protein, called Axin (Zeng *et al.* (1997) *Cell* 90:181-192). In the absence of a Wnt ligand, these proteins promote phosphorylation and then degradation of β -catenin, whereas Wnt signaling inactivates GSK3, thus preventing β -catenin degradation. As a result, β -catenin is translocated to the nucleus, where it forms a complex with TCF transcription factors and activates target gene expression (Cadigan and Nusse, *supra*). Deregulation of this pathway can lead to carcinogenesis (reviewed by Gumbiner, *supra*), emphasizing the long-recognized connection between Wnts, normal development and cancer. This connection has been further established recently with the identification the c-Myc proto-oncogene as a target of Wnt signaling (He *et al.* (1998) *Science* 281:1509-3512).

While the outcome of Wnt signaling may be influenced by multiple intracellular regulatory mechanisms, recent studies have identified several classes of secreted factors which can modulate Wnt action outside of the cell. These include Cerberus, a secreted Wnt inhibitor implicated in head development (Bouwmeester *et al.* (1996) *Nature* 382:595-601), and a family of proteins related to the extracellular domain of Frizzled. These Frizzled-related proteins ("FRPs") (Rattner *et al.* (1997) *Proc. Natl. Acad. Sci. USA* 94:2859-2863), also known as secreted apoptosis-related proteins ("SARPs"), are encoded by several independently discovered genes including FrzA/FRP1, SDF5/FRP2, FrzB/FRP3, FRP4 and Sizzled (Melkonyan *et al.* (1997) *Proc. Natl. Acad. Sci. USA* 94:13636-13641; Finch *et al.* (1997) *Proc. Natl. Acad. Sci. USA* 94:6770-6775; Wang *et al.* (1997) *Cell* 88:747-766; Leyns *et al.* (1997) *Cell* 88:747-756; Mayr *et al.* (1997) *Mech. Dev.* 63:109-325; and Salic *et al.* (1997) *Development* 124:4739-4748). These proteins inhibit the ability of Xwnt8 to induce a secondary axis in frog embryos (for review see Zorn (1997) *Curr. Biol.* 7:R501-504), and are thought to compete for binding of Wnt ligands to the Frizzled receptors. Data on binding of certain FRPs to Xwnt8 (Wang *et al.*, (1997) *Biochem. Biophys. Res. Comm.* 236:502-504; and Leyns *et al.*, *supra*) and Wg corroborate this notion (Rattner *et al.*, *supra*).

- 3 -

It is now recognized that many of these families of signaling molecules have a dual role to play in both the development of an organism as well as in promoting or maintaining the differentiated state of tissues in the adult animal. Furthermore, major families of signaling molecules have been implicated in controlling proliferation of cells in mature adult tissue, for example, during normal cell turnover in the adult organism as well as in tissue regeneration activated as a result of damage to the adult tissue. Given the important role of these signalling molecules such as the Wnts and FRPs in both developing and adult tissues, there exists a need for identifying novel modulators of such molecules for use in regulating a variety of cellular processes.

Summary of the Invention

The present invention is based, at least in part, on the discovery of nucleic acid molecules which encode a novel family of secreted human proteins, referred to herein as the human Dickkopf proteins or "hDkks" (formerly referred to as the "Cysteine-Rich Secreted Proteins", "CRSPs", "CRISPYs", or "CRSP proteins). The Dkk molecules of the present invention are useful as modulating agents in regulating a variety of cellular processes. Accordingly, in one aspect, this invention provides isolated nucleic acid molecules encoding Dkk proteins or biologically active portions thereof, as well as nucleic acid fragments suitable as primers or hybridization probes for the detection of Dkk-encoding nucleic acids. In another aspect, this invention provides isolated nucleic acid molecules encoding Dkk-related proteins (*e.g.*, Soggy proteins) or biologically active portions thereof, as well as nucleic acid fragments suitable as primers or hybridization probes for the detection of Dkk- or Soggy-encoding nucleic acids.

In one embodiment, a Dkk nucleic acid molecule is 60% homologous to the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:3, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452 or complement thereof. In yet another embodiment, a Dkk nucleic acid molecule is 80% homologous to the nucleotide sequence shown in SEQ ID NO:4, SEQ ID NO:6, or a complement thereof. In yet another embodiment, a Dkk nucleic acid molecule is 60% homologous to the nucleotide sequence shown in SEQ ID NO:7, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as

- 4 -

Accession Number 98633, or a complement thereof. In yet another embodiment, a Dkk nucleic acid molecule is 85% homologous to the nucleotide sequence shown in SEQ ID NO:7, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or a complement thereof. In yet
5 another embodiment, a Dkk nucleic acid molecule is 70% homologous to the nucleotide sequence shown in SEQ ID NO:20, SEQ ID NO:22, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a complement thereof. In yet another embodiment, a nucleic acid molecule of the present invention (*e.g.*, a Dkk-related nucleic acid molecule) is 90% homologous to the
10 nucleotide sequence shown in SEQ ID NO:13, SEQ ID NO:15, or a complement thereof.

In a preferred embodiment, an isolated Dkk nucleic acid molecule has the nucleotide sequence shown SEQ ID NO:3, or a complement thereof. In another embodiment, a Dkk nucleic acid molecule further comprises nucleotides 1-37 of SEQ ID NO:1. In yet another preferred embodiment, a Dkk nucleic acid molecule further
15 comprises nucleotides 1088-2479 of SEQ ID NO:1. In another preferred embodiment, an isolated Dkk nucleic acid molecule has the nucleotide sequence shown in SEQ ID NO:1.

In another preferred embodiment, an isolated Dkk nucleic acid molecule has the nucleotide sequence shown SEQ ID NO:6, or a complement thereof. In another
20 embodiment, a Dkk nucleic acid molecule further comprises nucleotides 1-124 of SEQ ID NO:4. In yet another preferred embodiment, a Dkk nucleic acid molecule further comprises nucleotides 797-848 of SEQ ID NO:4. In another preferred embodiment, an isolated Dkk nucleic acid molecule has the nucleotide sequence shown in SEQ ID NO:4.

In another preferred embodiment, an isolated Dkk nucleic acid molecule has the
25 nucleotide sequence shown SEQ ID NO:9, or a complement thereof. In another embodiment, a Dkk nucleic acid molecule further comprises nucleotides 1-108 of SEQ ID NO:7. In yet another preferred embodiment, a Dkk nucleic acid molecule further comprises nucleotides 907-1536 of SEQ ID NO:7. In another preferred embodiment, an isolated Dkk nucleic acid molecule has the nucleotide sequence shown in SEQ ID NO:7.

In another preferred embodiment, an isolated Dkk nucleic acid molecule has the nucleotide sequence shown SEQ ID NO:22, or a complement thereof. In another embodiment, a Dkk nucleic acid molecule further comprises nucleotides 1-723 of SEQ ID NO:20. In yet another preferred embodiment, a Dkk nucleic acid molecule further
5 comprises nucleotides 1501-3687 of SEQ ID NO:20. In yet another preferred embodiment, an isolated Dkk nucleic acid molecule has the nucleotide sequence shown in SEQ ID NO:20.

In another preferred embodiment, an isolated nucleic acid molecule of the present invention (*e.g.*, a Dkk-related nucleic acid molecule) has the nucleotide sequence
10 shown SEQ ID NO:15, or a complement thereof. In another embodiment, a nucleic acid molecule further comprises nucleotides 1-74 of SEQ ID NO:13. In yet another preferred embodiment, a nucleic acid molecule further comprises nucleotides 801-928 of SEQ ID NO:13. In yet another preferred embodiment, an isolated nucleic acid molecule has the nucleotide sequence shown in SEQ ID NO:13.

15 In another embodiment, a Dkk nucleic acid molecule includes a nucleotide sequence encoding a protein having an amino acid sequence sufficiently homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:21. In another preferred embodiment, a Dkk nucleic acid molecule includes a nucleotide sequence encoding a protein having an amino acid sequence at least 60%
20 homologous to the amino acid sequence of SEQ ID NO:2. In yet another preferred embodiment, a Dkk nucleic acid molecule includes a nucleotide sequence encoding a protein having an amino acid sequence at least 60% homologous to the amino acid sequence of SEQ ID NO:5. In yet another preferred embodiment, a Dkk nucleic acid molecule includes a nucleotide sequence encoding a protein having an amino acid
25 sequence at least 60% homologous to the amino acid sequence of SEQ ID NO:8. In yet another preferred embodiment, a Dkk nucleic acid molecule includes a nucleotide sequence encoding a protein having an amino acid sequence at least 75% homologous to the amino acid sequence of SEQ ID NO:8. In yet another preferred embodiment, a Dkk nucleic acid molecule includes a nucleotide sequence encoding a protein having an
30 amino acid sequence at least 65% homologous to the amino acid sequence of SEQ ID NO:21. In another embodiment, a nucleic acid molecule of the present invention (*e.g.*, a

- 6 -

Dkk-related nucleic acid molecule) includes a nucleotide sequence encoding a protein having an amino acid sequence sufficiently homologous to the amino acid sequence of SEQ ID NO:14 (*e.g.*, encodes a protein having an amino acid sequence which is 60% homologous to the amino acid sequence of SEQ ID NO:14).

5 In another embodiment, an isolated nucleic acid molecule of the present invention encodes a Dkk protein which includes a signal sequence and at least one cysteine-rich region, and is secreted. In another embodiment, an isolated nucleic acid molecule of the present invention encodes a Dkk protein which includes a signal sequence and a cysteine-rich region, wherein the cysteine-rich region comprises at least
10 one cysteine-rich domain, and is secreted. In yet another embodiment, a Dkk nucleic acid molecule encodes a Dkk protein and is a naturally occurring nucleotide sequence.

 In another embodiment, an isolated nucleic acid molecule of the present invention encodes a Dkk-related protein (*e.g.*, a Soggy protein) which includes a signal sequence, lacks cysteine-rich domains, and is secreted. In another embodiment, an
15 isolated nucleic acid molecule of the present invention encodes a Dkk-related protein (*e.g.*, a Soggy protein) which includes a signal sequence and a Soggy domain, and is secreted. In yet another embodiment, a nucleic acid molecule of the present invention encodes a Dkk-related protein and is a naturally occurring nucleotide sequence.

 Another embodiment of the invention features nucleic acid molecules which
20 specifically detect Dkk nucleic acid molecules relative to nucleic acid molecules encoding non-Dkk proteins (or specifically detect Dkk-related nucleic acid molecules). For example, in one embodiment, a nucleic acid molecule hybridizes under stringent conditions to a nucleic acid molecule consisting of nucleotides 470-2479 of nucleotide sequence shown in SEQ ID NO:1, to nucleotides 1-475 of nucleotide sequence shown in
25 SEQ ID NO:4, or to nucleotides 1-600 of nucleotide sequence shown in SEQ ID NO:7, or hybridizes under stringent conditions to the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, to the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or to the nucleotide sequence of the DNA insert of the plasmid deposited with
30 ATCC as Accession Number _____. In another embodiment, the nucleic acid molecule is at least 500 nucleotides in length and hybridizes under stringent conditions to a

nucleic acid molecule comprising the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20 or a complement thereof.

Another embodiment of the invention provides an isolated nucleic acid molecule which is antisense to the coding strand of a Dkk nucleic acid or Dkk-related nucleic acid. Another embodiment of the invention provides an isolated nucleic acid molecules in a form suitable for expression of mRNA. In another embodiment, the isolated nucleic acid molecules are in a form suitable for expression of protein. In yet another embodiment, the isolated nucleic acid molecules are free from vector sequences.

Another aspect of the invention provides a vector comprising a Dkk nucleic acid molecule or Dkk-related nucleic acid molecule. In certain embodiments, the vector is a recombinant expression vector. In another embodiment, the invention provides a host cell containing a vector of the invention. The invention also provides a method for producing a Dkk protein or Dkk-related protein by culturing in a suitable medium, a host cell of the invention containing a recombinant expression vector such that a Dkk protein or Dkk-related protein is produced.

Another aspect of this invention features isolated or recombinant Dkk proteins and polypeptides or Dkk-related proteins and polypeptides. In one embodiment, an isolated Dkk protein has a signal sequence and a cysteine-rich region which comprises two cysteine-rich domains, and is secreted. In another embodiment, an isolated Dkk protein has an amino acid sequence sufficiently homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:21. In a preferred embodiment, a Dkk protein has an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ ID NO:2. In another preferred embodiment, a Dkk protein has an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ ID NO:5. In another preferred embodiment, a Dkk protein has an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ ID NO:8. In another preferred embodiment, a Dkk protein has an amino acid sequence at least about 75% homologous to the amino acid sequence of SEQ ID NO:8. In another preferred embodiment, a Dkk protein has an amino acid sequence at least about 65% homologous to the amino acid sequence of SEQ ID NO:21. In another embodiment, a Dkk protein has the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID

NO:8, or SEQ ID NO:21. In another preferred embodiment, a protein of the present invention has an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ ID NO:14. In another embodiment, a protein has the amino acid sequence of SEQ ID NO:14.

5 Another embodiment of the invention features an isolated Dkk protein which is encoded by a nucleic acid molecule having a nucleotide sequence at least about 60% homologous to a nucleotide sequence of SEQ ID NO:1, or a complement thereof. Another embodiment of the invention features an isolated Dkk protein which is encoded by a nucleic acid molecule having a nucleotide sequence at least about 80% homologous
10 to a nucleotide sequence of SEQ ID NO:4, or a complement thereof. Another embodiment of the invention features an isolated Dkk protein which is encoded by a nucleic acid molecule having a nucleotide sequence at least about 60% homologous to a nucleotide sequence of SEQ ID NO:7, or a complement thereof. Another embodiment of the invention features an isolated Dkk protein which is encoded by a nucleic acid
15 molecule having a nucleotide sequence at least about 85% homologous to a nucleotide sequence of SEQ ID NO:7, or a complement thereof. Another embodiment of the invention features an isolated Dkk protein which is encoded by a nucleic acid molecule having a nucleotide sequence at least about 70% homologous to a nucleotide sequence of SEQ ID NO:20, or a complement thereof. Another embodiment of the invention
20 features an isolated protein which is encoded by a nucleic acid molecule having a nucleotide sequence at least about 90% homologous to a nucleotide sequence of SEQ ID NO:13, or a complement thereof. This invention further features an isolated protein which is encoded by a nucleic acid molecule having a nucleotide sequence which hybridizes under stringent hybridization conditions to a nucleic acid molecule
25 comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, or a complement thereof.

 The proteins of the present invention, or biologically active portions thereof, can be operatively linked to a non-Dkk polypeptide or non-Dkk-related polypeptide to form fusion proteins. The invention further features antibodies that specifically bind Dkk or
30 Dkk-related proteins, such as monoclonal or polyclonal antibodies. In addition, the

- 9 -

proteins or biologically active portions thereof can be incorporated into pharmaceutical compositions, which optionally include pharmaceutically acceptable carriers.

In another aspect, the present invention provides a method for detecting Dkk expression (or the expression of a Dkk-related molecule) in a biological sample by
5 contacting the biological sample with an agent capable of detecting a nucleic acid molecule, protein or polypeptide of the present invention such that the presence of a Dkk (of Dkk-related) nucleic acid molecule, protein or polypeptide is detected in the biological sample.

In another aspect, the present invention provides a method for detecting the
10 presence of a Dkk activity (or Dkk-related activity) in a biological sample by contacting the biological sample with an agent capable of detecting an indicator of Dkk activity (or Dkk-related activity) such that the presence of the activity is detected in the biological sample.

In another aspect, the invention provides a method for modulating Dkk activity
15 (or Dkk-related activity) comprising contacting the cell with an agent that modulates the activity such that the activity in the cell is modulated. In one embodiment, the agent inhibits Dkk activity (or Dkk-related activity). In another embodiment, the agent stimulates Dkk activity (or Dkk-related activity). In one embodiment, the agent is an antibody that specifically binds to a Dkk (or Dkk-related) protein. In another
20 embodiment, the agent modulates expression of a protein (*e.g.*, a Dkk or a Dkk-related protein) by modulating transcription of a gene or translation of a mRNA of the present invention. In yet another embodiment, the agent is a nucleic acid molecule having a nucleotide sequence that is antisense to the coding strand of a mRNA or gene of the present invention.

In one embodiment, the methods of the present invention are used to treat a
25 subject having a disorder characterized by aberrant expression or activity of a protein or nucleic acid of the invention by administering to the subject an agent which is a modulator of Dkk or a Dkk-related molecule. In one embodiment, the modulator is a Dkk or Dkk-related protein. In another embodiment the modulator is a Dkk or Dkk-
30 related nucleic acid molecule. In yet another embodiment, the modulator is an antibody peptide, peptidomimetic, or other small molecule. In a preferred embodiment, the

disorder characterized by aberrant protein or nucleic acid expression is a developmental, differentiative, or proliferative disorder.

The present invention also provides a diagnostic assay for identifying the presence or absence of a genetic alteration characterized by at least one of (i) aberrant
5 modification or mutation of a gene encoding a Dkk or Dkk-related protein; (ii) mis-regulation of said gene; and (iii) aberrant post-translational modification of a Dkk or Dkk-related protein, wherein a wild-type form of said gene encodes an protein with a Dkk or Dkk-related activity.

In another aspect the invention provides a method for identifying a compound
10 that binds to or modulates the activity of a Dkk or Dkk-related protein, by providing a indicator composition comprising a Dkk or Dkk-related protein having a biological activity, contacting the indicator composition with a test compound, and determining the effect of the test compound on the activity in the indicator composition to identify a compound that modulates the activity of a Dkk or Dkk-related protein.

15 Other features and advantages of the invention will be apparent from the following detailed description and claims.

Brief Description of the Drawings

Figure 1A-B depicts the cDNA sequence and predicted amino acid sequence of
20 human Dkk-3. The nucleotide sequence corresponds to nucleic acids 1 to 2479 of SEQ ID NO:1. The amino acid sequence corresponds to amino acids 1 to 350 of SEQ ID NO:2.

Figure 2 depicts the cDNA sequence and predicted amino acid sequence of
human Dkk-4. The nucleotide sequence corresponds to nucleic acids 1 to 848 of SEQ
25 ID NO:4. The amino acid sequence corresponds to amino acids 1 to 224 of SEQ ID NO:5.

Figure 3 depicts the cDNA sequence and predicted amino acid sequence of
human Dkk-1. The nucleotide sequence corresponds to nucleic acids 1 to 1536 of SEQ
ID NO:7. The amino acid sequence corresponds to amino acids 1 to 266 of SEQ ID
30 NO:8.

Figure 4A-B depicts the cDNA sequence and predicted amino acid sequence of full-length human Dkk-2. The nucleotide sequence corresponds to nucleic acids 1 to 3687 of SEQ ID NO:20. The amino acid sequence corresponds to amino acids 1 to 259 of SEQ ID NO:21.

5 *Figure 5A-B* depicts the cDNA sequence and predicted amino acid sequence of murine Dkk-3. The nucleotide sequence corresponds to nucleic acids 1 to 2380 of SEQ ID NO:16. The amino acid sequence corresponds to amino acids 1 to 349 of SEQ ID NO:17.

10 *Figure 6* depicts a multiple sequence alignment of the amino acid sequences of hDkk-1 (corresponding the SEQ ID NO:8), mDkk-1 (having Accession No. AF030433), *Xenopus* Dkk-1 ("xDkk-1") (having Accession No. AF030434), hDkk-2 (corresponding to SEQ ID NO:21), hDkk-3 (corresponding to SEQ ID NO:2), mDkk-3 (corresponding to SEQ ID NO:17), chicken Dkk-3 ("cDkk-3") (having Accession No. D26311), and hDkk-4 (corresponding to SEQ ID NO:5). The alignment was performed using the
15 ClustalW algorithm as implemented in the GCG program PILEUP. The alignment provides information regarding the relationship between the Dkk proteins of the instant invention. Predicted signal peptides are underlined, N-glycosylation sites are indicated by a thick bar, CRD-1 by an open box, CRD-2 by a shaded box. The proteolytic cleavage site within hDkk4 is indicated by an arrow.

20 *Figure 7* depicts the cDNA sequence and predicted amino acid sequence of human Soggy. The nucleotide sequence corresponds to nucleic acids 1 to 928 of SEQ ID NO:13. The amino acid sequence corresponds to amino acids 1 to 242 of SEQ ID NO:14.

25 *Figure 8* depicts the cDNA sequence and predicted amino acid sequence of murine Soggy-1. The nucleotide sequence corresponds to nucleic acids 1 to 835 of SEQ ID NO:26. The amino acid sequence corresponds to amino acids 1 to 230 of SEQ ID NO:27.

30 *Figure 9* is a schematic diagram illustrating the Dkk and Dkk-related proteins of the instant invention. The figure depicts the structural domains of the human Dkks and Soggy. Signal peptides are indicated by darkened boxes. The cysteine-rich domains of

a Dkk cysteine-rich region are depicted as CRD-1 and CRD-2. Branches indicate sites of N-glycosylation.

Figure 10 depicts a multiple sequence alignment of hSoggy-1 (corresponding to SEQ ID NO:14), murine Soggy-1 (corresponding to SEQ ID NO:27), hDkk-3 (corresponding to SEQ ID NO:2), and mDkk-3 (corresponding to SEQ ID NO:17). The alignment was generated as described in the legend to Figure 6. The alignment provides details regarding the relationship between the Dkk-3 and Soggy-1 proteins of the instant invention. Predicted signal peptides are underlined, N-glycosylation sites are indicated by a thick bar. CRD-1 and CRD-2 within Dkk-3 are indicated for reference by open and shaded boxes.

Figure 11 depicts a multiple sequence alignment of the carboxy-terminal cysteine-rich domains of hDkk-1, hDkk-2, hDkk-3, hDkk-4 with human colipase (having accession No. J02883). The carboxy-terminal cysteine-rich domains of the Dkk proteins are indicated by an open box. The alignment was generated using PILEUP (gap penalties of 12 for opening and 12 for extending). A minor adjustment was necessary since PILEUP inserts a single gap in hDkk-1 and hDkk-2 between Gly56 and Ser57, even with a gap opening penalty of 15. The conserved residues are indicated. The disulfide-bonding pattern typical for the colipase family and predicted for the Dkk family is indicated below the alignment.

Figure 12 is a schematic diagram depicting the relationship between the hDkk-3 nucleotide sequence (corresponding to SEQ ID NO:1) and those of RIG and RIG-like 7-1 (Accession Nos. U32331 and AF034208, respectively). Thick bars indicate regions of sequence identity between hDkk-3 and RIG or RIG-like 7-1 mRNAs. As between RIG and hDkk-3, there exists a short region of identity within the 3' untranslated regions of the mRNAs when the mRNAs are aligned in reverse orientation. As between hDkk-3 and RIG-like 7-1, there exists a longer region of identity, however, RIG-like 7-1 lacks a signal sequence and, accordingly, is not predicted to be secreted.

Detailed Description of the Invention

The present invention is based on the discovery of novel molecules, referred to herein as Dkk protein and nucleic acid molecules, which comprise a family of molecules

having certain conserved structural and functional features. The term "family" when referring to the protein and nucleic acid molecules of the invention is intended to mean two or more proteins or nucleic acid molecules having a common structural domain and having sufficient amino acid or nucleotide sequence homology as defined herein. Such

5 family members can be naturally-occurring and can be from either the same or different species. For example, a family can contain a first protein of human origin, as well as other, distinct proteins of human origin or alternatively, can contain homologues of non-human origin. Members of a family may also have common functional characteristics.

In one embodiment, a Dkk family member is identified based on the presence of

10 at least one "cysteine-rich domain" in the protein molecule or corresponding amino acid sequence. As defined herein, a "cysteine-rich domain" refers to a portion of a Dkk protein (*e.g.*, hDkk-3) which is rich in cysteine residues. In a preferred embodiment, a "cysteine-rich domain" is a protein domain having an amino acid sequence of about 45-85 amino acids of which preferably 10 amino acids are cysteine residues located at the

15 same relative amino acid position as the cysteine residues in human Dkk-3 having SEQ ID NO:2 (*e.g.*, amino acid residues 147-195 of SEQ ID NO:2). In another embodiment, a "cysteine-rich domain" has 30-100 amino acids, preferably about 35-95 amino acids, more preferably about 40-90 amino acids, more preferably about 50-80 amino acids, even more preferably about 55-75, 60-70, or 65 amino acids, of which at least about 3-

20 20, preferably about 5-15, or more preferably about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 amino acids are cysteine residues.

A preferred Dkk protein of the present invention has a first cysteine-rich domain ("CRD-1") referred to herein as an "amino-terminal cysteine-rich domain" or "N-terminal cysteine-rich domain" and a second cysteine-rich domain ("CRD-2"), referred

25 to herein as a "carboxy-terminal cysteine-rich domain" or "C-terminal cysteine-rich domain". As defined herein, an "amino-terminal cysteine-rich domain" is a protein domain having an amino acid sequence of about 45-55 amino acids of which preferably 10 amino acids are cysteine residues located at the same relative position as the cysteine residues in an amino-terminal cysteine-rich domain of human Dkk-3 having SEQ ID

30 NO:2 (*e.g.*, amino acid residues 147-195 of SEQ ID NO:2). In another embodiment, an "amino-terminal cysteine-rich domain" has 30-70, preferably 35-65, more preferably

about 40-60, and even more preferably about 46, 47, 48, 49, 50, 51, 52, 53, or 54 amino acids, of which at least about 3-20, preferably about 5-15, or more preferably about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 amino acids are cysteine residues. In a preferred embodiment, an amino-terminal cysteine-rich domain has the following

5 consensus sequence: C - X(2) - D - X(2) - C - X(5) - C - X(8-13) - C - X(2) - C - X(6) - C - X(5) - C - C - X(4) - C - X(4) - C (SEQ ID NO:23). The consensus sequences described herein are described according to standard Prosite Signature designation (*e.g.*, all amino acids are indicated according to their universal single letter designation; X designates any amino acid; X(n) designates any n amino acids, *e.g.*, X (2) designates any

10 2 amino acids; and [LIVM] indicates any one of the amino acids appearing within the brackets, *e.g.*, any one of L, I, V, or M, in the alternative, any one of Leu, Ile, Val, or Met.)

As defined herein, a "carboxy-terminal cysteine-rich domain" is a protein domain having an amino acid sequence of about 80-85 amino acids of which preferably 10

15 amino acids are cysteine residues located at the same relative position as the cysteine residues in a carboxy-terminal cysteine-rich domain of human Dkk-3 having SEQ ID NO:2 (*e.g.*, amino acid residues 201-284 of SEQ ID NO:2). In another embodiment, a "carboxy-terminal cysteine-rich domain" has 65-100, preferably 70-95, more preferably about 75-90, and even more preferably about 81, 82, 83, or 84 amino acids, of which at

20 least about 3-20, preferably about 5-15, or more preferably about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 amino acids are cysteine residues. In a preferred embodiment, a carboxy-terminal cysteine-rich domain has the following consensus sequence: C - X(4) - D - C - X(2) - G - X - C - C - X(8-10) - C - X - P - X(4) - G - X(2) - C - X(16-24) - C - X - C - X(2) - P - X(4) - G - X(2) - C - X(16-24) - C - X - C - X(2) -

25 G - L - X - C - X(10-17) - C (SEQ ID NO:24).

A preferred protein of the present invention is a hDkk-3 protein (human Dkk-3) containing an amino-terminal cysteine-rich domain including about amino acids 147-195 of SEQ ID NO:2, having 10 cysteine residues, and a carboxy-terminal cysteine-rich domain including about amino acids 201-284 of SEQ ID NO:2, having 10 cysteine

30 residues (the positions of the cysteine residues are depicted in Figure 6). In another embodiment, a hDkk-4 (human Dkk-4) protein contains an amino-terminal cysteine-rich

domain including about amino acids 41-90 of SEQ ID NO:5, having 10 cysteine residues, and a carboxy-terminal cysteine-rich domain including about amino acids 138-218 of SEQ ID NO:5, having 10 cysteine residues (the positions of the cysteine residues are depicted in Figure 6). In another embodiment, a hDkk-1 protein (human Dkk-1) contains an amino-terminal cysteine-rich domain including about amino acids 85-138 of SEQ ID NO:8, having 10 cysteine residues, and a carboxy-terminal cysteine-rich domain including about amino acids 182-263 of SEQ ID NO:8, having 10 cysteine residues (the positions of the cysteine residues are depicted in Figure 6). In another embodiment, a hDkk-2 protein (human Dkk-2) contains an amino-terminal cysteine-rich domain including about amino acids 78-127 of SEQ ID NO:21, having 10 cysteine residues, and a carboxy-terminal cysteine-rich domain including about amino acids 176-256 of SEQ ID NO:21, having 10 cysteine residues (the positions of the cysteine residues are depicted in Figure 6).

Alignment of the human Dkk proteins with human colipase (having Accession No. J02883) indicates that the carboxy-terminal cysteine-rich domains of the human Dkk proteins have a pattern of cysteines typical of colipase (Figure 11 and Avarind and Koonin, *supra*). Within colipase, these cysteine residues are involved in disulfide bonding which gives rise to a structure termed the "colipase fold". The "colipase fold" is typical of a range of small proteins which are involved in protein-protein interactions including, but not limited to the colipases, snake and scorpion toxins and protease inhibitors (Hubbard *et al.* (1997) *Nucleic Acids Res.* 25:236-239. These proteins have a series of short β strands with large connecting loops, which are held together by disulfide bonds. The disulfide-bonding pattern typical for colipase and predicted for the Dkk family is indicated below the alignment of Figure 11. Conserved hydrophobic residues between the Dkks and human colipase suggest that the Dkks, like the colipases, interact with lipids (*e.g.*, Leu51 of human colipase, SEQ ID NO:25 which corresponds to Leu271 of hDkk-3 (SEQ ID NO:2); Leu200 of hDkk-4 (SEQ ID NO:5); Leu243 of hDkk-1 (SEQ ID NO:8); and Leu237 of hDkk-2 (SEQ ID NO:21). The carboxy-terminal cysteine-rich domain of the Dkk family, may function in the membrane association of Dkk, which in turn may be required for the inhibition of Wnt secretion or Wnt:7 transmembrane receptor interaction. In addition, inhibition of Wnt function by

- 16 -

the Dkk family may be closely associated with the cell membrane and the carboxy-terminal cysteine-rich domain of the Dkk family may mediate this association.

Furthermore, the amino-terminal cysteine-rich domain of the Dkk family may directly interact with Wnt or its receptor. Accordingly, a preferred Dkk protein of the present invention comprises a carboxy-terminal cysteine-rich domain. In one embodiment, a Dkk protein comprising a carboxy-terminal cysteine-rich domain lacks the amino-terminal cysteine-rich domain.

In a preferred embodiment, the cysteine residues of a cysteine-rich domain are located at the same relative amino acid position as the cysteine residues in human Dkk-3 having SEQ ID NO:2. In another preferred embodiment, the cysteine residues of a cysteine-rich domain are located at the same relative position as the cysteine residues in a cysteine-rich domain of human Dkk-3 having SEQ ID NO:2. For example, as shown in Figure 6, human Dkk-4 has at least about 10 cysteine residues located at the same relative amino acid position as the cysteine residues in human Dkk-3 having SEQ ID NO:2 (*e.g.*, cys151 in Dkk-4, SEQ ID NO:5, is located at the same relative amino acid position as cys214 in Dkk-3, SEQ ID NO:2; cys156 in Dkk-4, SEQ ID NO:5, is located at the same relative amino acid position as cys219 in Dkk-3, SEQ ID NO:2; and cys157 in Dkk-4, SEQ ID NO:5, is located at the same relative amino acid position as cys220 in Dkk-3, SEQ ID NO:2). Similarly, as shown in Figure 6, Dkk-1 has at least about 10 cysteine residues located at the same relative amino acid position as the cysteine residues in human Dkk-3 having SEQ ID NO:2. As also shown in Figure 6, Dkk-2 has at least about 10 cysteine residues located at the same relative amino acid position as the cysteine residues in human Dkk-3 having SEQ ID NO:2. Table I sets forth at least 20 cysteine residues in each of hDkk-4, hDkk-1, and hDkk-2 which are located in the same relative position as 20 cysteine residues in hDkk-3.

Table I:

cysteine	aa position in hDkk-3	aa position in hDkk-4	aa position in hDkk-1	aa position in hDkk-2
1	147	41	85	78
2	153	47	91	84
3	159	53	97	90
4	168	63	111	100
5	171	66	114	103
6	178	73	121	110
7	184	79	127	116
8	185	80	128	117
9	190	85	133	122
10	195	90	138	127
11	208	145	189	183
12	214	151	195	189
13	219	156	200	194
14	220	157	201	195
15	231	166	210	204
16	241	176	220	214
17	265	194	237	231
18	267	196	239	233
19	273	202	245	239
20	284	218	263	256

The first 10 rows of Table I contain 10 cysteine residues that are included within
5 the first, or amino-terminal, cysteine-rich domain of each of hDkks-3, -4, -1, and -2.
The last 10 rows of Table I contain 10 cysteine residues that are included within the
second, or carboxy-terminal, cysteine-rich domain of each of hDkks-3, -4, -1, and -2.

Preferred Dkk proteins have more than one cysteine-rich domain, more preferably have at least two cysteine-rich domains and, thus, have a cysteine-rich region. As used herein, the term "cysteine-rich region" refers to a protein domain which includes at least two cysteine-rich domains and has an amino acid sequence of about 120-200 amino acid residues of which at least about 20 of the amino acids are cysteine residues. In another embodiment, a "cysteine-rich region" has preferably about 140-180 amino acid residues, and even more preferably at least about 135-175 amino acids of which at least about 10-30, preferably about 15-20, and more preferably about 16, 17, 18, or 19 of the amino acids are cysteine residues. In a preferred embodiment, a cysteine-rich region is located in the C-terminal region of a Dkk protein. For example, in one embodiment, a hDkk-3 protein contains a cysteine rich region containing about amino acids 147-284 of SEQ ID NO:2, having 20 cysteine residues at the positions indicated in Figure 6. In another embodiment, a hDkk-4 protein contains a cysteine rich region containing about amino acids 41-218 of SEQ ID NO:5, having 20 cysteine residues at the positions indicated in Figure 6. In another embodiment, a hDkk-1 protein contains a cysteine rich region containing about amino acids 85-263 of SEQ ID NO:8, having 20 cysteine residues at the positions indicated in Figure 6. In another embodiment, a hDkk-2 protein contains a cysteine rich region containing about amino acids 78-256 of SEQ ID NO:21, having 20 cysteine residues at the positions indicated in Figure 6.

In another embodiment, in addition to cysteine-rich domains, the cysteine-rich region contains a spacer region which separates the first and second cysteine-rich domains. As used herein, the "spacer region" refers to amino acid residues which are located between the first and second cysteine-rich domains of a cysteine-rich region and includes amino acid residues located C-terminal to the first cysteine-rich domain and N-terminal to the second cysteine-rich domain. As defined herein, a "spacer region" refers to a protein domain of about 5-70 amino acids, preferably about 10-65 amino acids, more preferably about 15-60 amino acids, even more preferably about 20-55 amino acids, and even more preferably about 25-50, 30-45 or 35-40 amino acids. For example, hDkk-3 protein contains a spacer region of about amino acids 196-200 of SEQ ID NO:2; hDkk-4 protein contains a spacer region of about amino acids 91-137 of SEQ ID NO:5; hDkk-1 protein contains a spacer region of about amino acids 139-181 of SEQ ID NO:8;

and hDkk-2 protein contains a spacer region of about amino acids 128-175 of SEQ ID NO:21. The spacer regions of hDkk-1, hDkk-2 and hDkk-4 are remarkably conserved in length (*e.g.*, the spacer region of hDkk-1 consists of 43 amino acid residues, the spacer region of hDkk-2 consists of 48 amino acid residues and the spacer region of hDkk-4 consists of 47 amino acid residues, suggesting that the close proximity of CRD-1 and CRD-2 is important in Dkk function. Accordingly, in one embodiment, the spacer region functions to spatially restrict the separation of CRD-1 from CRD-2.

In another embodiment of the invention, the Dkk protein has at least one cysteine-rich domain, preferably a cysteine-rich region, and a signal sequence. As used herein, a "signal sequence" refers to a peptide containing about 18-24 amino acids which occurs at the N-terminus of secretory and integral membrane proteins and which contains at least about 40-70% hydrophobic amino acid residues (*e.g.*, alanine, valine, leucine, isoleucine, phenylalanine, tyrosine, tryptophan, or proline). In another embodiment, a signal sequence contains at least about 8-34, 9-33, 10-32, 11-31, 12-30, 13-29, 14-28 amino acid residues, preferably about 15-27 amino acid residues, more preferably about 16-26 amino acid residues, more preferably about 17-25 amino acid residues, and more preferably about 18-24, 19-23, 20-22, or 21 amino acid residues, and has at least about about 50-65%, and more preferably about 55-60% hydrophobic amino acid residues (*e.g.*, alanine, valine, leucine, isoleucine, phenylalanine, tyrosine, tryptophan, or proline). Such a "signal sequence", also referred to in the art as a "signal peptide", serves to direct a protein containing such a sequence to a lipid bilayer. For example, in one embodiment, a hDkk-3 protein contains a signal sequence of about amino acids 1-23 of SEQ ID NO:2. In another embodiment, a hDkk-4 protein contains a signal sequence of about amino acids 1-19 of SEQ ID NO:5. In another embodiment, a hDkk-1 protein contains a signal sequence of about amino acids 1-20 of SEQ ID NO:8. In another embodiment, a hDkk-2 protein contains a signal sequence of about amino acids 1-33 of SEQ ID NO:21. A preferred Dkk protein of the present invention is a human protein (*e.g.*, encoded by a nucleotide sequence corresponding to a naturally-occurring human gene).

- 20 -

Accordingly, one embodiment of the invention features a Dkk protein having at least one cysteine-rich domain, preferably at least one cysteine-rich region. Another embodiment features a Dkk protein having at least one cysteine-rich region, wherein the cysteine-rich region includes at least one cysteine-rich domain. Another embodiment features a Dkk protein having at least one cysteine-rich region, wherein the cysteine-rich region includes at least two cysteine-rich domains. Another embodiment features a protein or domain within a protein having 20, 30, 40, 50, 60, 70, 80, 90, 95, or 99% homology to a cysteine-rich domain of a Dkk protein of the invention (*e.g.*, hDkk-3, hDkk-4, hDkk-1, or hDkk-2).

Yet another embodiment of the invention features a Dkk protein having at least one cysteine-rich domain, preferably at least one cysteine-rich region and a signal peptide. Another embodiment features a Dkk protein having at least one cysteine-rich domain, preferably at least one cysteine-rich region and a signal peptide, wherein the cysteine-rich region includes at least two cysteine-rich domains. Another embodiment features a Dkk protein having at least one cysteine-rich domain, preferably at least one cysteine-rich region and a signal peptide, wherein the cysteine-rich region includes at least two cysteine-rich domains and a spacer.

Yet another aspect of the invention features Dkk proteins having domains and/or regions which are conserved among a subset of Dkk proteins but are not necessarily conserved among all Dkk family members. In one embodiment, a Dkk protein (*e.g.*, Dkk-3) has an "extended N-terminal region" which is extended in length as compared to, for example, the "N-terminal regions" of other Dkk family members (*e.g.*, Dkk-4, Dkk-1, and Dkk-2). As defined herein, an "N-terminal region" of a Dkk proteins consists of amino acid residues found between the signal peptide and CRD-1 of a Dkk protein.

Preferably, the first amino acid residue of an N-terminal region of Dkk is the first residue of a mature Dkk protein and the last residue of an N-terminal region of Dkk is the residue preceeding the first cysteine residue of CRD-1. In a preferred embodiment, an N-terminal region is about 1-20 amino acid residues in length, preferably about 21-30, 31-40, 41-50, 51-60, 61-70, 71-80, 81-90, 91-100, 101-110, 111-120, 121-130, 131-140, 141-150, 151-160 or more amino acid residues in length. In contrast, an "extended N-terminal region" is at least about 71-80, 81-90, 91-100, 101-110, 111-120, 121-130,

131-140, 141-150, 151-160 or more amino acid residues in length. For example, in one embodiment, a hDkk-4 protein includes an "N-terminal region" of about amino acids 20-40 of SEQ ID NO:5 (21 amino acid residues in length). In another embodiment, a hDkk-1 protein includes an N-terminal region of about amino acids 21-84 of SEQ ID NO:8 (64 amino acid residues in length). In another embodiment, a hDkk-2 protein includes an "N-terminal region" of about amino acids 34-77 of SEQ ID NO:21 (44 amino acid residues in length). In another embodiment, a hDkk-3 protein has an "extended N-terminal region" of about amino acids 23-146 of SEQ ID NO:2 (124 amino acid residues in length).

10 In another embodiment, a Dkk protein (*e.g.*, Dkk-3) has an "acidic C-terminal region" which includes amino acid residues found C-terminal to CRD-2 of a Dkk protein. Preferably, the first amino acid residue of an acidic C-terminal region is the residue following the last cysteine of CRD-2 and the last residue of an acidic C-terminal region is the last residue of a Dkk protein. In a preferred embodiment, an acidic C-terminal region is about 65-66 amino acid residues in length and has about 27-25% acidic amino acid residues (*e.g.*, glutamic acid or aspartic acid). In another preferred embodiment, an acidic C-terminal region is about 55-80 amino acid residues in length, preferably about 60-75 amino acid residues in length, and more preferably about 64-70 amino acid residues in length and has about 21-35% acidic amino acid residues, preferably about 23-33% acidic amino acid residues, and more preferably about 25-31% acidic amino acid residues. Preferably, an acidic C-terminal region is involved in protein-protein interactions. For example, in one embodiment, a hDkk-3 protein has an acidic C-terminal region from about amino acids 285-350 of SEQ ID NO:2.

25 Preferred Dkk molecules of the present invention have an amino acid sequence sufficiently homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:21. As used herein, the term "sufficiently homologous" refers to a first amino acid or nucleotide sequence which contains a sufficient or minimum number of identical or equivalent (*e.g.*, an amino acid residue which has a similar side chain) amino acid residues or nucleotides to a second amino acid or nucleotide sequence such that the first and second amino acid or nucleotide sequences share common structural domains and/or a common functional activity. For example,

30

amino acid or nucleotide sequences which share common structural domains have at least about 40% homology, preferably 50% homology, more preferably 60%-70% homology across the amino acid sequences of the domains and contain at least one, preferably two, more preferably three, and even more preferably four, five or six

5 structural domains, are defined herein as sufficiently homologous. Furthermore, amino acid or nucleotide sequences which share at least 40%, preferably 50%, more preferably 60, 70, or 80% homology and share a common functional activity are defined herein as sufficiently homologous.

As used interchangeably herein, a "Dkk activity", "biological activity of Dkk" or
10 "functional activity of Dkk", refers to an activity exerted by a Dkk protein, polypeptide or nucleic acid molecule (*e.g.*, an activity on a Dkk responsive cell) as determined *in vivo*, or *in vitro*, according to standard techniques. In one embodiment, a Dkk activity is a direct activity, such as an association with a Dkk-target molecule. As used herein, a "target molecule" is a molecule with which a Dkk protein binds or interacts in nature,
15 such that Dkk-mediated function is achieved. A Dkk target molecule can be a non-Dkk molecule or a Dkk protein or polypeptide of the present invention. In an exemplary embodiment, a Dkk target molecule is a membrane-bound protein (*e.g.*, a cell-surface receptor or "Dkk receptor") or a modified form of such a protein which has been altered such that the protein is soluble (*e.g.*, recombinantly produced such that the protein does
20 not express a membrane-binding domain). In another embodiment, a Dkk target is a second soluble protein molecule (*e.g.*, a "Dkk binding partner" or "Dkk substrate"). In such an exemplary embodiment, a Dkk binding partner can be a second soluble non-Dkk protein or a second Dkk protein molecule of the present invention. Alternatively, a Dkk activity is an indirect activity, such as a cellular signaling activity mediated by
25 interaction of the Dkk protein with a second protein (*e.g.*, a Dkk receptor). As used herein, the term "Dkk receptor" refers to a protein or protein complex, to which a Dkk protein, *e.g.*, human Dkk, can bind. A receptor can be a cell surface receptor, *e.g.*, a peptide, growth factor, or nuclear hormone receptor. Dkk receptors can be isolated by methods known in the art and further described herein. Interaction of a Dkk protein with
30 a Dkk receptor can result in transduction of a signal from the cell surface to the nucleus. The signal transduced can be, an increase in intracellular calcium, an increase in

phosphatidylinositol or other molecule, and can result in, *e.g.*, in phosphorylation of specific proteins, a modulation of gene transcription and any of the other biological activities set forth herein.

In a preferred embodiment, a Dkk activity is at least one or more of the following activities: (i) interaction of a Dkk protein with and/or binding to a second molecule, (*e.g.*, a protein, such as a Dkk receptor, a soluble form of a Dkk receptor, a receptor for a member of the *wnt* family of signaling proteins, or a non-Dkk signaling molecule, for example, a lipid included in a cell membrane); (ii) interaction of a Dkk protein with an intracellular protein *via* a membrane-bound Dkk receptor; (iii) complex formation between a soluble Dkk protein and a second soluble Dkk binding partner (*e.g.*, a non-Dkk protein molecule or a second Dkk protein molecule); (iv) interaction with other extracellular proteins (*e.g.*, regulation of *wnt*-dependent cellular adhesion to extracellular matrix components); (v) binding to and eliminating an undesirable molecule (*e.g.*, a detoxifying activity or defense function); and/or (vi) an enzymatic activity. In yet another preferred embodiment, a Dkk activity is at least one or more of the following activities: (1) modulation of cellular signal transduction, either *in vitro* or *in vivo* (*e.g.*, modulation, *e.g.*, antagonism, of the activity of members of the *wnt* family of secreted proteins or suppression of *wnt*-dependent signal transduction, for example suppression of Wnt 2b, Wnt3 and/or Wnt8-dependent signal transduction by hDkk-1 and/or hDkk-4); (2) regulation of communication between cells (*e.g.*, regulation of *wnt*-dependent cell-cell interactions); (3) regulation of expression of genes whose expression is modulated by binding of Dkk (*e.g.*, hDkk-3) to a receptor; (4) regulation of gene transcription in a cell involved in development or differentiation, either *in vitro* or *in vivo* (*e.g.*, induction of cellular differentiation); (5) regulation of gene transcription in a cell involved in development or differentiation, wherein at least one gene encodes a differentiation-specific protein; (6) regulation of gene transcription in a cell involved in development or differentiation, wherein at least one gene encodes a second secreted protein; (7) regulation of gene transcription in a cell involved in development or differentiation, wherein at least one gene encodes a signal transduction molecule; (8) regulation of cellular proliferation, either *in vitro* or *in vivo* (*e.g.*, induction of cellular proliferation or inhibition of proliferation as in the case of suppression of tumorigenesis (*e.g.*, suppression

- 24 -

of glial cell tumor growth, for example, glioblastoma growth)); (9) formation and maintenance of ordered spatial arrangements of differentiated tissues in vertebrates, both adult and embryonic (*e.g.*, induction of head formation during vertebrate development or maintenance of hematopoietic progenitor cells); (10) modulation of cell death, such as stimulation of cell survival; (11) regulating cell migration; and/or (12) immune modulation.

As referred to herein, "differentiation-specific proteins" include proteins involved in the transition of a cell from the undifferentiated to the differentiated phenotype. For example, such proteins can be differentiation specific structural proteins or differentiation-specific transcription factors. Such differentiation-specific proteins are generally expressed at higher levels in cells which are making the transition from the undifferentiated to the differentiated phenotype (*e.g.*, during embryonic development or during regeneration of mature tissue in the adult animal), or are expressed at higher levels in fully-differentiated or terminally-differentiated cells as compared to their undifferentiated counterparts. Also, as referred to herein, "differentiation-specific genes" include nucleic acid molecules which encode differentiation-specific proteins.

Accordingly, another embodiment of the invention features isolated Dkk proteins and polypeptides having a Dkk activity. Preferred Dkk proteins have at least one cysteine-rich region and a Dkk activity. In another preferred embodiment, the Dkk protein has at least one cysteine-rich region, wherein the cysteine-rich region comprises at least one cysteine-rich domain, and a Dkk activity. In another preferred embodiment, the Dkk protein has at least one cysteine-rich region, wherein the cysteine-rich region comprises at least two cysteine-rich domains, and a Dkk activity. In yet another preferred embodiment, a Dkk protein further comprises a signal sequence. In still another preferred embodiment, a Dkk protein has a cysteine-rich region, a Dkk activity, and an amino acid sequence sufficiently homologous to an amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:21.

A preferred Dkk fragment comprises a carboxy-terminal cysteine-rich domain. In one embodiment, a Dkk fragment comprises a carboxy-terminal cysteine-rich domain and retains a biological activity of a Dkk protein. In yet another embodiment, a Dkk fragment lacks an amino-terminal cysteine-rich domain.

The human Dkk-3 cDNA, which is approximately 2479 nucleotides in length, encodes a protein which is approximately 350 amino acid residues in length. The human Dkk-3 protein contains an N-terminal signal sequence and a cysteine-rich region comprising two cysteine-rich domains. A Dkk cysteine-rich region can be found at least, for example, from about amino acids 147-284 of SEQ ID NO:2. The hDkk-3 cysteine-rich region comprises an amino-terminal cysteine-rich domain from about amino acids 147-195 of SEQ ID NO:2 and a carboxy-terminal cysteine-rich domain from about amino acids 201-284 of SEQ ID NO:2. The human Dkk-3 protein is a secreted protein which further contains a signal sequence at about amino acids 1-21, 1-22, 1-23, or 1-24 of SEQ ID NO:2. Accordingly, a mature human Dkk-3 protein begins at about amino acid residue 22, 23, 24, or 25 of SEQ ID NO:2 and is about 329, 328, 327, or 326 amino acids in length. The prediction of such a signal peptide can be made, for example, utilizing the computer algorithm SIGNALP (Nielsen, *et al.*, (1997) *Protein Engineering* 10:1-6).

The human Dkk-4 cDNA, which is approximately 848 nucleotides in length, encodes a protein which is approximately 224 amino acid residues in length. The human Dkk-4 protein contains an N-terminal signal sequence and a cysteine-rich region comprising two cysteine-rich domains. A Dkk cysteine-rich region can be found at least, for example, from about amino acids 41-218 of SEQ ID NO:5. The hDkk-4 cysteine-rich region comprises an amino-terminal cysteine-rich domain from about amino acids 41-90 of SEQ ID NO:5 and a carboxy-terminal cysteine-rich domain from about amino acids 138-218 of SEQ ID NO:5. The human Dkk-4 protein is a secreted protein which further contains a signal sequence at about amino acids 1-17, 1-18, 1-19, or 1-20 of SEQ ID NO:5. Accordingly, a mature human Dkk-4 protein begins at about amino acid residue 18, 19, 20, or 21 of SEQ ID NO:5 and is about 207, 206, 205, or 204 amino acids in length. A preferred fragment of hDkk-4 comprises amino acid residues 134-224 of SEQ ID NO:5. In another embodiment, a preferred fragment of hDkk-4 consists of amino acid residues 134-224 of SEQ ID NO:5.

The human Dkk-1 cDNA, which is approximately 1536 nucleotides in length, encodes a protein which is approximately 266 amino acid residues in length. The human Dkk-1 protein contains an N-terminal signal sequence and a cysteine-rich region

- 26 -

comprising two cysteine-rich domains. A Dkk cysteine-rich region can be found at least, for example, from about amino acids 85-263 of SEQ ID NO:8. The hDkk-1 cysteine-rich region comprises an amino-terminal cysteine-rich domain from about amino acids 85-138 of SEQ ID NO:8 and a carboxy-terminal cysteine-rich domain from about amino acids 182-263 of SEQ ID NO:8. The human Dkk-1 protein is a secreted protein which further contains a signal sequence at about amino acids 1-18, 1-19, 1-20, or 1-21 of SEQ ID NO:8. Accordingly, a mature human Dkk-1 protein begins at about amino acid residue 19, 20, 21, or 22 of SEQ ID NO:8 and is about 248, 247, 246, or 245 amino acids in length.

The human Dkk-2 cDNA, which is approximately 3687 nucleotides in length, encodes a protein which is approximately 259 amino acid residues in length. The human Dkk-2 protein contains a cysteine-rich region comprising two cysteine-rich domains. A Dkk cysteine-rich region can be found at least, for example, from about amino acids 78-256 of SEQ ID NO:21. The hDkk-2 cysteine-rich region comprises an amino-terminal cysteine-rich domain from about amino acids 78-127 of SEQ ID NO:21 and a carboxy-terminal cysteine-rich domain from about amino acids 176-256 of SEQ ID NO:21. The human Dkk-2 protein is a secreted protein which further contains a signal sequence at about amino acids 1-31, 1-32, 1-33, or 1-34 of SEQ ID NO:21. Accordingly, a mature human Dkk-2 protein begins at about amino acid residue 32, 33, 34, or 35 of SEQ ID NO:21 and is about 228, 227, 226, or 225 amino acids in length.

Dkk proteins of the present invention can be used to identify additional Dkk-related proteins or family members. For example, a protein having homology to hDkk-3 was identified using the nucleotide sequence encoding the N-terminal unique region of hDkk-3 to search a nucleotide sequence database. A human cDNA clone (Accession No.: AA397836) was identified from the dBEST database as having homology to hDkk-3 and was fully sequenced. The encoded protein is referred to herein as human "Soggy-1" or "Dkk-like-N". The nucleotide and predicted amino acid sequence of human Soggy-1 are depicted in Figure 7. The nucleotide sequence of human Soggy-1 (SEQ ID NO:13) encodes a protein having 242 amino acids (SEQ ID NO:14). The nucleotide sequence of human Soggy-1 includes a 5' untranslated region containing nucleotides 1-74 of SEQ ID NO:13, a coding region containing nucleotides 75-800 of SEQ ID NO:13

(corresponding to nucleotides 1-726 of SEQ ID NO:15), and a 3' untranslated region containing nucleotides 801-928 of SEQ ID NO:13. The Soggy-1 protein (amino acid residues 32-132) has 25% identity to an N-terminal domain of human Dkk-3 (consisting of amino acid residues 22-140) as determined by ALIGN, Myers and Miller, (1989)

5 *CABIOS*, utilizing a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4.

Two murine cDNA clones were further identified from the database and fully sequenced. Combining the sequence information from these two clones resulted in a full-length sequence for murine Soggy-1. The nucleotide sequence and predicted amino
10 acid sequence of murine Soggy-1 are depicted in Figure 8. The nucleotide sequence of murine Soggy-1 (SEQ ID NO:26) encodes a protein having 230 amino acids (SEQ ID NO:27). The nucleotide sequence of murine Soggy-1 includes a 5' untranslated region containing nucleotides 1-56 of SEQ ID NO:26, a coding region containing nucleotides 57-746 of SEQ ID NO:26 (corresponding to SEQ ID NO:26), and a 3' untranslated
15 region containing nucleotides 747-835 of SEQ ID NO:26. Human and murine Soggy-1 proteins display 59% overall identity. An alignment of human and murine Soggy proteins to human and murine Dkk-3 proteins is depicted in Figure 10.

In one embodiment, a Soggy protein is identified based on the presence of at least one soggy domain or "SGY" domain in the protein or corresponding nucleic acid
20 molecule. As defined herein, a "SGY domain" includes a protein domain of a Soggy protein (*e.g.*, hSoggy-1) having an amino acid sequence of about 45-56 amino acids and having at least about 25-40% identity with amino acid residues 90-140 of hDkk-3 (leu90-glu140 of SEQ ID NO:2). In another embodiment, a "SGY domain" has 46-55, preferably 47-54, more preferably about 48-53, and even more preferably about 49-52 or
25 50-51 amino acids, and has at least about 27-38%, preferably about 28-37%, more preferably about 29-36%, even more preferably about 30-35%, and even more preferably about 31-34%, or 32-33% identity with amino acid residues 90-140 of hDkk-3 (Leu90-Glu140 of SEQ ID NO:2). In yet another embodiment, a "SGY domain" has the following consensus sequence: L - P - X(3) - H - X - E - X(7) - G - N - X - T - X(3) - H -
30 X(4) - K - X - T - X - N - X(2) - G - X(4) - S - E - X - V - X(2) - S - X(4) - E (SEQ ID NO:29). For example, human Soggy-1 has a SGY domain from about amino acid

- 28 -

residues 81-131 (Leu 81-Glu131 of SEQ ID NO:14) having 33% identity with amino acid residues 90-140 of hDkk-3 (leu90-glu140 of SEQ ID NO:2). Likewise, murine Soggy-1 has a SGY domain from about amino acid residues 71-120 (Leu 71-Glu120 of SEQ ID NO:27) having 33% identity with amino acid residues 90-140 of hDkk-3 (leu90-glu140 of SEQ ID NO:2). The SGY domains of human and murine Soggy-1 are depicted by shaded boxes in Figure 10.

In another embodiment of the invention, a Soggy protein has at least one SGY domain and a signal sequence. For example, in one embodiment, a hSoggy-1 protein contains a signal sequence of about amino acids 1-29, 1-30, 1-31, or 1-32 of SEQ ID NO:14. Accordingly, a mature hSoggy-1 protein begins at about amino acid residue 30, 31, 32, or 33 of SEQ ID NO:14 and is about 213, 212, 211, or 210 amino acids in length. In another embodiment, a mSoggy-1 protein contains a signal sequence of about amino acids 1-19, 1-20, 1-21, or 1-22 of SEQ ID NO:27. Accordingly, a mature mSoggy-1 protein begins at about amino acid residue 211, 210, 209, or 208 of SEQ ID NO:28 and is about 213, 212, 211, or 210 amino acids in length.

Various aspects of the invention are described in further detail in the following subsections:

I. Isolated Nucleic Acid Molecules

One aspect of the invention pertains to isolated nucleic acid molecules that encode Dkk proteins or biologically active portions thereof, as well as nucleic acid fragments sufficient for use as hybridization probes to identify Dkk-encoding nucleic acids (*e.g.*, Dkk mRNA) and fragments for use as PCR primers for the amplification or mutation of Dkk nucleic acid molecules. As used herein, the term "nucleic acid molecule" is intended to include DNA molecules (*e.g.*, cDNA or genomic DNA) and RNA molecules (*e.g.*, mRNA) and analogs of the DNA or RNA generated using nucleotide analogs. The nucleic acid molecule can be single-stranded or double-stranded, but preferably is double-stranded DNA.

An "isolated" nucleic acid molecule is one which is separated from other nucleic acid molecules which are present in the natural source of the nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (*i.e.*,

- 29 -

sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated Dkk nucleic acid molecule can contain less than about 5 kb, 4kb, 3kb, 2kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived. An isolated chromosome is not an isolated nucleic acid molecule as defined herein. Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized.

A nucleic acid molecule of the present invention, *e.g.*, a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a portion thereof, can be isolated using standard molecular biology techniques and the sequence information provided herein. Using all or portion of the nucleic acid sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, as a hybridization probe, Dkk nucleic acid molecules can be isolated using standard hybridization and cloning techniques (*e.g.*, as described in Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989).

Moreover, a nucleic acid molecule encompassing all or a portion of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as

- 30 -

Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, can be isolated by the polymerase chain reaction (PCR) using synthetic oligonucleotide primers designed based upon the sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____.

A nucleic acid of the invention can be amplified using cDNA, mRNA or alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis. Furthermore, oligonucleotides corresponding to Dkk nucleotide sequences can be prepared by standard synthetic techniques, *e.g.*, using an automated DNA synthesizer.

In a preferred embodiment, an isolated nucleic acid molecule of the invention comprises the nucleotide sequence shown in SEQ ID NO:1. The sequence of SEQ ID NO:1 corresponds to the human Dkk-3 cDNA. This cDNA comprises sequences encoding the human Dkk-3 protein (*i.e.*, "the coding region", from nucleotides 38-1087), as well as 5' untranslated sequences (nucleotides 1 to 37) and 3' untranslated sequences (nucleotides 1088 to 2479). Alternatively, the nucleic acid molecule can comprise only the coding region of SEQ ID NO:1 (*e.g.*, nucleotides 38 to 1087, corresponding to SEQ ID NO:3). A plasmid containing the full-length nucleotide sequence encoding hDkk-3 was deposited with the American Type Culture Collection (ATCC), presently in Manassas Virginia, on June 11, 1997 and assigned Accession Number 98452.

In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises the nucleotide sequence shown in SEQ ID NO:4. The sequence of SEQ ID NO:4 corresponds to the human Dkk-4 cDNA. This cDNA comprises sequences encoding the human Dkk-4 protein (*i.e.*, "the coding region", from nucleotides 125-796), as well as 5' untranslated sequences (nucleotides 1 to 124) and 3'

- 31 -

untranslated sequences (nucleotides 797 to 848). Alternatively, the nucleic acid molecule can comprise only the coding region of SEQ ID NO:4 (*e.g.*, nucleotides 125 to 796, corresponding to SEQ ID NO:6).

5 In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises the nucleotide sequence shown in SEQ ID NO:7. The sequence of SEQ ID NO:7 corresponds to the human Dkk-1 cDNA. This cDNA comprises sequences encoding the human Dkk-1 protein (*i.e.*, "the coding region", from nucleotides 109-906), as well as 5' untranslated sequences (nucleotides 1 to 108) and 3' untranslated sequences (nucleotides 907-1536). Alternatively, the nucleic acid molecule
10 can comprise only the coding region of SEQ ID NO:7 (*e.g.*, nucleotides 109-906, corresponding to SEQ ID NO:9). A plasmid containing the full-length nucleotide sequence encoding hDkk-1 was deposited with the American Type Culture Collection (ATCC), presently in Manassas Virginia, on January 16, 1998 and assigned Accession Number 98633.

15 In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises the nucleotide sequence shown in SEQ ID NO:20. The sequence of SEQ ID NO:20 corresponds to the human Dkk-2 cDNA. This cDNA comprises sequences encoding the human Dkk-2 protein (*i.e.*, "the coding region", from nucleotides 724-1500), 5' untranslated sequences (nucleotides 1-723), as well as 3' untranslated sequences (nucleotides 1501-3687). Alternatively, the nucleic acid
20 molecule can comprise only the coding region of SEQ ID NO:20 (*e.g.*, nucleotides 724 to 1500, corresponding to SEQ ID NO:22). A plasmid, clone fthul33, containing the full-length nucleotide sequence encoding hDkk-2 was deposited with the American Type Culture Collection (ATCC), presently in Manassas Virginia, on March 2, 1999 and
25 assigned Accession Number _____.

In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises the nucleotide sequence shown in SEQ ID NO:13. The sequence of SEQ ID NO:13 corresponds to the human Soggy cDNA. This cDNA comprises sequences encoding the human Soggy protein (*i.e.*, "the coding region", from
30 nucleotides 75 to 800), as well as 5' untranslated sequences (nucleotides 1 to 74) and 3' untranslated sequences (nucleotides 801 to 928). Alternatively, the nucleic acid

- 32 -

molecule can comprise only the coding region of SEQ ID NO:13 (*e.g.*, nucleotides 75 to 800, corresponding to SEQ ID NO:15).

In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises a nucleic acid molecule which is a complement of the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a portion of any of these nucleotide sequences. A nucleic acid molecule which is complementary to the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, is one which is sufficiently complementary to the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, such that it can hybridize to the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, thereby forming a stable duplex.

In still another preferred embodiment, an isolated nucleic acid molecule of the present invention comprises a nucleotide sequence which is at least about 30-35%, preferably about 40-45%, more preferably about 50-55%, even more preferably about

- 33 -

60-65%, and even more preferably at least about 70-75%, 80-85%, 90-95% or more homologous to the nucleotide sequences shown in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:20, or SEQ ID NO:22, the nucleotide sequence of the DNA insert of the
5 plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a portion of any of these nucleotide sequences.

In one aspect, the present invention features isolated nucleic acid molecules
10 which are linear (*e.g.*, linear fragments of double-stranded DNA, linear strands of single-stranded DNA, single-stranded RNA molecules, and oligonucleotides). Another aspect of the present invention features circular nucleic acid molecules (*e.g.*, double-stranded DNA molecules, for example, plasmid molecules including the nucleotide sequences shown in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7,
15 SEQ ID NO:9, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:20, or SEQ ID NO:22, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a
20 portion of any of these nucleotide sequences).

In one embodiment, the isolated nucleic acid molecules of the present invention are DNA molecules which are in a form suitable for expression (*e.g.*, suitable for expression of corresponding messenger RNA or mRNA). In another embodiment, the isolated nucleic acid molecules are DNA molecules which are in a form suitable for
25 expression of corresponding protein (*e.g.*, in a form, for example, in a vector, which is capable of expressing protein, *e.g.*, in the appropriate orientation for expression from regulatory elements and/or in-frame with appropriate regulatory elements). In another embodiment, the isolated nucleic acids are in a form suitable for determination of nucleic acid sequence (*e.g.*, in a form suitable for sequencing, for example, is a
30 sequencing vector including a M13, T7, T3 and SP6 promoter. Examples of sequencing vectors include, but are not limited to pBluescript (Stratagene™), pT7T3D (Pharmacia™)

and pCR2.1 (InVitrogen). In yet another embodiment, the isolated nucleic acid molecules are free from vector sequences. In a preferred embodiment, an isolated nucleic acid molecule is free from sequencing vector sequences.

Moreover, the nucleic acid molecule of the invention can comprise only a portion
5 of the nucleic acid sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as
10 Accession Number _____, for example a fragment which can be used as a probe or primer or a fragment encoding a biologically active portion of a Dkk protein or Dkk-related protein. The nucleotide sequence determined from the cloning of the human Dkk genes allows for the generation of probes and primers designed for use in identifying and/or cloning Dkk homologues in other cell types, *e.g.*, from other tissues, as well as
15 Dkk homologues from other mammals and Dkk-related proteins. The probe/primer typically comprises substantially purified oligonucleotide. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, preferably about 25, more preferably about 40, 50 or 75 consecutive nucleotides of a sense sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ
20 ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, of an anti-sense sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA
25 insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or of a naturally occurring mutant of SEQ ID
30 NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number

- 35 -

98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____. In an exemplary embodiment, a nucleic acid molecule of the present invention comprises a nucleotide
5 sequence which hybridizes under stringent hybridization conditions to a nucleic acid molecule consisting of nucleotides 470-2479 of SEQ ID NO:1 or to a nucleic acid molecule consisting of nucleotides 1-475 of SEQ ID NO:4.

Probes based on human nucleotide sequences (*e.g.*, the human Dkk nucleotide sequence) can be used to detect transcripts or genomic sequences encoding the same or
10 homologous proteins. For instance, primers based on the nucleic acid represented in SEQ ID NOs:1 or 3 can be used in PCR reactions to clone Dkk homologs (*e.g.*, hDkk-3 homologues). In a preferred embodiment of the invention, Dkk homologs are cloned by PCR amplification (*e.g.*, RT-PCR) using primers hybridizing to a portion of the nucleotide sequence encoding the Dkk cysteine rich domain. Likewise, probes based on
15 the subject Dkk sequences can be used to detect transcripts or genomic sequences encoding the same or homologous proteins. In preferred embodiments, the probe further comprises a label group attached thereto, *e.g.*, the label group can be a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used as a part of a diagnostic test kit for identifying cells or tissue which misexpress a Dkk
20 protein, such as by measuring a level of a Dkk-encoding nucleic acid in a sample of cells from a subject *e.g.*, detecting Dkk mRNA levels or determining whether a genomic Dkk gene has been mutated or deleted.

A nucleic acid fragment encoding a "biologically active portion of a Dkk or Dkk-related protein" can be prepared by isolating a portion of SEQ ID NO:1, SEQ ID NO:4,
25 SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, which encodes a polypeptide having a biological
30 activity (the biological activities of the Dkk and Dkk-related proteins have previously

been described), expressing the encoded portion of the protein (*e.g.*, by recombinant expression *in vitro*) and assessing the activity of the encoded portion of the protein.

The invention further encompasses nucleic acid molecules that differ from the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, due to degeneracy of the genetic code and thus encode the same proteins as those encoded by the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____. In another embodiment, an isolated nucleic acid molecule of the invention has a nucleotide sequence encoding a protein having an amino acid sequence shown in SEQ ID NO:2, SEQ ID NO: 5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21.

In addition to the human nucleotide sequences shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, it will be appreciated by those skilled in the art that DNA sequence polymorphisms that lead to changes in the amino acid sequences of the Dkk or Dkk-related proteins may exist within a population (*e.g.*, the human population). Such genetic polymorphism in the Dkk or Dkk-related genes may exist among individuals within a population due to natural allelic variation. As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules comprising an open reading frame encoding a protein, preferably a mammalian Dkk or Dkk-related protein. Such natural allelic variations can typically result in 1-5% variance

in the nucleotide sequence of a Dkk or Dkk-related gene. Any and all such nucleotide variations and resulting amino acid polymorphisms in genes that are the result of natural allelic variation and that do not alter the functional activity of a Dkk or Dkk-related protein are intended to be within the scope of the invention.

5 Moreover, nucleic acid molecules encoding Dkk or Dkk-related proteins from other species, and thus which have a nucleotide sequence which differs from the human sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid
10 deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, are intended to be within the scope of the invention. For example, a murine Dkk-3 cDNA has been identified based of the nucleotide sequence of human Dkk-3. The nucleotide sequence of murine Dkk-3 (SEQ ID NO:16) encodes a hDkk-3 protein having 349
15 amino acids. The nucleotide and amino acid sequences of murine Dkk-3 are depicted in Figure 5. The coding region of murine Dkk-3 is represented by SEQ ID NO:18. A plasmid containing the full-length nucleotide sequence encoding mDkk-3 was deposited with the American Type Culture Collection (ATCC), presently in Manassas, Virginia, on January 16, 1998 and assigned Accession Number 98634. Likewise, a murine Dkk-
20 related protein (Soggy-1) has been identified based of the nucleotide sequence of human Dkk-3. The nucleotide sequence of murine Soggy-1 (SEQ ID NO:26) encodes a protein having 230 amino acids (SEQ ID NO:27). The nucleotide and amino acid sequences of murine Soggy-1 are depicted in Figure 8. The coding region of murine Soggy-1 is represented by SEQ ID NO:28.

25 Nucleic acid molecules corresponding to natural allelic variants and homologues of the Dkk or Dkk-related cDNAs of the invention can be isolated based on their homology to the human nucleic acids disclosed herein using the human cDNA, or a portion thereof, as a hybridization probe according to standard hybridization techniques under stringent hybridization conditions. Examples of tissues and/or libraries suitable
30 for isolation of the subject nucleic acids include brain, spinal chord and heart tissue. cDNA encoding a Dkk protein (*e.g.*, a hDkk-3 protein) can be obtained by isolating total

mRNA from a cell, *e.g.*, a vertebrate cell, a mammalian cell, or a human cell, including embryonic cells. Double stranded cDNAs can then be prepared from the total mRNA, and subsequently inserted into a suitable plasmid or bacteriophage vector using any one of a number of known techniques. The gene encoding a hDkk-3 protein can also be
5 cloned using established polymerase chain reaction techniques in accordance with the nucleotide sequence information provided by the invention. The nucleic acid of the invention can be DNA or RNA or analogs thereof.

Accordingly, in another embodiment, an isolated nucleic acid molecule of the invention is at least 15 nucleotides in length and hybridizes under stringent conditions to
10 the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid
15 deposited with ATCC as Accession Number _____. In another embodiment, the nucleic acid is at least 30, 50, 100, 250, 300, 400 or 500 nucleotides in length. As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences at least 60% homologous to each other typically remain hybridized to each other. Preferably, the conditions are
20 such that sequences at least about 70%, more preferably at least about 80%, even more preferably at least about 85% or 90% homologous to each other typically remain hybridized to each other. Such stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. A preferred, non-limiting example of stringent hybridization
25 conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65°C. Preferably, an isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to the sequence of SEQ ID NO:1 corresponds to a naturally-occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an
30 RNA or DNA molecule having a nucleotide sequence that occurs in nature (*e.g.*, encodes a natural protein).

In addition to naturally-occurring allelic variants of the Dkk or Dkk-related sequences that may exist in the population, the skilled artisan will further appreciate that changes can be introduced by mutation into the nucleotide sequences of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, thereby leading to changes in the amino acid sequence of the encoded Dkk proteins, without altering the functional ability of the Dkk proteins. For example, nucleotide substitutions leading to amino acid substitutions (particularly conservative amino acid substitutions) at "non-essential" amino acid residues can be made in the sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____. A "non-essential" amino acid residue is a residue that can be altered from the wild-type sequence of Dkk (or wild-type Dkk-related sequence) (*e.g.*, the sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21) without altering the biological activity, whereas an "essential" amino acid residue is required for biological activity. For example, amino acid residues that are conserved among the Dkk or Dkk-related proteins of the present invention (*e.g.*, cysteine residues within cysteine-rich domains), are predicted to be particularly unamenable to alteration. Furthermore, amino acid residues that are conserved between Dkk protein and other proteins having cysteine-rich domains are not likely to be amenable to alteration.

Accordingly, another aspect of the invention pertains to nucleic acid molecules encoding Dkk or Dkk-related proteins that contain changes in amino acid residues that are not essential for activity. Such proteins differ in amino acid sequence from SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21 yet retain biological activity. In one embodiment, the isolated nucleic acid molecule comprises a

- 40 -

nucleotide sequence encoding a protein, wherein the protein comprises an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21. Preferably, the protein encoded by the nucleic acid molecule is at least about 65-70% homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21, more preferably at least about 75-80% homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21, even more preferably at least about 85-90% homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21, and most preferably at least about 95% homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21.

An isolated nucleic acid molecule encoding a Dkk protein homologous to the protein of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21 can be created by introducing one or more nucleotide substitutions, additions or deletions into the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, such that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein. Mutations can be introduced into SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Preferably, conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (*e.g.*,

lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*,
5 threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine). Thus, for example, a predicted nonessential amino acid residue in a Dkk protein (*e.g.*, one not located in a cysteine-rich domain) is preferably replaced with another amino acid residue from the same side chain family. Alternatively, in another embodiment, mutations can be introduced randomly along all or part of a Dkk or
10 Dkk-related coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for biological activity to identify mutants that retain activity. Following mutagenesis of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of
15 the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, the encoded protein can be expressed recombinantly and the activity of the protein can be determined.

In a preferred embodiment, a mutant Dkk or Dkk-related protein can be assayed
20 for intracellular calcium, an increase in phosphatidylinositol or other molecule, and can result, *e.g.*, in phosphorylation of specific proteins, a modulation of gene transcription and any of the other biological activities set forth herein.

In a preferred embodiment, a mutant Dkk or Dkk-related protein can also be assayed for the ability to (1) modulate cellular signal transduction, either *in vitro* or *in*
25 *vivo*; (2) regulate communication between cells; (3) regulate expression of genes whose expression is modulated by binding of Dkk (*e.g.*, hDkk-3) to a receptor; (4) regulate gene transcription in a cell involved in development or differentiation, either *in vitro* or *in vivo*; (5) regulate cellular proliferation, either *in vitro* or *in vivo*; (6) form and/or maintain ordered spatial arrangements of differentiated tissues in vertebrates; (7)
30 modulate cell death (*e.g.* cell survival); (8) regulate cell migration; and/or (9) modulate immune system function.

In addition to the nucleic acid molecules encoding Dkk or Dkk-related proteins described above, another aspect of the invention pertains to isolated nucleic acid molecules which are antisense thereto. An "antisense" nucleic acid comprises a nucleotide sequence which is complementary to a "sense" nucleic acid encoding a protein, *e.g.*, complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence. Accordingly, an antisense nucleic acid can hydrogen bond to a sense nucleic acid. The antisense nucleic acid can be complementary to an entire Dkk coding strand, or to only a portion thereof. In one embodiment, an antisense nucleic acid molecule is antisense to a "coding region" of the coding strand of a nucleotide sequence encoding Dkk. The term "coding region" refers to the region of the nucleotide sequence comprising codons which are translated into amino acid residues (*e.g.*, the coding region of human Dkk-3 corresponds to SEQ ID NO:3, the coding region of human Dkk-4 corresponds to SEQ ID NO:6, the coding region of human Dkk-1 corresponds to SEQ ID NO:9, the coding region of human Dkk-2 corresponds to SEQ ID NO:22, and the coding region of human Soggy corresponds to SEQ ID NO:15). In another embodiment, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence encoding a Dkk or Dkk-related protein. The term "noncoding region" refers to 5' and 3' sequences which flank the coding region that are not translated into amino acids (*i.e.*, also referred to as 5' and 3' untranslated regions).

Given the coding strand sequences disclosed herein (*e.g.*, SEQ ID NO:3, SEQ ID NO:6, SEQ ID NO:9, SEQ ID NO:15, or SEQ ID NO:22), antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick base pairing. The antisense nucleic acid molecule can be complementary to the entire coding region of a Dkk or Dkk-related mRNA, but more preferably is an oligonucleotide which is antisense to only a portion of the coding or noncoding region of the mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of Dkk mRNA. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, or 90 nucleotides in length. In a preferred embodiment, an oligonucleotide is about 30-90, preferably about 40-80, more preferably about 50-70 nucleotides in length and is antisense to a portion of

SEQ ID NO:1 from about nucleotides 1-150. In another embodiment, an oligonucleotide is antisense to a portion of SEQ ID NO:4 from about nucleotides 25-225. In another embodiment, an oligonucleotide is antisense to a portion of SEQ ID NO:7 from about nucleotides 1-200. In another embodiment, an oligonucleotide is antisense to a portion of SEQ ID NO:20 from about nucleotides 625-825. In yet another embodiment, an oligonucleotide is antisense to a portion of SEQ ID NO:13 from about nucleotides 1-175.

An antisense nucleic acid of the invention can be constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (*e.g.*, an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, *e.g.*, phosphorothioate derivatives, acridine substituted nucleotides, can be used. Alternatively, the antisense nucleic acid molecule can be synthesized to increase transport across cellular membranes, *e.g.*, methylphosphonate derivatives. The antisense molecules can include a 3'-terminal cap (*e.g.*, a 3'-aminopropyl modification), a biotin moiety, or even a 3'-3' terminal linkage.

Examples of modified nucleotides which can be used to generate the antisense nucleic acid include 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the

antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (*i.e.*, RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

5 The antisense nucleic acid molecules of the invention are typically administered to a subject or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a Dkk or Dkk-related protein to thereby inhibit expression of the protein, *e.g.*, by inhibiting transcription and/or translation. The hybridization can be by conventional nucleotide complementarity to form a stable
10 duplex, or, for example, in the case of an antisense nucleic acid molecule which binds to DNA duplexes, through specific interactions in the major groove of the double helix. An example of a route of administration of antisense nucleic acid molecules of the invention include direct injection at a tissue site. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then administered systemically.
15 For example, for systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface, *e.g.*, by linking the antisense nucleic acid molecules to peptides or antibodies which bind to cell surface receptors or antigens. The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular
20 concentrations of the antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

 In yet another embodiment, the antisense nucleic acid molecule of the invention is an α -anomeric nucleic acid molecule. An α -anomeric nucleic acid molecule forms
25 specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gaultier *et al.*, (1987) *Nucleic Acids Res.* 15:6625-6641). The antisense nucleic acid molecule can also comprise a 2'-O-methylribonucleotide (Inoue *et al.*, (1987) *Nucleic Acids Res.* 15:6131-6148) or a chimeric RNA-DNA analogue (Inoue *et al.*, (1987) *FEBS Lett.* 215:327-330).

In still another embodiment, an antisense nucleic acid of the invention is a ribozyme. Ribozymes are catalytic RNA molecules with ribonuclease activity which are capable of cleaving a single-stranded nucleic acid, such as an mRNA, to which they have a complementary region. Thus, ribozymes (*e.g.*, hammerhead ribozymes
5 (described in Haselhoff and Gerlach (1988) *Nature* 334:585-591)) can be used to catalytically cleave Dkk or Dkk-related mRNA transcripts to thereby inhibit translation of Dkk or Dkk-related mRNA. A ribozyme having specificity for a Dkk- or Dkk-related-encoding nucleic acid can be designed based upon the nucleotide sequence of a Dkk or Dkk-related cDNA disclosed herein (*i.e.*, SEQ ID NO:1, SEQ ID NO:4, SEQ ID
10 NO:7, SEQ ID NO:13, SEQ ID NO:20, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633). For example, a derivative of a *Tetrahymena* L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be
15 cleaved in a Dkk-encoding mRNA. See, *e.g.*, Cech *et al.*, U.S. Patent No. 4,987,071; and Cech *et al.*, U.S. Patent No. 5,116,742. Alternatively, Dkk (or Dkk-related) mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See, *e.g.*, Bartel, D. and Szostak, J.W. (1993) *Science* 261:1411-1418.

20 Alternatively, gene expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region of the Dkk or Dkk-related gene (*e.g.*, the promoter and/or enhancers) to form triple helical structures that prevent transcription of the gene in target cells. See generally, Helene, C. (1991) *Anticancer Drug Des.* 6(6):569-84; Helene, C. *et al.*, (1992) *Ann. N.Y. Acad. Sci.* 660:27-36; and Maher, L.J.
25 (1992) *Bioassays* 14(12):807-15.

In yet another embodiment, the nucleic acid molecules of the present invention can be modified at the base moiety, sugar moiety or phosphate backbone to improve, *e.g.*, the stability, hybridization, or solubility of the molecule. For example, the deoxyribose phosphate backbone of the nucleic acid molecules can be modified to
30 generate peptide nucleic acids (see Hyrup B. *et al.* (1996) *Bioorganic & Medicinal Chemistry* 4 (1): 5-23). As used herein, the terms "peptide nucleic acids" or "PNAs"

refer to nucleic acid mimics, *e.g.*, DNA mimics, in which the deoxyribose phosphate backbone is replaced by a pseudopeptide backbone and only the four natural nucleobases are retained. The neutral backbone of PNAs has been shown to allow for specific hybridization to DNA and RNA under conditions of low ionic strength. The synthesis of PNA oligomers can be performed using standard solid phase peptide synthesis protocols as described in Hyrup B. *et al.* (1996) *supra*; Perry-O'Keefe *et al.* PNAS 93: 14670-675.

PNAs of Dkk or Dkk-related nucleic acid molecules can be used therapeutic and diagnostic applications. For example, PNAs can be used as antisense or antigene agents for sequence-specific modulation of gene expression by, for example, inducing transcription or translation arrest or inhibiting replication. PNAs of Dkk or Dkk-related nucleic acid molecules can also be used in the analysis of single base pair mutations in a gene, (*e.g.*, by PNA-directed PCR clamping); as 'artificial restriction enzymes' when used in combination with other enzymes, (*e.g.*, S1 nucleases (Hyrup B. (1996) *supra*)); or as probes or primers for DNA sequencing or hybridization (Hyrup B. *et al.* (1996) *supra*; Perry-O'Keefe *supra*).

In another embodiment, PNAs of Dkk can be modified, (*e.g.*, to enhance their stability or cellular uptake), by attaching lipophilic or other helper groups to PNA, by the formation of PNA-DNA chimeras, or by the use of liposomes or other techniques of drug delivery known in the art. For example, PNA-DNA chimeras of Dkk nucleic acid molecules can be generated which may combine the advantageous properties of PNA and DNA. Such chimeras allow DNA recognition enzymes, (*e.g.*, RNase H and DNA polymerases), to interact with the DNA portion while the PNA portion would provide high binding affinity and specificity. PNA-DNA chimeras can be linked using linkers of appropriate lengths selected in terms of base stacking, number of bonds between the nucleobases, and orientation (Hyrup B. (1996) *supra*). The synthesis of PNA-DNA chimeras can be performed as described in Hyrup B. (1996) *supra* and Finn P.J. *et al.* (1996) *Nucleic Acids Res.* 24 (17): 3357-63. For example, a DNA chain can be synthesized on a solid support using standard phosphoramidite coupling chemistry and modified nucleoside analogs, *e.g.*, 5'-(4-methoxytrityl)amino-5'-deoxy-thymidine phosphoramidite, can be used as a between the PNA and the 5' end of DNA (Mag, M. *et*

al. (1989) *Nucleic Acid Res.* 17: 5973-88). PNA monomers are then coupled in a stepwise manner to produce a chimeric molecule with a 5' PNA segment and a 3' DNA segment (Finn P.J. *et al.* (1996) *supra*). Alternatively, chimeric molecules can be synthesized with a 5' DNA segment and a 3' PNA segment (Peterser, K.H. *et al.* (1975) 5 *Bioorganic Med. Chem. Lett.* 5: 1119-11124).

In other embodiments, the oligonucleotide may include other appended groups such as peptides (*e.g.*, for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (see, *e.g.*, Letsinger *et al.* (1989) *Proc. Natl. Acad. Sci. US.* 86:6553-6556; Lemaitre *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:648-652; 10 PCT Publication No. W088/09810, published December 15, 1988) or the blood-brain barrier (see, *e.g.*, PCT Publication No. W089/10134, published April 25, 1988). In addition, oligonucleotides can be modified with hybridization-triggered cleavage agents (See, *e.g.*, Krol *et al.* (1988) *BioTechniques* 6:958-976) or intercalating agents. (See, *e.g.*, Zon (1988) *Pharm. Res.* 5:539-549). To this end, the oligonucleotide may be 15 conjugated to another molecule, (*e.g.*, a peptide, hybridization triggered cross-linking agent, transport agent, or hybridization-triggered cleavage agent).

II. Isolated Dkk Proteins and Anti-Dkk Antibodies

One aspect of the invention pertains to isolated Dkk proteins, Dkk-related 20 proteins and biologically active portions thereof, as well as polypeptide fragments suitable for use as immunogens to raise antibodies. In one embodiment, native Dkk or Dkk-related proteins can be isolated from cells or tissue sources by an appropriate purification scheme using standard protein purification techniques. In another embodiment, proteins are produced by recombinant DNA techniques. Alternative to 25 recombinant expression, a Dkk or Dkk-related protein or polypeptide can be synthesized chemically using standard peptide synthesis techniques.

An "isolated" or "purified" protein or biologically active portion thereof is substantially free of cellular material or other contaminating proteins from the cell or tissue source from which the Dkk or Dkk-related protein is derived, or substantially free 30 from chemical precursors or other chemicals when chemically synthesized. The language "substantially free of cellular material" includes preparations of protein in

which the protein is separated from cellular components of the cells from which it is isolated or recombinantly produced. In one embodiment, the language "substantially free of cellular material" includes preparations of Dkk or Dkk-related protein having less than about 30% (by dry weight) of non-Dkk protein or non-Dkk-related protein (also referred to herein as a "contaminating protein"), more preferably less than about 20% of non-Dkk protein or non-Dkk-related protein, still more preferably less than about 10% of non-Dkk protein or non-Dkk-related protein, and most preferably less than about 5% non-Dkk protein or non-Dkk-related protein. When the Dkk or Dkk-related protein or biologically active portion thereof is recombinantly produced, it is also preferably substantially free of culture medium, *i.e.*, culture medium represents less than about 20%, more preferably less than about 10%, and most preferably less than about 5% of the volume of the protein preparation.

The language "substantially free of chemical precursors or other chemicals" includes preparations of Dkk or Dkk-related protein in which the protein is separated from chemical precursors or other chemicals which are involved in the synthesis of the protein. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of Dkk protein having less than about 30% (by dry weight) of chemical precursors, non-Dkk chemicals, or non-Dkk-related chemicals, more preferably less than about 20% chemical precursors, non-Dkk chemicals, or non-Dkk-related chemicals, still more preferably less than about 10% chemical precursors, non-Dkk chemicals, or non-Dkk-related chemicals, and most preferably less than about 5% chemical precursors, non-Dkk chemicals, or non-Dkk-related chemicals.

Biologically active portions of a Dkk or Dkk-related protein include peptides comprising amino acid sequences sufficiently homologous to or derived from the amino acid sequence of the Dkk or Dkk-related protein, *e.g.*, the amino acid sequence shown in SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21, which include less amino acids than the full length proteins, and exhibit at least one activity of a Dkk or Dkk-related protein. Typically, biologically active portions comprise a domain or motif with at least one activity of the Dkk or Dkk-related protein. A biologically active portion of a protein can be a polypeptide which is, for example, 10, 25, 50, 100 or more amino acids in length.

In one embodiment, a biologically active portion of a Dkk protein comprises at least a cysteine-rich region. In another embodiment, a biologically active portion of a Dkk protein comprises at least a cysteine-rich region, wherein the cysteine-rich region includes at least one cysteine-rich domain. In yet another embodiment, a biologically active portion of a Dkk protein comprises at least a signal sequence.

In another embodiment, a biologically active portion of a Dkk-related protein (*e.g.*, a Soggy protein) comprises at least a Soggy domain. In yet another embodiment, a biologically active portion of a Dkk-related protein comprises at least a signal sequence.

In an alternative embodiment, a biologically active portion of a Dkk or Dkk-related protein comprises an amino acid sequence lacking a signal sequence.

It is to be understood that a preferred biologically active portion of a Dkk or Dkk-related protein of the present invention may contain at least one of the above-identified structural domains. A more preferred biologically active portion of a Dkk or Dkk-related protein may contain at least two of the above-identified structural domains.

An even more preferred biologically active portion of a protein may contain at least three of the above-identified structural domains. A particularly preferred biologically active portion of a protein of the present invention may contain at least four of the above-identified structural domains.

Moreover, other biologically active portions, in which other regions of the protein are deleted, can be prepared by recombinant techniques and evaluated for one or more of the functional activities of a native Dkk or Dkk-related protein.

In a preferred embodiment, the Dkk protein has an amino acid sequence shown in SEQ ID NO:2 or an amino acid sequence at least about 55% homologous to SEQ ID NO:2. In another preferred embodiment, the Dkk protein has an amino acid sequence shown in SEQ ID NO:5 or an amino acid sequence at least about 35% homologous to SEQ ID NO:5. In another preferred embodiment, the Dkk protein has an amino acid sequence shown in SEQ ID NO:8 or an amino acid sequence at least about 85% homologous to SEQ ID NO:8. In another preferred embodiment, the Dkk protein has an amino acid sequence shown in SEQ ID NO:21 or an amino acid sequence at least about 35% homologous to SEQ ID NO:21. In another preferred embodiment, the protein has an amino acid sequence shown in SEQ ID NO:14 or an amino acid sequence at least

about 60% homologous to SEQ ID NO:14. In still another preferred embodiment, a protein of the present invention comprises an amino acid sequence which is at least about 30-35%, preferably about 40-45%, more preferably about 50-55%, even more preferably about 60-65%, and even more preferably at least about 70-75%, 80-85%, 90-
5 95% or more homologous to the amino acid sequences shown in SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21.

In other embodiments, the protein is substantially homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21, and, preferably, retains the functional activity of the protein of SEQ ID NO:2, SEQ ID NO:5, SEQ ID
10 NO:8, SEQ ID NO:14, or SEQ ID NO:21, yet differs in amino acid sequence due to natural allelic variation or mutagenesis, as described in detail in subsection I above.

Accordingly, in another embodiment, the protein is a protein which comprises an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21 and, preferably,
15 retains the functional activity of the proteins of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:11, respectively. Preferably, the protein is at least about 70% homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21, more preferably at least about 80% homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21, even more
20 preferably at least about 90% homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21, and most preferably at least about 95% or more homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21.

To determine the percent homology of two amino acid sequences or of two
25 nucleic acids, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence and non-homologous sequences can be disregarded for comparison purposes). In one embodiment, an alignment is a global alignment, *e.g.*, an overall sequence alignment. In another embodiment, an
30 alignment is a local alignment. In a preferred embodiment, the length of a sequence aligned for comparison purposes is at least 30%, preferably at least 40%, more

preferably at least 50%, even more preferably at least 60%, and even more preferably at least 70%, 80%, or 90% of the length of the reference sequence to which it is aligned (*e.g.*, when aligning a second sequence to the Dkk amino acid sequence of SEQ ID NO:2, at least 105, preferably at least 145, more preferably at least 175, even more preferably at least 210, and even more preferably at least 245, 280 or 315 amino acid residues are aligned). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid or nucleic acid "identity" is equivalent to amino acid or nucleic acid "homology"). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. In an embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch (*J. Mol. Biol.* (48):444-453 (1970)) algorithm which has been incorporated into the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, 5, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (available at <http://www.gcg.com>), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. In another embodiment, the percent identity between two amino acid or nucleotide sequences is determined using the algorithm of E. Meyers and W. Miller (*CABIOS*, 4:11-17 (1989)) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

The nucleic acid and protein sequences of the present invention can further be used as a "query sequence" to perform a search against public databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, *et al.* (1990) *J.*

5 *Mol. Biol.* 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to Dkk nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to Dkk protein molecules of the invention. To obtain
10 gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, (1997) *Nucleic Acids Res.* 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (*e.g.*, XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>.

The invention also provides Dkk or Dkk-related chimeric or fusion proteins. As
15 used herein, a "chimeric protein" or "fusion protein" comprises a Dkk or Dkk-related polypeptide operatively linked to a non-Dkk polypeptide or non-Dkk-related polypeptide. A "Dkk polypeptide" or "Dkk-related polypeptide" refers to a polypeptide having an amino acid sequence corresponding to Dkk or a Dkk-related protein, whereas a "non-Dkk polypeptide" or "non-Dkk-related polypeptide" refers to a polypeptide
20 having an amino acid sequence corresponding to a protein which is not substantially homologous to the Dkk or Dkk-related protein, *e.g.*, a protein which is different from the Dkk or Dkk-related protein and which is derived from the same or a different organism. Within a Dkk or Dkk-related fusion protein the Dkk or Dkk-related polypeptide can correspond to all or a portion of a Dkk or Dkk-related protein. In a preferred
25 embodiment, a Dkk or Dkk-related fusion protein comprises at least one biologically active portion of a Dkk protein. In another preferred embodiment, a Dkk or Dkk-related fusion protein comprises at least two biologically active portions of a Dkk or Dkk-related protein. In another preferred embodiment, a Dkk or Dkk-related fusion protein comprises at least three biologically active portions of a Dkk or Dkk-related protein.
30 Within the fusion protein, the term "operatively linked" is intended to indicate that the Dkk or Dkk-related polypeptide and the non-Dkk or non-Dkk-related polypeptide are

fused in-frame to each other. The non-Dkk or non-Dkk-related polypeptide can be fused to the N-terminus or C-terminus of the Dkk or Dkk-related polypeptide.

For example, in one embodiment, the fusion protein is a GST-Dkk fusion protein in which the Dkk sequences are fused to the C-terminus of the GST sequences. Such fusion proteins can facilitate the purification of recombinant Dkk.

In another embodiment, the fusion protein is a Dkk or Dkk-related protein containing a heterologous signal sequence at its N-terminus. For example, the native Dkk signal sequence (i.e., about amino acids 1 to 23 of SEQ ID NO:2) can be removed and replaced with a signal sequence from another protein. In certain host cells (*e.g.*, mammalian host cells), expression and/or secretion of Dkk or Dkk-related proteins can be increased through use of a heterologous signal sequence.

In yet another embodiment, the fusion protein is a Dkk-immunoglobulin fusion protein in which the Dkk sequences comprising primarily the Dkk cysteine-rich regions are fused to sequences derived from a member of the immunoglobulin protein family.

Soluble derivatives have also been made of cell surface glycoproteins in the immunoglobulin gene superfamily consisting of an extracellular domain of the cell surface glycoprotein fused to an immunoglobulin constant (Fc) region (see *e.g.*, Capon, *et al.* (1989) *Nature* 337:525-531 and Capon U.S. Patents 5,116,964 and 5,428,130 [CD4-IgG1 constructs]; Linsley, P.S. *et al.* (1991) *J. Exp. Med.* 173:721-730 [a CD28-IgG1 construct and a B7-1-IgG1 construct]; and Linsley, P.S. *et al.* (1991) *J. Exp. Med.* 174:561-569 and U.S. Patent 5,434,131[a CTLA4-IgG1]). Such fusion proteins have proven useful for modulating receptor-ligand interactions. Soluble derivatives of cell surface proteins of the tumor necrosis factor receptor (TNFR) superfamily proteins have been made consisting of an extracellular domain of the cell surface receptor fused to an immunoglobulin constant (Fc) region (See for example Moreland *et al.* (1997) *N. Engl. J. Med.* 337(3):141-147; van der Poll *et al.* (1997) *Blood* 89(10):3727-3734; and Ammann *et al.* (1997) *J. Clin. Invest.* 99(7):1699-1703.)

The Dkk-immunoglobulin fusion proteins of the invention can be incorporated into pharmaceutical compositions and administered to a subject to inhibit an interaction between a Dkk ligand and a Dkk receptor on the surface of a cell, to thereby suppress Dkk-mediated signal transduction *in vivo*. The Dkk-immunoglobulin fusion proteins

can be used to affect the bioavailability of a Dkk cognate receptor. Inhibition of the Dkk ligand/Dkk interaction may be useful therapeutically for both the treatment of differentiative or proliferative disorders, as well as modulating (*e.g.*, promoting or inhibiting) developmental responses, cell adhesion, and/or cell fate. Moreover, the Dkk-immunoglobulin fusion proteins of the invention can be used as immunogens to produce anti-Dkk antibodies in a subject, to purify Dkk ligands and in screening assays to identify molecules which inhibit the interaction of Dkk with a Dkk ligand.

Preferably, a Dkk or Dkk-related chimeric or fusion protein of the invention is produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques, for example by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and reamplified to generate a chimeric gene sequence (see, for example, *Current Protocols in Molecular Biology*, eds. Ausubel *et al.*, John Wiley & Sons: 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (*e.g.*, a GST polypeptide). A Dkk-encoding nucleic acid or nucleic acid encoding a Dkk-related protein can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the protein.

The present invention also pertains to variants of the Dkk or Dkk-related proteins which function as either agonists (mimetics) or as antagonists. Variants of the Dkk or Dkk-related proteins can be generated by mutagenesis, *e.g.*, discrete point mutation or truncation of a Dkk or Dkk-related protein. An agonist of the Dkk or Dkk-related proteins can retain substantially the same, or a subset, of the biological activities of the naturally occurring form of a Dkk or Dkk-related protein. An antagonist of a Dkk or Dkk-related protein can inhibit one or more of the activities of the naturally occurring

- 55 -

form of the protein by, for example, competitively binding to a downstream or upstream member of a cellular signaling cascade which includes the Dkk or Dkk-related protein. Thus, specific biological effects can be elicited by treatment with a variant of limited function. In one embodiment, treatment of a subject with a variant having a subset of the biological activities of the naturally occurring form of the protein has fewer side effects in a subject relative to treatment with the naturally occurring form of the Dkk or Dkk-related protein.

In one embodiment, variants of a Dkk or Dkk-related protein which function as either agonists (mimetics) or as antagonists can be identified by screening combinatorial libraries of mutants, *e.g.*, truncation mutants, of a Dkk or Dkk-related protein for protein agonist or antagonist activity. In one embodiment, a variegated library of Dkk or Dkk-related variants is generated by combinatorial mutagenesis at the nucleic acid level and is encoded by a variegated gene library. A variegated library of Dkk or Dkk-related variants can be produced by, for example, enzymatically ligating a mixture of synthetic oligonucleotides into gene sequences such that a degenerate set of potential Dkk or Dkk-related sequences is expressible as individual polypeptides, or alternatively, as a set of larger fusion proteins (*e.g.*, for phage display) containing the set of Dkk or Dkk-related sequences therein. There are a variety of methods which can be used to produce libraries of potential Dkk or Dkk-related variants from a degenerate oligonucleotide sequence. Chemical synthesis of a degenerate gene sequence can be performed in an automatic DNA synthesizer, and the synthetic gene then ligated into an appropriate expression vector. Use of a degenerate set of genes allows for the provision, in one mixture, of all of the sequences encoding the desired set of potential Dkk or Dkk-related sequences. Methods for synthesizing degenerate oligonucleotides are known in the art (see, *e.g.*, Narang, S.A. (1983) *Tetrahedron* 39:3; Itakura *et al.*, (1984) *Annu. Rev. Biochem.* 53:323; Itakura *et al.*, (1984) *Science* 198:1056; Ike *et al.*, (1983) *Nucleic Acid Res.* 11:477).

In addition, libraries of fragments of a Dkk or Dkk-related protein coding sequence can be used to generate a variegated population of Dkk or Dkk-related fragments for screening and subsequent selection of variants of a Dkk or Dkk-related protein. In one embodiment, a library of coding sequence fragments can be generated by

treating a double stranded PCR fragment of a Dkk coding sequence with a nuclease under conditions wherein nicking occurs only about once per molecule, denaturing the double stranded DNA, renaturing the DNA to form double stranded DNA which can include sense/antisense pairs from different nicked products, removing single stranded
5 portions from reformed duplexes by treatment with S1 nuclease, and ligating the resulting fragment library into an expression vector. By this method, an expression library can be derived which encodes N-terminal, C-terminal and internal fragments of various sizes of the Dkk protein.

Several techniques are known in the art for screening gene products of
10 combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene products having a selected property. Such techniques are adaptable for rapid screening of the gene libraries generated by the combinatorial mutagenesis of Dkk or Dkk-related proteins. The most widely used techniques, which are amenable to high through-put analysis, for screening large gene libraries typically include cloning the
15 gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and expressing the combinatorial genes under conditions in which detection of a desired activity facilitates isolation of the vector encoding the gene whose product was detected. Recursive ensemble mutagenesis (REM), a new technique which enhances the frequency of functional mutants in the libraries, can be used in
20 combination with the screening assays to identify Dkk variants (Arkin and Yourvan (1992) *PNAS* 89:7811-7815; Delgrave *et al.*, (1993) *Protein Engineering* 6(3):327-331).

In one embodiment, cell based assays can be exploited to analyze a variegated Dkk or Dkk-related library. For example, a library of expression vectors can be transfected into a cell line which ordinarily responds to a particular ligand in a Dkk-
25 dependent manner. The transfected cells are then contacted with the ligand and the effect of expression of the mutant on signaling by the ligand can be detected, *e.g.*, by measuring any of a number of immune cell responses. Plasmid DNA can then be recovered from the cells which score for inhibition, or alternatively, potentiation of ligand induction, and the individual clones further characterized.

An isolated Dkk protein, Dkk-related protein, or a portion or fragment thereof, can be used as an immunogen to generate antibodies that bind Dkk or Dkk-related proteins using standard techniques for polyclonal and monoclonal antibody preparation. A full-length Dkk or Dkk-related protein can be used or, alternatively, the invention
5 provides antigenic peptide fragments for use as immunogens. The antigenic peptide of Dkk comprises at least 8 amino acid residues of the amino acid sequence shown in SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21 and encompasses an epitope of Dkk or Dkk-related protein such that an antibody raised against the peptide forms a specific immune complex with the protein. Preferably, the
10 antigenic peptide comprises at least 10 amino acid residues, more preferably at least 15 amino acid residues, even more preferably at least 20 amino acid residues, and most preferably at least 30 amino acid residues.

Preferred epitopes encompassed by the antigenic peptide are regions of Dkk or Dkk-related proteins that are located on the surface of the protein, *e.g.*, hydrophilic
15 regions.

A Dkk or Dkk-related immunogen typically is used to prepare antibodies by immunizing a suitable subject, (*e.g.*, rabbit, goat, mouse or other mammal) with the immunogen. An appropriate immunogenic preparation can contain, for example, recombinantly expressed Dkk or Dkk-related protein or a chemically synthesized Dkk or
20 Dkk-related polypeptide. The preparation can further include an adjuvant, such as Freund's complete or incomplete adjuvant, or similar immunostimulatory agent. Immunization of a suitable subject with an immunogenic Dkk preparation, for example, induces a polyclonal anti-Dkk antibody response.

Accordingly, another aspect of the invention pertains to anti-Dkk antibodies as
25 well as antibodies to Dkk-related proteins. The term "antibody" as used herein refers to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, *i.e.*, molecules that contain an antigen binding site which specifically binds (immunoreacts with) an antigen, such as Dkk or Dkk-related antigens. Examples of immunologically active portions of immunoglobulin molecules include F(ab) and
30 F(ab')₂ fragments which can be generated by treating the antibody with an enzyme such as pepsin. The invention provides polyclonal and monoclonal antibodies that bind Dkk

or Dkk-related polypeptides. The term "monoclonal antibody" or "monoclonal antibody composition", as used herein, refers to a population of antibody molecules that contain only one species of an antigen binding site capable of immunoreacting with a particular epitope of Dkk or a or Dkk-related protein. A monoclonal antibody composition thus typically displays a single binding affinity for a particular Dkk or Dkk-related protein with which it immunoreacts.

Polyclonal antibodies can be prepared as described above by immunizing a suitable subject with a Dkk or Dkk-related immunogen. The antibody titer in the immunized subject can be monitored over time by standard techniques, such as with an enzyme linked immunosorbent assay (ELISA) using immobilized Dkk or Dkk-related protein. If desired, the antibody molecules directed against Dkk or Dkk-related protein can be isolated from the mammal (*e.g.*, from the blood) and further purified by well known techniques, such as protein A chromatography to obtain the IgG fraction. At an appropriate time after immunization, *e.g.*, when the antibody titers are highest, antibody-producing cells can be obtained from the subject and used to prepare monoclonal antibodies by standard techniques, such as the hybridoma technique originally described by Kohler and Milstein (1975) *Nature* 256:495-497) (see also, Brown *et al.*, (1981) *J. Immunol.* 127:539-46; Brown *et al.*, (1980) *J. Biol. Chem.* 255:4980-83; Yeh *et al.*, (1976) *PNAS* 76:2927-31; and Yeh *et al.*, (1982) *Int. J. Cancer* 29:269-75), the more recent human B cell hybridoma technique (Kozbor *et al.*, (1983) *Immunol Today* 4:72), the EBV-hybridoma technique (Cole *et al.*, (1985), *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96) or trioma techniques. The technology for producing monoclonal antibody hybridomas is well known (see generally R. H. Kenneth, in *Monoclonal Antibodies: A New Dimension In Biological Analyses*, Plenum Publishing Corp., New York, New York (1980); E. A. Lerner (1981) *Yale J. Biol. Med.*, 54:387-402; M.L. Gefter *et al.*, (1977) *Somatic Cell Genet.* 3:231-36). Briefly, an immortal cell line (typically a myeloma) is fused to lymphocytes (typically splenocytes) from a mammal immunized with a Dkk or Dkk-related immunogen as described above, and the culture supernatants of the resulting hybridoma cells are screened to identify a hybridoma producing a monoclonal antibody that binds Dkk or Dkk-related protein.

Any of the many well known protocols used for fusing lymphocytes and immortalized cell lines can be applied for the purpose of generating a monoclonal antibody (see, *e.g.*, G. Galfre *et al.*, (1977) *Nature* 266:55052; Gefter *et al.*, *Somatic Cell Genet.*, cited *supra*; Lerner, *Yale J. Biol. Med.*, cited *supra*; Kenneth, *Monoclonal Antibodies*, cited *supra*). Moreover, the ordinarily skilled worker will appreciate that there are many variations of such methods which also would be useful. Typically, the immortal cell line (*e.g.*, a myeloma cell line) is derived from the same mammalian species as the lymphocytes. For example, murine hybridomas can be made by fusing lymphocytes from a mouse immunized with an immunogenic preparation of the present invention with an immortalized mouse cell line. Preferred immortal cell lines are mouse myeloma cell lines that are sensitive to culture medium containing hypoxanthine, aminopterin and thymidine ("HAT medium"). Any of a number of myeloma cell lines can be used as a fusion partner according to standard techniques, *e.g.*, the P3-NS1/1-Ag4-1, P3-x63-Ag8.653 or Sp2/O-Ag14 myeloma lines. These myeloma lines are available from ATCC. Typically, HAT-sensitive mouse myeloma cells are fused to mouse splenocytes using polyethylene glycol ("PEG"). Hybridoma cells resulting from the fusion are then selected using HAT medium, which kills unfused and unproductively fused myeloma cells (unfused splenocytes die after several days because they are not transformed). Hybridoma cells producing a monoclonal antibody of the invention are detected by screening the hybridoma culture supernatants for antibodies that bind Dkk or Dkk-related protein, *e.g.*, using a standard ELISA assay.

Alternative to preparing monoclonal antibody-secreting hybridomas, a monoclonal antibody can be identified and isolated by screening a recombinant combinatorial immunoglobulin library (*e.g.*, an antibody phage display library) with Dkk or Dkk-related protein to thereby isolate immunoglobulin library members that bind Dkk or Dkk-related protein. Kits for generating and screening phage display libraries are commercially available (*e.g.*, the Pharmacia *Recombinant Phage Antibody System*, Catalog No. 27-9400-01; and the Stratagene *SurfZAP™ Phage Display Kit*, Catalog No. 240612). Additionally, examples of methods and reagents particularly amenable for use in generating and screening antibody display library can be found in, for example, Ladner *et al.*, U.S. Patent No. 5,223,409; Kang *et al.*, PCT International Publication No.

WO 92/18619; Dower *et al.*, PCT International Publication No. WO 91/17271; Winter *et al.*, PCT International Publication WO 92/20791; Markland *et al.*, PCT International Publication No. WO 92/15679; Breitling *et al.*, PCT International Publication WO 93/01288; McCafferty *et al.*, PCT International Publication No. WO 92/01047; Garrard *et al.*, PCT International Publication No. WO 92/09690; Ladner *et al.*, PCT International Publication No. WO 90/02809; Fuchs *et al.*, (1991) *Bio/Technology* 9:1370-1372; Hay *et al.*, (1992) *Hum. Antibod. Hybridomas* 3:81-85; Huse *et al.*, (1989) *Science* 246:1275-1281; Griffiths *et al.*, (1993) *EMBO J* 12:725-734; Hawkins *et al.*, (1992) *J. Mol. Biol.* 226:889-896; Clarkson *et al.*, (1991) *Nature* 352:624-628; Gram *et al.*, (1992) *PNAS* 89:3576-3580; Garrard *et al.*, (1991) *Bio/Technology* 9:1373-1377; Hoogenboom *et al.*, (1991) *Nuc. Acid Res.* 19:4133-4137; Barbas *et al.*, (1991) *PNAS* 88:7978-7982; and McCafferty *et al.*, *Nature* (1990) 348:552-554.

Additionally, recombinant antibodies, such as chimeric and humanized monoclonal antibodies, comprising both human and non-human portions, which can be made using standard recombinant DNA techniques, are within the scope of the invention. Such chimeric and humanized monoclonal antibodies can be produced by recombinant DNA techniques known in the art, for example using methods described in Robinson *et al.*, International Application No. PCT/US86/02269; Akira, *et al.*, European Patent Application 184,187; Taniguchi, M., European Patent Application 171,496; Morrison *et al.*, European Patent Application 173,494; Neuberger *et al.*, PCT International Publication No. WO 86/01533; Cabilly *et al.*, U.S. Patent No. 4,816,567; Cabilly *et al.*, European Patent Application 125,023; Better *et al.*, (1988) *Science* 240:1041-1043; Liu *et al.*, (1987) *PNAS* 84:3439-3443; Liu *et al.*, (1987) *J. Immunol.* 139:3521-3526; Sun *et al.*, (1987) *PNAS* 84:214-218; Nishimura *et al.*, (1987) *Canc. Res.* 47:999-1005; Wood *et al.*, (1985) *Nature* 314:446-449; and Shaw *et al.*, (1988) *J. Natl. Cancer Inst.* 80:1553-1559; Morrison, S. L. (1985) *Science* 229:1202-1207; Oi *et al.*, (1986) *BioTechniques* 4:214; Winter U.S. Patent 5,225,539; Jones *et al.*, (1986) *Nature* 321:552-525; Verhoeven *et al.*, (1988) *Science* 239:1534; and Beidler *et al.*, (1988) *J. Immunol.* 141:4053-4060.

An antibody (*e.g.*, monoclonal antibody) can be used to isolate Dkk or Dkk-related protein by standard techniques, such as affinity chromatography or immunoprecipitation. An antibody can facilitate the purification of natural Dkk or Dkk-related protein from cells and of recombinantly produced Dkk or Dkk-related protein expressed in host cells. Moreover, an antibody can be used to detect Dkk or Dkk-related protein (*e.g.*, in a cellular lysate or cell supernatant) in order to evaluate the abundance and pattern of expression of the Dkk or Dkk-related protein. Antibodies can be used diagnostically to monitor protein levels in tissue as part of a clinical testing procedure, *e.g.*, to, for example, determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling (*i.e.*, physically linking) the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

III. Recombinant Expression Vectors and Host Cells

Another aspect of the invention pertains to vectors, preferably expression vectors, containing a nucleic acid encoding Dkk or a nucleic acid encoding a Dkk-related protein (or a portion thereof). As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (*e.g.*, bacterial vectors having a bacterial origin of replication and episomal

mammalian vectors). Other vectors (*e.g.*, non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "expression vectors". In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" can be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (*e.g.*, replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

The recombinant expression vectors of the invention comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, which is operatively linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, "operably linked" is intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner which allows for expression of the nucleotide sequence (*e.g.*, in an *in vitro* transcription/translation system or in a host cell when the vector is introduced into the host cell). The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (*e.g.*, polyadenylation signals). Such regulatory sequences are described, for example, in Goeddel; *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Regulatory sequences include those which direct constitutive expression of a nucleotide sequence in many types of host cell and those which direct expression of the nucleotide sequence only in certain host cells (*e.g.*, tissue-specific regulatory sequences). It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. The expression vectors of the invention can be introduced into host cells to thereby produce proteins or peptides, including fusion proteins or peptides, encoded by nucleic acids as

- 63 -

described herein (*e.g.*, Dkk proteins, Dkk-related proteins, mutant forms of Dkk or Dkk-related proteins, fusion proteins, etc.).

The recombinant expression vectors of the invention can be designed for expression of Dkk or Dkk-related proteins in prokaryotic or eukaryotic cells. For example, Dkk can be expressed in bacterial cells such as *E. coli*, insect cells (using baculovirus expression vectors) yeast cells or mammalian cells. Suitable host cells are discussed further in Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Alternatively, the recombinant expression vector can be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

Expression of proteins in prokaryotes is most often carried out in *E. coli* with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein encoded therein, usually to the amino terminus of the recombinant protein. Such fusion vectors typically serve three purposes: 1) to increase expression of recombinant protein; 2) to increase the solubility of the recombinant protein; and 3) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase. Typical fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith, D.B. and Johnson, K.S. (1988) *Gene* 67:31-40), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein.

Purified fusion proteins can be utilized in activity assays, in ligand binding (*e.g.*, direct assays or competitive assays described in detail below), to generate antibodies specific for Dkk or Dkk-related proteins, as examples. In a preferred embodiment, a Dkk or Dkk-related fusion expressed in a retroviral expression vector of the present invention can be utilized to infect bone marrow cells which are subsequently

transplanted into irradiated recipients. The pathology of the subject recipient is then examined after sufficient time has passed (e.g six (6) weeks).

Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amann *et al.*, (1988) *Gene* 69:301-315) and pET 11d (Studier *et al.*, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 60-89). Target gene expression from the pTrc vector relies on host RNA polymerase transcription from a hybrid trp-lac fusion promoter. Target gene expression from the pET 11d vector relies on transcription from a T7 gn10-lac fusion promoter mediated by a coexpressed viral RNA polymerase (T7 gn1). This viral polymerase is supplied by host strains BL21(DE3) or HMS174(DE3) from a resident λ prophage harboring a T7 gn1 gene under the transcriptional control of the lacUV 5 promoter.

One strategy to maximize recombinant protein expression in *E. coli* is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant protein (Gottesman, S., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 119-128). Another strategy is to alter the nucleic acid sequence of the nucleic acid to be inserted into an expression vector so that the individual codons for each amino acid are those preferentially utilized in *E. coli* (Wada *et al.*, (1992) *Nucleic Acids Res.* 20:2111-2118). Such alteration of nucleic acid sequences of the invention can be carried out by standard DNA synthesis techniques.

In another embodiment, the Dkk or Dkk-related expression vector is a yeast expression vector. Examples of vectors for expression in yeast *S. cerevisiae* include pYepSec1 (Baldari, *et al.*, (1987) *Embo J.* 6:229-234), pMFa (Kurjan and Herskowitz, (1982) *Cell* 30:933-943), pJRY88 (Schultz *et al.*, (1987) *Gene* 54:113-123), pYES2 (Invitrogen Corporation, San Diego, CA), and picZ (InVitrogen Corp, San Diego, CA).

Alternatively, Dkk or Dkk-related protein can be expressed in insect cells using baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., Sf 9 cells) include the pAc series (Smith *et al.*, (1983) *Mol. Cell Biol.* 3:2156-2165) and the pVL series (Lucklow and Summers (1989) *Virology* 170:31-39).

In yet another embodiment, a nucleic acid of the invention is expressed in mammalian cells using a mammalian expression vector. Examples of mammalian expression vectors include pCDM8 (Seed, B. (1987) *Nature* 329:840) and pMT2PC (Kaufman *et al.*, (1987) *EMBO J.* 6:187-195). When used in mammalian cells, the expression vector's control functions are often provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma, Adenovirus 2, cytomegalovirus and Simian Virus 40. For other suitable expression systems for both prokaryotic and eukaryotic cells see chapters 16 and 17 of Sambrook, J., Fritsch, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual. 2nd, ed.*, Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.

In another embodiment, the recombinant mammalian expression vector is capable of directing expression of the nucleic acid preferentially in a particular cell type (*e.g.*, tissue-specific regulatory elements are used to express the nucleic acid). Tissue-specific regulatory elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert *et al.*, (1987) *Genes Dev.* 1:268-277), lymphoid-specific promoters (Calame and Eaton (1988) *Adv. Immunol.* 43:235-275), in particular promoters of T cell receptors (Winoto and Baltimore (1989) *EMBO J.* 8:729-733) and immunoglobulins (Banerji *et al.*, (1983) *Cell* 33:729-740; Queen and Baltimore (1983) *Cell* 33:741-748), neuron-specific promoters (*e.g.*, the neurofilament promoter; Byrne and Ruddle (1989) *PNAS* 86:5473-5477), pancreas-specific promoters (Edlund *et al.*, (1985) *Science* 230:912-916), and mammary gland-specific promoters (*e.g.*, milk whey promoter; U.S. Patent No. 4,873,316 and European Application Publication No. 264,166). Developmentally-regulated promoters are also encompassed, for example the murine hox promoters (Kessel and Gruss (1990) *Science* 249:374-379) and the α -fetoprotein promoter (Campes and Tilghman (1989) *Genes Dev.* 3:537-546).

The invention further provides a recombinant expression vector comprising a DNA molecule of the invention cloned into the expression vector in an antisense orientation. That is, the DNA molecule is operatively linked to a regulatory sequence in a manner which allows for expression (by transcription of the DNA molecule) of an

- 66 -

RNA molecule which is antisense to Dkk mRNA or a Dkk-related mRNA. Regulatory sequences operatively linked to a nucleic acid cloned in the antisense orientation can be chosen which direct the continuous expression of the antisense RNA molecule in a variety of cell types, for instance viral promoters and/or enhancers, or regulatory sequences can be chosen which direct constitutive, tissue specific or cell type specific expression of antisense RNA. The antisense expression vector can be in the form of a recombinant plasmid, phagemid or attenuated virus in which antisense nucleic acids are produced under the control of a high efficiency regulatory region, the activity of which can be determined by the cell type into which the vector is introduced. For a discussion of the regulation of gene expression using antisense genes see Weintraub, H. *et al.*, Antisense RNA as a molecular tool for genetic analysis, *Reviews - Trends in Genetics*, Vol. 1(1) 1986.

Another aspect of the invention pertains to host cells into which a recombinant expression vector of the invention has been introduced. The terms "host cell" and "recombinant host cell" are used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

A host cell can be any prokaryotic or eukaryotic cell. For example, Dkk protein can be expressed in bacterial cells such as *E. coli*, insect cells, yeast or mammalian cells (such as Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to those skilled in the art.

Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (*e.g.*, DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation. Suitable methods for transforming or transfecting host cells can be found in Sambrook, *et al.*, (*Molecular Cloning: A*

Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989), and other laboratory manuals.

For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may
5 integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (*e.g.*, resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Preferred selectable markers include those which confer resistance to drugs, such as G418, hygromycin and methotrexate. Nucleic acid encoding a selectable marker can be
10 introduced into a host cell on the same vector as that encoding Dkk or can be introduced on a separate vector. Cells stably transfected with the introduced nucleic acid can be identified by drug selection (*e.g.*, cells that have incorporated the selectable marker gene will survive, while the other cells die).

A host cell of the invention, such as a prokaryotic or eukaryotic host cell in
15 culture, can be used to produce (*i.e.*, express) a Dkk or Dkk-related protein. Accordingly, the invention further provides methods for producing Dkk or Dkk-related proteins using the host cells of the invention. In one embodiment, the method comprises culturing the host cell of invention (into which a recombinant expression vector encoding Dkk or a Dkk-related protein has been introduced) in a suitable medium such
20 that protein is produced. In another embodiment, the method further comprises isolating Dkk or a Dkk-related protein from the medium or the host cell.

The host cells of the invention can also be used to produce nonhuman transgenic animals. For example, in one embodiment, a host cell of the invention is a fertilized oocyte or an embryonic stem cell into which Dkk-coding sequences (or Dkk-related
25 coding sequences) have been introduced. Such host cells can then be used to create non-human transgenic animals in which exogenous Dkk sequences (or Dkk-related sequences) have been introduced into their genome or homologous recombinant animals in which endogenous Dkk sequences (or Dkk-related sequences) have been altered. Such animals are useful for studying the function and/or activity of Dkk or Dkk-related
30 proteins and for identifying and/or evaluating modulators of Dkk or Dkk-related protein activity. As used herein, a "transgenic animal" is a non-human animal, preferably a

- 68 -

mammal, more preferably a rodent such as a rat or mouse, in which one or more of the cells of the animal includes a transgene. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, amphibians, etc. A transgene is exogenous DNA which is integrated into the genome of a cell from which a transgenic animal develops and which remains in the genome of the mature animal, thereby directing the expression of an encoded gene product in one or more cell types or tissues of the transgenic animal. As used herein, a "homologous recombinant animal" is a non-human animal, preferably a mammal, more preferably a mouse, in which an endogenous Dkk or Dkk-related gene has been altered by homologous recombination between the endogenous gene and an exogenous DNA molecule introduced into a cell of the animal, *e.g.*, an embryonic cell of the animal, prior to development of the animal.

A transgenic animal of the invention can be created, for example, by introducing Dkk-encoding nucleic acid into the male pronuclei of a fertilized oocyte, *e.g.*, by microinjection, retroviral infection, and allowing the oocyte to develop in a pseudopregnant female foster animal. The human Dkk cDNA sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, or SEQ ID NO:20 can be introduced as a transgene into the genome of a non-human animal. Alternatively, a nonhuman homologue of a human Dkk gene, such as a mouse Dkk gene, can be isolated based on hybridization to the human Dkk cDNA (described further in subsection I above) and used as a transgene. Intronic sequences and polyadenylation signals can also be included in the transgene to increase the efficiency of expression of the transgene. A tissue-specific regulatory sequence(s) can be operably linked to the Dkk transgene to direct expression of Dkk protein to particular cells. Methods for generating transgenic animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009, both by Leder *et al.*, U.S. Patent No. 4,873,191 by Wagner *et al.*, and in Hogan, B., *Manipulating the Mouse Embryo*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986). Similar methods are used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of the Dkk transgene in its genome and/or expression of Dkk mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional

animals carrying the transgene. Moreover, transgenic animals carrying a transgene encoding Dkk can further be bred to other transgenic animals carrying other transgenes.

To create a homologous recombinant animal, a vector is prepared which contains at least a portion of a Dkk gene into which a deletion, addition or substitution has been introduced to thereby alter, *e.g.*, functionally disrupt, the Dkk gene. The Dkk gene can be a human gene (*e.g.*, the cDNA of SEQ ID NO:3, SEQ ID NO: 6, SEQ ID NO:9 or SEQ ID NO:22), but more preferably, is a non-human homologue of a human Dkk gene. For example, a mouse Dkk gene of SEQ ID NO:16 can be used to construct a homologous recombination vector suitable for altering an endogenous Dkk gene in the mouse genome. In a preferred embodiment, the vector is designed such that, upon homologous recombination, the endogenous Dkk gene is functionally disrupted (*i.e.*, no longer encodes a functional protein; also referred to as a "knock out" vector). Alternatively, the vector can be designed such that, upon homologous recombination, the endogenous Dkk gene is mutated or otherwise altered but still encodes functional protein (*e.g.*, the upstream regulatory region can be altered to thereby alter the expression of the endogenous Dkk protein). In the homologous recombination vector, the altered portion of the Dkk gene is flanked at its 5' and 3' ends by additional nucleic acid of the Dkk gene to allow for homologous recombination to occur between the exogenous Dkk gene carried by the vector and an endogenous Dkk gene in an embryonic stem cell. The additional flanking Dkk nucleic acid is of sufficient length for successful homologous recombination with the endogenous gene. Typically, several kilobases of flanking DNA (both at the 5' and 3' ends) are included in the vector (see *e.g.*, Thomas, K.R. and Capecchi, M. R. (1987) *Cell* 51:503 for a description of homologous recombination vectors). The vector is introduced into an embryonic stem cell line (*e.g.*, by electroporation) and cells in which the introduced Dkk gene has homologously recombined with the endogenous Dkk gene are selected (see *e.g.*, Li, E. *et al.*, (1992) *Cell* 69:915). The selected cells are then injected into a blastocyst of an animal (*e.g.*, a mouse) to form aggregation chimeras (see *e.g.*, Bradley, A. in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E.J. Robertson, ed. (IRL, Oxford, 1987) pp. 113-152). A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term. Progeny harboring the

homologously recombined DNA in their germ cells can be used to breed animals in which all cells of the animal contain the homologously recombined DNA by germline transmission of the transgene. Methods for constructing homologous recombination vectors and homologous recombinant animals are described further in Bradley, A.

5 (1991) *Current Opinion in Biotechnology* 2:823-829 and in PCT International Publication Nos.: WO 90/11354 by Le Mouellec *et al.*; WO 91/01140 by Smithies *et al.*; WO 92/0968 by Zijlstra *et al.*; and WO 93/04169 by Berns *et al.* It is also within the scope of the present invention to practice the above-described transgenic methodology utilizing nucleic acid molecules which encode Dkk-related proteins.

10 In another embodiment, transgenic non-humans animals can be produced which contain selected systems which allow for regulated expression of the transgene. One example of such a system is the *cre/loxP* recombinase system of bacteriophage P1. For a description of the *cre/loxP* recombinase system, see, *e.g.*, Lakso *et al.*, (1992) *PNAS* 89:6232-6236. Another example of a recombinase system is the FLP recombinase
15 system of *Saccharomyces cerevisiae* (O'Gorman *et al.*, (1991) *Science* 251:1351-1355. If a *cre/loxP* recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the *Cre* recombinase and a selected protein are required. Such animals can be provided through the construction of "double" transgenic animals, *e.g.*, by mating two transgenic animals, one containing a transgene encoding a
20 selected protein and the other containing a transgene encoding a recombinase.

Clones of the non-human transgenic animals described herein can also be produced according to the methods described in Wilmot, I. *et al.*, (1997) *Nature* 385:810-813. In brief, a cell, *e.g.*, a somatic cell, from the transgenic animal can be isolated and induced to exit the growth cycle and enter G₀ phase. The quiescent cell can
25 then be fused, *e.g.*, through the use of electrical pulses, to an enucleated oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops to morula or blastocyte and then transferred to pseudopregnant female foster animal. The offspring borne of this female foster animal will be a clone of the animal from which the cell, *e.g.*, the somatic cell, is
30 isolated.

IV. Pharmaceutical Compositions

The Dkk and Dkk-related nucleic acid molecules, Dkk and Dkk-related proteins, and anti-Dkk or anti-Dkk-related protein antibodies (also referred to herein as "active compounds") of the invention can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier. As used herein the language "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, *e.g.*, intravenous, intradermal, subcutaneous, oral (*e.g.*, inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic

water, Cremophor EL™ (BASF, Parsippany, NJ) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound (*e.g.*, a Dkk protein, Dkk-related protein or antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is

- 73 -

applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as

5 microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

10 For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant, *e.g.*, a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be
15 permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in
20 the art.

The compounds can also be prepared in the form of suppositories (*e.g.*, with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

In one embodiment, the active compounds are prepared with carriers that will
25 protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art.
30 The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected

cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD50/ED50. Compounds which exhibit large therapeutic indices are preferred. While compounds that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC50 (*i.e.*, the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine

useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

The nucleic acid molecules of the invention can be inserted into vectors and used as gene therapy vectors. Gene therapy vectors can be delivered to a subject by, for example, intravenous injection, local administration (see U.S. Patent 5,328,470) or by stereotactic injection (see *e.g.*, Chen *et al.*, (1994) *PNAS* 91:3054-3057). The pharmaceutical preparation of the gene therapy vector can include the gene therapy vector in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from recombinant cells, *e.g.*, retroviral vectors, the pharmaceutical preparation can include one or more cells which produce the gene delivery system.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

V. Uses and Methods of the Invention

The molecules of the present invention (*e.g.*, nucleic acid molecules, proteins, protein homologues, and antibodies described herein) can be used in one or more of the following methods: a) screening assays; b) predictive medicine (*e.g.*, diagnostic assays, prognostic assays, monitoring clinical trials, and pharmacogenetics); and c) methods of treatment (*e.g.*, therapeutic and prophylactic). As described herein, a Dkk protein of the invention has one or more of the following activities: intracellular calcium, an increase in phosphatidylinositol or other molecule, and can result, *e.g.*, in phosphorylation of specific proteins, a modulation of gene transcription and any of the other biological activities set forth herein.

In a preferred embodiment, a Dkk activity is at least one or more of the following activities: (i) interaction of a Dkk protein with and/or binding to a second molecule, (*e.g.*, a protein, such as a Dkk (*e.g.*, hDkk-3) receptor, a soluble form of a Dkk receptor, a receptor for a member of the *wnt* family of signaling proteins, or a non-Dkk signaling molecule); (ii) interaction of a Dkk protein with an intracellular protein via a membrane-bound Dkk receptor; (iii) complex formation between a soluble Dkk protein and a

- 76 -

second soluble Dkk binding partner (*e.g.*, a non-Dkk protein molecule or a second Dkk protein molecule); (iv) interaction with other extracellular proteins (*e.g.*, regulation of *wnt*-dependent cellular adhesion to extracellular matrix components); (v) binding to and eliminating an undesirable molecule (*e.g.*, a detoxifying activity or defense function);

5 and/or (vi) an enzymatic activity, and can thus be used in, for example, (1) modulation of cellular signal transduction, either *in vitro* or *in vivo* (*e.g.*, antagonism of the activity of members of the *wnt* family of secreted proteins or suppression of *wnt*-dependent signal transduction); (2) regulation of communication between cells (*e.g.*, regulation of *wnt*-dependent cell-cell interactions); (3) regulation of expression of genes

10 whose expression is modulated by binding of Dkk (*e.g.*, hDkk-3) to a receptor; (4) regulation of gene transcription in a cell involved in development or differentiation, either *in vitro* or *in vivo* (*e.g.*, induction of cellular differentiation); (5) regulation of gene transcription in a cell involved in development or differentiation, wherein at least one gene encodes a differentiation-specific protein; (6) regulation of gene transcription

15 in a cell involved in development or differentiation, wherein at least one gene encodes a second secreted protein; (7) regulation of gene transcription in a cell involved in development or differentiation, wherein at least one gene encodes a signal transduction molecule; (8) regulation of cellular proliferation, either *in vitro* or *in vivo* (*e.g.*, induction of cellular proliferation or inhibition of proliferation, for example, inhibition of

20 tumorigenesis (*e.g.*, inhibition of glioblastoma proliferation)); (9) formation and maintenance of ordered spatial arrangements of differentiated tissues in vertebrates, both adult and embryonic (*e.g.*, induction of head formation during vertebrate development or maintenance of hematopoietic progenitor cells); (10) modulation of cell death, such as stimulation of cell survival; (11) regulating cell migration; and/or (12) immune

25 modulation.

Accordingly one embodiment of the present invention involves a method of use (*e.g.*, a diagnostic assay, prognostic assay, or a prophylactic/therapeutic method of treatment) wherein a molecule of the present invention (*e.g.*, a Dkk protein, Dkk nucleic acid, or a Dkk modulator) is used, for example, to diagnose, prognose and/or treat a

30 disease and/or condition in which any of the aforementioned activities (*i.e.*, activities (i) - (vi) and (1) - (12) in the above paragraph) is indicated. In another embodiment, the

- 77 -

present invention involves a method of use (*e.g.*, a diagnostic assay, prognostic assay, or a prophylactic/therapeutic method of treatment) wherein a molecule of the present invention (*e.g.*, a Dkk protein, Dkk nucleic acid, or a Dkk modulator) is used, for example, for the diagnosis, prognosis, and/or treatment of subjects, preferably a human subject, in which any of the aforementioned activities is pathologically perturbed. In a preferred embodiment, the methods of use (*e.g.*, diagnostic assays, prognostic assays, or prophylactic/therapeutic methods of treatment) involve administering to a subject, preferably a human subject, a molecule of the present invention (*e.g.*, a Dkk protein, Dkk nucleic acid, or a Dkk modulator) for the diagnosis, prognosis, and/or therapeutic treatment. In another embodiment, the methods of use (*e.g.*, diagnostic assays, prognostic assays, or prophylactic/therapeutic methods of treatment) involve administering to a human subject a molecule of the present invention (*e.g.*, a Dkk protein, Dkk nucleic acid, or a Dkk modulator).

Other embodiments of the invention pertain to the use of isolated nucleic acid molecules of the invention can be used, for example, to express Dkk or Dkk-related protein (*e.g.*, via a recombinant expression vector in a host cell in gene therapy applications), to detect Dkk or Dkk-related mRNA (*e.g.*, in a biological sample) or a genetic alteration in a Dkk or Dkk-related gene, and to modulate Dkk or Dkk-related activity, as described further below. In addition, the Dkk or Dkk-related proteins can be used to screen drugs or compounds which modulate the Dkk activity as well as to treat disorders characterized by insufficient or excessive production of Dkk or Dkk-related protein or production of Dkk or Dkk-related protein forms which have decreased or aberrant activity compared to Dkk or Dkk-related wild type protein (*e.g.*, developmental disorders or proliferative diseases such as cancer as well as diseases, ocular disorders (*e.g.*, blindness) conditions or disorders characterized by abnormal cell differentiation and/or survival, an abnormal extracellular structure, or an abnormality in a defense mechanism). Moreover, the antibodies of the invention can be used to detect and isolate Dkk or Dkk-related proteins, regulate the bioavailability of Dkk or Dkk-related proteins, and modulate Dkk or Dkk-related activity. The term "an aberrant activity", as applied to an activity of a protein such as Dkk (*e.g.*, hDkk-3), refers to an activity which differs from the activity of the wild-type or native protein or which differs from the activity of

the protein in a healthy subject. An activity of a protein can be aberrant because it is stronger than the activity of its native counterpart. Alternatively, an activity can be aberrant because it is weaker or absent related to the activity of its native counterpart. An aberrant activity can also be a change in an activity. For example an aberrant protein
5 can interact with a different protein relative to its native counterpart. A cell can have an aberrant Dkk (*e.g.*, hDkk-3) activity due to overexpression or underexpression of the gene encoding Dkk.

A. Screening Assays:

10 The invention provides a method (also referred to herein as a "screening assay") for identifying modulators, *i.e.*, candidate or test compounds or agents (*e.g.*, peptides, peptidomimetics, small molecules or other drugs) which bind to Dkk or Dkk-related proteins or have a stimulatory or inhibitory effect on, for example, Dkk or Dkk-related expression or activity. Modulators can include, for example, agonists and/or
15 antagonists. The term "agonist", as used herein, is meant to refer to an agent that mimics or upregulates (*e.g.* potentiates or supplements) a Dkk or Dkk-related (*e.g.*, hDkk-3) bioactivity. An agonist can be a compound which mimics a bioactivity of a Dkk or Dkk-related protein, such as transduction of a signal from a Dkk receptor, by, *e.g.*, interacting with a hDkk-3 receptor. An agonist can also be a compound that upregulates
20 expression of a Dkk or Dkk-related gene. An agonist can also be a compound which modulates the expression or activity of a protein which is located downstream, for example, of a Dkk receptor, thereby mimicking or enhancing the effect of binding of Dkk to a Dkk receptor.

"Antagonist" as used herein is meant to refer to an agent that inhibits, decreases
25 or suppresses a bioactivity (*e.g.*, hDkk-3). An antagonist can be a compound which decreases signalling from a Dkk or Dkk-related protein, *e.g.*, a compound that is capable of binding to hDkk-3 or to a hDkk-3 receptor. A preferred antagonist inhibits the interaction between a Dkk or Dkk-related protein and another molecule, such as a Dkk receptor. Alternatively, an antagonist can be a compound that downregulates expression
30 of a Dkk or Dkk-related gene. An antagonist can also be a compound which modulates

- 79 -

the expression or activity of a protein which is located downstream of a Dkk receptor, thereby antagonizing the effect of binding of Dkk to a Dkk receptor.

In one embodiment, the invention provides assays for screening candidate or test compounds which bind to or modulate the activity of a Dkk or Dkk-related protein or polypeptide or biologically active portion thereof. In another embodiment, the invention provides assays for screening candidate or test compounds which bind to or modulate the activity of a Dkk receptor. The test compounds of the present invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the 'one-bead one-compound' library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds (Lam, K.S. (1997) *Anticancer Drug Des.* 12:145).

Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt *et al.* (1993) *Proc. Natl. Acad. Sci. U.S.A.* 90:6909; Erb *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:11422; Zuckermann *et al.* (1994). *J. Med. Chem.* 37:2678; Cho *et al.*, (1993) *Science* 261:1303; Carrell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2059; Carrell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2061; and in Gallop *et al.* (1994) *J. Med. Chem.* 37:1233.

Libraries of compounds may be presented in solution (*e.g.*, Houghten (1992) *Biotechniques* 13:412-421), or on beads (Lam (1991) *Nature* 354:82-84), chips (Fodor (1993) *Nature* 364:555-556), bacteria (Ladner USP 5,223,409), spores (Ladner USP '409), plasmids (Cull *et al.* (1992) *Proc Natl Acad Sci USA* 89:1865-1869) or on phage (Scott and Smith (1990) *Science* 249:386-390); (Devlin (1990) *Science* 249:404-406); (Cwirla *et al.* (1990) *Proc. Natl. Acad. Sci.* 87:6378-6382); (Felici (1991) *J. Mol. Biol.* 222:301-310); (Ladner *supra.*).

In one embodiment, an assay is a cell-based assay in which a cell which expresses a Dkk receptor on the cell surface is contacted with a test compound and the ability of the test compound to bind to a Dkk receptor determined. The cell, for

- 80 -

example, can be of mammalian origin or a yeast cell. Determining the ability of the test compound to bind to a Dkk receptor can be accomplished, for example, by coupling the test compound with a radioisotope or enzymatic label such that binding of the test compound to the Dkk receptor can be determined by detecting the labeled compound in a complex. For example, test compounds can be labeled with ^{125}I , ^{35}S , ^{14}C , or ^3H , either directly or indirectly, and the radioisotope detected by direct counting of radioemission or by scintillation counting. Alternatively, test compounds can be enzymatically labeled with, for example, horseradish peroxidase, alkaline phosphatase, or luciferase, and the enzymatic label detected by determination of conversion of an appropriate substrate to product.

It is also within the scope of this invention to determine the ability of a test compound to interact with a Dkk receptor without the labeling of any of the interactants. For example, a microphysiometer can be used to detect the interaction of a test compound with a Dkk receptor without the labeling of either the test compound or the receptor. McConnell, H. M. *et al.* (1992) *Science* 257:1906-1912. As used herein, a "microphysiometer" (*e.g.*, Cytosensor™) is an analytical instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indicator of the interaction between ligand and receptor.

In a preferred embodiment, the assay comprises contacting a cell which expresses a Dkk receptor on the cell surface with a Dkk protein or biologically-active portion thereof, to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with a Dkk receptor, wherein determining the ability of the test compound to interact with a Dkk receptor comprises determining the ability of the test compound to preferentially bind to the Dkk receptor as compared to the ability of Dkk, or a biologically active portion thereof, to bind to the receptor.

In another embodiment, an assay is a cell-based assay comprising contacting a cell expressing a Dkk target molecule with a test compound and determining the ability of the test compound to modulate (*e.g.* stimulate or inhibit) the activity of the Dkk target molecule. Determining the ability of the test compound to modulate the activity of a

Dkk target molecule can be accomplished, for example, by determining the ability of the Dkk protein to bind to or interact with the Dkk target molecule.

Determining the ability of the Dkk protein to bind to or interact with a Dkk target molecule can be accomplished by one of the methods described above for determining
5 direct binding. In a preferred embodiment, determining the ability of the Dkk protein to bind to or interact with a Dkk target molecule can be accomplished by determining the activity of the target molecule. For example, the activity of the target molecule can be determined by detecting induction of a cellular second messenger of the target (*i.e.* intracellular Ca^{2+} , diacylglycerol, IP_3 , etc.), detecting catalytic/enzymatic activity of
10 the target an appropriate substrate, detecting the induction of a reporter gene (comprising a Dkk-responsive regulatory element operatively linked to a nucleic acid encoding a detectable marker, *e.g.*, luciferase), or detecting a cellular response, for example, development, differentiation or rate of proliferation.

In yet another embodiment, an assay of the present invention is a cell-free assay
15 in which a Dkk or Dkk-related protein or biologically active portion thereof is contacted with a test compound and the ability of the test compound to bind to the Dkk or Dkk-related protein or biologically active portion thereof is determined. Binding of the test compound to the Dkk or Dkk-related protein can be determined either directly or indirectly as described above. In a preferred embodiment, the assay includes contacting
20 the Dkk or Dkk-related protein or biologically active portion thereof with a known compound which binds Dkk or the Dkk-related protein to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with a Dkk or Dkk-related protein, wherein determining the ability of the test compound to interact with a Dkk or Dkk-related protein comprises
25 determining the ability of the test compound to preferentially bind to Dkk or a Dkk-related protein or biologically active portion thereof as compared to the known compound.

In another embodiment, the assay is a cell-free assay in which a Dkk or Dkk-related protein or biologically active portion thereof is contacted with a test compound
30 and the ability of the test compound to modulate (*e.g.*, stimulate or inhibit) the activity of the Dkk or Dkk-related protein or biologically active portion thereof is determined.

Determining the ability of the test compound to modulate the activity of a Dkk or Dkk-related protein can be accomplished, for example, by determining the ability of the Dkk or Dkk-related protein to bind to a target molecule (*e.g.*, a Dkk-target molecule) by one of the methods described above for determining direct binding. Determining the ability
5 of the Dkk or Dkk-related protein to bind to a target molecule can also be accomplished using a technology such as real-time Biomolecular Interaction Analysis (BIA).

Sjolander, S. and Urbaniczky, C. (1991) *Anal. Chem.* 63:2338-2345 and Szabo *et al.*

(1995) *Curr. Opin. Struct. Biol.* 5:699-705. As used herein, "BIA" is a technology for studying biospecific interactions in real time, without labeling any of the interactants

10 (*e.g.*, BIAcore™). Changes in the optical phenomenon surface plasmon resonance (SPR) can be used as an indication of real-time reactions between biological molecules.

In an alternative embodiment, determining the ability of the test compound to modulate the activity of a Dkk or Dkk-related protein can be accomplished by determining the ability of the Dkk or Dkk-related protein to further modulate the activity
15 of a target molecule (*e.g.*, a Dkk-target molecule). For example, the catalytic/enzymatic activity of the target molecule on an appropriate substrate can be determined as previously described.

In yet another embodiment, the cell-free assay involves contacting a Dkk or Dkk-related protein or biologically active portion thereof with a known compound which
20 binds the Dkk or Dkk-related protein to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with the Dkk or Dkk-related protein, wherein determining the ability of the test compound to interact with the Dkk or Dkk-related protein comprises determining the ability of the Dkk or Dkk-related protein to preferentially bind to or modulate the
25 activity of a target molecule (*e.g.*, a Dkk target molecule).

In many drug screening programs which test libraries of compounds and natural extracts, high throughput assays are desirable in order to maximize the number of compounds surveyed in a given period of time. Assays which are performed in cell-free systems, such as may be derived with purified or semi-purified proteins, are often
30 preferred as "primary" screens in that they can be generated to permit rapid development and relatively easy detection of an alteration in a molecular target which is mediated by

a test compound. Moreover, the effects of cellular toxicity and/or bioavailability of the test compound can be generally ignored in the *in vitro* system, the assay instead being focused primarily on the effect of the drug on the molecular target as may be manifest in an alteration of binding affinity with upstream or downstream elements. Accordingly, in

5 an exemplary screening assay of the present invention, the compound of interest is contacted with a Dkk (*e.g.*, hDkk-3) protein or a Dkk (*e.g.*, hDkk-3) binding partner, *e.g.*, a receptor. The receptor can be soluble or the receptor can be present on a cell surface. To the mixture of the compound and the Dkk protein or Dkk binding partner is then added a composition containing a Dkk binding partner or a Dkk protein,
10 respectively. Detection and quantification of complexes of Dkk proteins and Dkk binding partners provide a means for determining a compound's efficacy at inhibiting (or potentiating) complex formation between Dkk and a binding partner. The efficacy of the compound can be assessed by generating dose response curves from data obtained using various concentrations of the test compound. Moreover, a control assay can also
15 be performed to provide a baseline for comparison. In the control assay, isolated and purified Dkk polypeptide or binding partner is added to a composition containing the Dkk binding partner or Dkk polypeptide, and the formation of a complex is quantitated in the absence of the test compound.

The cell-free assays of the present invention are amenable to use of both soluble
20 and/or membrane-bound forms of isolated proteins (*e.g.* Dkk proteins or biologically active portions thereof or Dkk target molecules). In the case of cell-free assays in which a membrane-bound form an isolated protein is used (*e.g.*, a Dkk target molecule or receptor) it may be desirable to utilize a solubilizing agent such that the membrane-bound form of the isolated protein is maintained in solution. Examples of such
25 solubilizing agents include non-ionic detergents such as n-octylglucoside, n-dodecylglucoside, n-dodecylmaltoside, octanoyl-N-methylglucamide, decanoyl-N-methylglucamide, Triton[®] X-100, Triton[®] X-114, Thesit[®], Isotridecypoly(ethylene glycol ether)_n, 3-[(3-cholamidopropyl)dimethylamminio]-1-propane sulfonate (CHAPS), 3-[(3-cholamidopropyl)dimethylamminio]-2-hydroxy-1-propane sulfonate
30 (CHAPSO), or N-dodecyl=N,N-dimethyl-3-ammonio-1-propane sulfonate.

In more than one embodiment of the above assay methods of the present invention, it may be desirable to immobilize either Dkk, a Dkk-related protein or a target molecule to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test compound to a Dkk or Dkk-related protein, or interaction of a Dkk or Dkk-related protein with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtitre plates, test tubes, and micro-centrifuge tubes. In one embodiment, a fusion protein can be provided which adds a domain that allows one or both of the proteins to be bound to a matrix. For example, glutathione-S-transferase/Dkk fusion proteins or glutathione-S-transferase/target fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtitre plates, which are then combined with the test compound or the test compound and either the non-adsorbed target protein or Dkk protein, and the mixture incubated under conditions conducive to complex formation (*e.g.*, at physiological conditions for salt and pH). Following incubation, the beads or microtitre plate wells are washed to remove any unbound components, the matrix immobilized in the case of beads, complex determined either directly or indirectly, for example, as described above. Alternatively, the complexes can be dissociated from the matrix, and the level of Dkk binding or activity determined using standard techniques.

Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either a Dkk protein, Dkk-related protein, or a Dkk target molecule can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated protein or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well known in the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, IL), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies reactive with Dkk, Dkk-related protein, or target molecules but which do not interfere with binding of the protein to its target molecule can be derivatized to the wells of the plate, and unbound target, Dkk, or Dkk-related protein trapped in the wells by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized

complexes, include immunodetection of complexes using antibodies reactive with the Dkk or Dkk-related protein or target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the Dkk or Dkk-related protein or target molecule.

5 In another embodiment, modulators of Dkk or Dkk-related expression are identified in a method wherein a cell is contacted with a candidate compound and the expression of Dkk or Dkk-related mRNA or protein in the cell is determined. The level of expression of mRNA or protein in the presence of the candidate compound is compared to the level of expression of mRNA or protein in the absence of the candidate
10 compound. The candidate compound can then be identified as a modulator of Dkk or Dkk-related expression based on this comparison. For example, when expression of Dkk mRNA or protein is greater (statistically significantly greater) in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of Dkk mRNA or protein expression. Alternatively, when expression of Dkk
15 mRNA or protein is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of Dkk mRNA or protein expression. The level of Dkk or Dkk-related mRNA or protein expression in the cells can be determined by methods described herein for detecting Dkk mRNA or protein.

20 In yet another aspect of the invention, the Dkk or Dkk-related proteins can be used as "bait proteins" in a two-hybrid assay or three-hybrid assay (see, *e.g.*, U.S. Patent No. 5,283,317; Zervos *et al.* (1993) *Cell* 72:223-232; Madura *et al.* (1993) *J. Biol. Chem.* 268:12046-12054; Bartel *et al.*, (1993) *Biotechniques* 14:920-924; Iwabuchi *et al.* (1993) *Oncogene* 8:1693-1696; and Brent WO94/10300), to identify other proteins,
25 which bind to or interact with Dkk or Dkk-related proteins ("binding proteins" or "bp") and modulate Dkk or Dkk-related activity. Such binding proteins are also likely to be involved in the propagation of signals by the Dkk or Dkk-related proteins as, for example, downstream elements of a Dkk-mediated signaling pathway. Alternatively, such binding proteins are likely to be cell-surface molecules associated with non-Dkk
30 expressing cells, wherein such binding proteins are involved in signal transduction.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. In one construct, the gene that codes for a Dkk protein is fused to a gene encoding the DNA binding domain of a known transcription factor (*e.g.*, GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact, *in vivo*, forming a Dkk-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (*e.g.*, LacZ) which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene which encodes the protein which interacts with the Dkk or Dkk-related protein.

This invention further pertains to novel agents identified by the above-described screening assays and to processes for producing such agents by use of these assays. Accordingly, in one embodiment, the present invention includes a compound or agent obtainable by a method comprising the steps of any one of the aforementioned screening assays (*e.g.*, cell-based assays or cell-free assays). For example, in one embodiment, the invention includes a compound or agent obtainable by a method comprising contacting a cell which expresses a target molecule with a test compound and the determining the ability of the test compound to bind to, or modulate the activity of, the target molecule. In another embodiment, the invention includes a compound or agent obtainable by a method comprising contacting a cell which expresses a target molecule with a Dkk or Dkk-related protein or biologically-active portion thereof, to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with, or modulate the activity of, the target molecule. In another embodiment, the invention includes a compound or agent obtainable by a method comprising contacting a Dkk or Dkk-related protein or biologically active portion thereof with a test compound and determining the ability of the test compound to

bind to, or modulate (*e.g.*, stimulate or inhibit) the activity of, the Dkk or Dkk-related protein or biologically active portion thereof. In yet another embodiment, the present invention includes a compound or agent obtainable by a method comprising contacting a Dkk or Dkk-related protein or biologically active portion thereof with a known

- 5 compound which binds the Dkk or Dkk-related protein to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with, or modulate the activity of the Dkk or Dkk-related protein.

Accordingly, it is within the scope of this invention to further use an agent
10 identified as described herein in an appropriate animal model. For example, an agent identified as described herein (*e.g.*, a Dkk modulating agent, an antisense Dkk nucleic acid molecule, a Dkk-specific antibody, or a Dkk-binding partner) can be used in an animal model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal
15 model to determine the mechanism of action of such an agent.

The present invention also pertains to uses of novel agents identified by the above-described screening assays for diagnoses, prognoses, and treatments as described herein. Accordingly, it is within the scope of the present invention to use such agents in the design, formulation, synthesis, manufacture, and/or production of a drug or
20 pharmaceutical composition for use in diagnosis, prognosis, or treatment, as described herein. For example, in one embodiment, the present invention includes a method of synthesizing or producing a drug or pharmaceutical composition by reference to the structure and/or properties of a compound obtainable by one of the above-described screening assays. For example, a drug or pharmaceutical composition can be
25 synthesized based on the structure and/or properties of a compound obtained by a method in which a cell which expresses a target molecule (*e.g.*, a Dkk target molecule) is contacted with a test compound and the ability of the test compound to bind to, or modulate the activity of, the target molecule is determined. In another exemplary embodiment, the present invention includes a method of synthesizing or producing a
30 drug or pharmaceutical composition based on the structure and/or properties of a compound obtainable by a method in which a Dkk or Dkk-related protein or biologically

active portion thereof is contacted with a test compound and the ability of the test compound to bind to, or modulate (*e.g.*, stimulate or inhibit) the activity of, the Dkk or Dkk-related protein or biologically active portion thereof is determined.

5 B. Detection Assays

Portions or fragments of the cDNA sequences identified herein (and the corresponding complete gene sequences) can be used in numerous ways as polynucleotide reagents. For example, these sequences can be used to: (i) map their respective genes on a chromosome; and, thus, locate gene regions associated with
10 genetic disease; (ii) identify an individual from a minute biological sample (tissue typing); and (iii) aid in forensic identification of a biological sample. These applications are described in the subsections below.

1. Chromosome Mapping

15 Once the sequence (or a portion of the sequence) of a gene has been isolated, this sequence can be used to map the location of the gene on a chromosome. This process is called chromosome mapping. Accordingly, portions or fragments of the Dkk or Dkk-related nucleotide sequences, described herein, can be used to map the location of the Dkk or Dkk-related genes on a chromosome. The mapping of the Dkk or Dkk-related
20 sequences to chromosomes is an important first step in correlating these sequences with genes associated with disease.

Briefly, Dkk or Dkk-related genes can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp in length) from the Dkk or Dkk-related nucleotide sequences. Computer analysis of the Dkk or Dkk-related sequences can be used to
25 predict primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers can then be used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the Dkk or Dkk-related sequences will yield an amplified fragment.

Somatic cell hybrids are prepared by fusing somatic cells from different mammals (*e.g.*, human and mouse cells). As hybrids of human and mouse cells grow and divide, they gradually lose human chromosomes in random order, but retain the mouse chromosomes. By using media in which mouse cells cannot grow, because they
5 lack a particular enzyme, but human cells can, the one human chromosome that contains the gene encoding the needed enzyme, will be retained. By using various media, panels of hybrid cell lines can be established. Each cell line in a panel contains either a single human chromosome or a small number of human chromosomes, and a full set of mouse chromosomes, allowing easy mapping of individual genes to specific human
10 chromosomes. (D'Eustachio P. *et al.*, (1983) *Science* 220:919-924). Somatic cell hybrids containing only fragments of human chromosomes can also be produced by using human chromosomes with translocations and deletions.

PCR mapping of somatic cell hybrids is a rapid procedure for assigning a particular sequence to a particular chromosome. Three or more sequences can be
15 assigned per day using a single thermal cycler. Using the Dkk or Dkk-related nucleotide sequences to design oligonucleotide primers, sublocalization can be achieved with panels of fragments from specific chromosomes. Other mapping strategies which can similarly be used to map a 9o, 1p, or 1v sequence to its chromosome include *in situ* hybridization (described in Fan, Y. *et al.*, (1990) *PNAS*, 87:6223-27), pre-screening with
20 labeled flow-sorted chromosomes, and pre-selection by hybridization to chromosome specific cDNA libraries.

Fluorescence *in situ* hybridization (FISH) of a DNA sequence to a metaphase chromosomal spread can further be used to provide a precise chromosomal location in one step. Chromosome spreads can be made using cells whose division has been
25 blocked in metaphase by a chemical such as colcemid that disrupts the mitotic spindle. The chromosomes can be treated briefly with trypsin, and then stained with Giemsa. A pattern of light and dark bands develops on each chromosome, so that the chromosomes can be identified individually. The FISH technique can be used with a DNA sequence as short as 500 or 600 bases. However, clones larger than 1,000 bases have a higher
30 likelihood of binding to a unique chromosomal location with sufficient signal intensity for simple detection. Preferably 1,000 bases, and more preferably 2,000 bases will

- 90 -

suffice to get good results at a reasonable amount of time. For a review of this technique, see Verma *et al.*, Human Chromosomes: A Manual of Basic Techniques (Pergamon Press, New York 1988).

Reagents for chromosome mapping can be used individually to mark a single chromosome or a single site on that chromosome, or panels of reagents can be used for marking multiple sites and/or multiple chromosomes. Reagents corresponding to noncoding regions of the genes actually are preferred for mapping purposes. Coding sequences are more likely to be conserved within gene families, thus increasing the chance of cross hybridizations during chromosomal mapping.

Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. (Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man, available on-line through Johns Hopkins University Welch Medical Library). The relationship between a gene and a disease, mapped to the same chromosomal region, can then be identified through linkage analysis (co-inheritance of physically adjacent genes), described in, for example, Egeland, J. *et al.* (1987) *Nature*, 325:783-787.

Moreover, differences in the DNA sequences between individuals affected and unaffected with a disease associated with a Dkk or Dkk-related gene, can be determined. If a mutation is observed in some or all of the affected individuals but not in any unaffected individuals, then the mutation is likely to be the causative agent of the particular disease. Comparison of affected and unaffected individuals generally involves first looking for structural alterations in the chromosomes, such as deletions or translocations that are visible from chromosome spreads or detectable using PCR based on that DNA sequence. Ultimately, complete sequencing of genes from several individuals can be performed to confirm the presence of a mutation and to distinguish mutations from polymorphisms.

2. Tissue Typing

The Dkk or Dkk-related sequences of the present invention can also be used to identify individuals from minute biological samples. The United States military, for example, is considering the use of restriction fragment length polymorphism (RFLP) for

- 91 -

identification of its personnel. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identification. This method does not suffer from the current limitations of "Dog Tags" which can be lost, switched, or stolen, making positive identification difficult. The sequences of the present invention are useful as additional DNA markers for RFLP (described in U.S. Patent 5,272,057).

Furthermore, the sequences of the present invention can be used to provide an alternative technique which determines the actual base-by-base DNA sequence of selected portions of an individual's genome. Thus, the Dkk or Dkk-related nucleotide sequences described herein can be used to prepare two PCR primers from the 5' and 3' ends of the sequences. These primers can then be used to amplify an individual's DNA and subsequently sequence it.

Panels of corresponding DNA sequences from individuals, prepared in this manner, can provide unique individual identifications, as each individual will have a unique set of such DNA sequences due to allelic differences. The sequences of the present invention can be used to obtain such identification sequences from individuals and from tissue. The Dkk or Dkk-related nucleotide sequences of the invention uniquely represent portions of the human genome. Allelic variation occurs to some degree in the coding regions of these sequences, and to a greater degree in the noncoding regions. It is estimated that allelic variation between individual humans occurs with a frequency of about once per each 500 bases. Each of the sequences described herein can, to some degree, be used as a standard against which DNA from an individual can be compared for identification purposes. Because greater numbers of polymorphisms occur in the noncoding regions, fewer sequences are necessary to differentiate individuals. The noncoding sequences of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20, can comfortably provide positive individual identification with a panel of perhaps 10 to 1,000 primers which each yield a noncoding amplified sequence of 100 bases. If predicted coding sequences, such as those in SEQ ID NO:3, SEQ ID NO:6, SEQ ID NO:9, SEQ ID NO:15 or SEQ ID NO:22 are used, a more appropriate number of primers for positive individual identification would be 500-2,000.

- 92 -

If a panel of reagents from Dkk or Dkk-related nucleotide sequences described herein is used to generate a unique identification database for an individual, those same reagents can later be used to identify tissue from that individual. Using the unique identification database, positive identification of the individual, living or dead, can be
5 made from extremely small tissue samples.

3. Use of Partial Dkk or Dkk-related Sequences in Forensic Biology

DNA-based identification techniques can also be used in forensic biology.

Forensic biology is a scientific field employing genetic typing of biological evidence
10 found at a crime scene as a means for positively identifying, for example, a perpetrator of a crime. To make such an identification, PCR technology can be used to amplify DNA sequences taken from very small biological samples such as tissues, *e.g.*, hair or skin, or body fluids, *e.g.*, blood, saliva, or semen found at a crime scene. The amplified sequence can then be compared to a standard, thereby allowing identification of the
15 origin of the biological sample.

The sequences of the present invention can be used to provide polynucleotide reagents, *e.g.*, PCR primers, targeted to specific loci in the human genome, which can enhance the reliability of DNA-based forensic identifications by, for example, providing another "identification marker" (*i.e.* another DNA sequence that is unique to a particular
20 individual). As mentioned above, actual base sequence information can be used for identification as an accurate alternative to patterns formed by restriction enzyme generated fragments. Sequences targeted to noncoding regions of SEQ ID NOs:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20 are particularly appropriate for this use as greater numbers of polymorphisms occur in the noncoding regions,
25 making it easier to differentiate individuals using this technique. Examples of polynucleotide reagents include the Dkk nucleotide sequences or portions thereof, *e.g.*, fragments derived from the noncoding regions of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, or SEQ ID NO:20, having a length of at least 20 bases, preferably at least 30 bases.

- 93 -

The Dkk or Dkk-related nucleotide sequences described herein can further be used to provide polynucleotide reagents, *e.g.*, labeled or labelable probes which can be used in, for example, an *in situ* hybridization technique, to identify a specific tissue, *e.g.*, brain tissue. This can be very useful in cases where a forensic pathologist is presented
5 with a tissue of unknown origin. Panels of such Dkk or Dkk-related probes can be used to identify tissue by species and/or by organ type.

In a similar fashion, these reagents, *e.g.*, Dkk or Dkk-related primers or probes can be used to screen tissue culture for contamination (*i.e.* screen for the presence of a mixture of different types of cells in a culture).

10

C. Predictive Medicine:

The present invention also pertains to the field of predictive medicine in which diagnostic assays, prognostic assays, and monitoring clinical trials are used for prognostic (predictive) purposes to thereby treat an individual prophylactically.

15 Accordingly, one aspect of the present invention relates to diagnostic assays for determining Dkk or Dkk-related protein and/or nucleic acid expression as well as Dkk or Dkk-related activity, in the context of a biological sample (*e.g.*, blood, serum, cells, tissue) to thereby determine whether an individual is afflicted with a disease or disorder, or is at risk of developing a disorder, associated with aberrant Dkk or Dkk-related
20 expression or activity, such as aberrant cell proliferation, differentiation, and/or survival resulting for example in a neurodegenerative disease (*e.g.*, Alzheimer's disease, Parkinson's disease, Huntington's chorea, amyotrophic lateral sclerosis and the like, as well as spinocerebellar degenerations) or cancer (for example, cancers of the epithelia (*e.g.*, carcinomas of the pancreas, stomach, liver, secretory glands (*e.g.*,
25 adenocarcinoma) bladder, lung, breast, skin (*e.g.*, malignant melanoma), reproductive tract including prostate gland, ovary, cervix and uterus); cancers of the hematopoietic and immune system (*e.g.*, leukemias and lymphomas); cancers of the central nervous, brain system and eye (*e.g.*, gliomas, glioblastoma, neuroblastoma and retinoblastoma); and cancers of connective tissues, bone, muscles and vasculature (*e.g.*, sarcomas)). The
30 invention also provides for prognostic (or predictive) assays for determining whether an individual is at risk of developing a disorder associated with Dkk or Dkk-related protein,

nucleic acid expression or activity. For example, mutations in a Dkk or Dkk-related gene can be assayed in a biological sample. Such assays can be used for prognostic or predictive purpose to thereby prophylactically treat an individual prior to the onset of a disorder characterized by or associated with Dkk or Dkk-related protein, nucleic acid
5 expression or activity.

Another aspect of the invention pertains to monitoring the influence of agents (*e.g.*, drugs, compounds) on the expression or activity of Dkk or Dkk-related in clinical trials.

These and other agents are described in further detail in the following sections.

10

1. Diagnostic Assays

An exemplary method for detecting the presence or absence of Dkk or Dkk-related protein or nucleic acid in a biological sample involves obtaining a biological sample from a test subject and contacting the biological sample with a compound or an
15 agent capable of detecting Dkk or Dkk-related protein or nucleic acid (*e.g.*, mRNA, genomic DNA) that encodes Dkk or Dkk-related protein such that the presence of Dkk or Dkk-related protein or nucleic acid is detected in the biological sample. A preferred agent for detecting Dkk or Dkk-related mRNA or genomic DNA is a labeled nucleic acid probe capable of hybridizing to Dkk or Dkk-related mRNA or genomic DNA. The
20 nucleic acid probe can be, for example, a full-length Dkk nucleic acid, such as the nucleic acid of SEQ ID NO: 1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:20, the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with
25 ATCC as Accession Number _____, or a portion thereof, such as an oligonucleotide of at least 15, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically hybridize under stringent conditions to Dkk or Dkk-related mRNA or genomic DNA. Other suitable probes for use in the diagnostic assays of the invention are described herein.

- 95 -

A preferred agent for detecting Dkk or Dkk-related protein is an antibody capable of binding to the protein, preferably an antibody with a detectable label. Antibodies can be polyclonal, or more preferably, monoclonal. An intact antibody, or a fragment thereof (*e.g.*, Fab or F(ab')₂) can be used. The term "labeled", with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (*i.e.*, physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include detection of a primary antibody using a fluorescently labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently labeled streptavidin. The term "biological sample" is intended to include tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject. That is, the detection method of the invention can be used to detect Dkk or Dkk-related mRNA, protein, or genomic DNA in a biological sample *in vitro* as well as *in vivo*. For example, *in vitro* techniques for detection of Dkk or Dkk-related mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro* techniques for detection of Dkk or Dkk-related protein include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence. *In vitro* techniques for detection of Dkk or Dkk-related genomic DNA include Southern hybridizations. Furthermore, *in vivo* techniques for detection of Dkk or Dkk-related protein include introducing into a subject a labeled antibody. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques.

In one embodiment, the biological sample contains protein molecules from the test subject. Alternatively, the biological sample can contain mRNA molecules from the test subject or genomic DNA molecules from the test subject. A preferred biological sample is a serum sample isolated by conventional means from a subject.

In another embodiment, the methods further involve obtaining a control biological sample from a control subject, contacting the control sample with a compound or agent capable of detecting Dkk or Dkk-related protein, mRNA, or genomic DNA, such that the presence of Dkk or Dkk-related protein, mRNA or genomic DNA is

- 96 -

detected in the biological sample, and comparing the presence of Dkk or Dkk-related protein, mRNA or genomic DNA in the control sample with the presence of Dkk or Dkk-related protein, mRNA or genomic DNA in the test sample.

The invention also encompasses kits for detecting the presence of Dkk or a Dkk-related protein in a biological sample. For example, the kit can comprise a labeled compound or agent capable of detecting Dkk or Dkk-related protein or mRNA in a biological sample; means for determining the amount of Dkk or Dkk-related protein or mRNA in the sample; and means for comparing the amount of Dkk or Dkk-related protein or mRNA in the sample with a standard. The compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect Dkk or Dkk-related protein or nucleic acid.

2. Prognostic Assays

The diagnostic methods described herein can furthermore be utilized to identify subjects having or at risk of developing a disease or disorder associated with aberrant Dkk expression or activity. For example, the assays described herein, such as the preceding diagnostic assays or the following assays, can be utilized to identify a subject having or at risk of developing a disorder associated with Dkk or Dkk-related protein, nucleic acid expression or activity such as a proliferative disorder, a differentiative or developmental disorder, a hematopoietic disorder as well as diseases, conditions or disorders characterized by abnormal cell survival, abnormal extracellular structure, or an abnormality in a defense mechanism. Alternatively, the prognostic assays can be utilized to identify a subject having or at risk for developing a differentiative or proliferative disease (*e.g.*, cancer). Thus, the present invention provides a method for identifying a disease or disorder associated with aberrant Dkk or Dkk-related expression or activity in which a test sample is obtained from a subject and Dkk or Dkk-related protein or nucleic acid (*e.g.*, mRNA, genomic DNA) is detected, wherein the presence of Dkk or Dkk-related protein or nucleic acid is diagnostic for a subject having or at risk of developing a disease or disorder associated with aberrant Dkk or Dkk-related expression or activity. As used herein, a "test sample" refers to a biological sample obtained from a

- 97 -

subject of interest. For example, a test sample can be a biological fluid (*e.g.*, serum), cell sample, or tissue.

Furthermore, the prognostic assays described herein can be used to determine whether a subject can be administered an agent (*e.g.*, an agonist, antagonist, peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate) to treat a disease or disorder associated with aberrant Dkk or Dkk-related expression or activity. For example, such methods can be used to determine whether a subject can be effectively treated with an agent for a disorder, such as a proliferative disorder, a differentiative or developmental disorder, a hematopoietic disorder, as well disorders characterized by abnormal cell survival, an abnormal extracellular structure, or an abnormality in a defense mechanism. Alternatively, such methods can be used to determine whether a subject can be effectively treated with an agent for a differentiative or proliferative disease (*e.g.*, cancer). Thus, the present invention provides methods for determining whether a subject can be effectively treated with an agent for a disorder associated with aberrant Dkk or Dkk-related expression or activity in which a test sample is obtained and Dkk or Dkk-related protein or nucleic acid expression or activity is detected (*e.g.*, wherein the abundance of Dkk or Dkk-related protein or nucleic acid expression or activity is diagnostic for a subject that can be administered the agent to treat a disorder associated with aberrant Dkk or Dkk-related expression or activity.)

The methods of the invention can also be used to detect genetic alterations in a Dkk or Dkk-related gene, thereby determining if a subject with the altered gene is at risk for a disorder characterized by aberrant development, aberrant cellular differentiation, aberrant cellular proliferation or an aberrant hematopoietic response. In preferred embodiments, the methods include detecting, in a sample of cells from the subject, the presence or absence of a genetic alteration characterized by at least one of an alteration affecting the integrity of a gene encoding a Dkk or Dkk-related-protein, or the mis-expression of the Dkk or Dkk-related gene. For example, such genetic alterations can be detected by ascertaining the existence of at least one of 1) a deletion of one or more nucleotides from a Dkk or Dkk-related gene; 2) an addition of one or more nucleotides to a Dkk or Dkk-related gene; 3) a substitution of one or more nucleotides of a Dkk or Dkk-related gene, 4) a chromosomal rearrangement of a Dkk or Dkk-related gene; 5) an

alteration in the level of a messenger RNA transcript of a Dkk or Dkk-related gene, 6) aberrant modification of a Dkk or Dkk-related gene, such as of the methylation pattern of the genomic DNA, 7) the presence of a non-wild type splicing pattern of a messenger RNA transcript of a Dkk or Dkk-related gene, 8) a non-wild type level of a Dkk or Dkk-related-protein, 9) allelic loss of a Dkk or Dkk-related gene, and 10) inappropriate post-translational modification of a Dkk or Dkk-related-protein. As described herein, there are a large number of assay techniques known in the art which can be used for detecting alterations in a Dkk or Dkk-related gene. A preferred biological sample is a tissue or serum sample isolated by conventional means from a subject.

10 In certain embodiments, detection of the alteration involves the use of a probe/primer in a polymerase chain reaction (PCR) (see, *e.g.*, U.S. Patent Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (see, *e.g.*, Landegran *et al.* (1988) *Science* 241:1077-1080; and Nakazawa *et al.* (1994) *PNAS* 91:360-364), the latter of which can be particularly
15 useful for detecting point mutations in the Dkk or Dkk-related-gene (see Abravaya *et al.* (1995) *Nucleic Acids Res.* 23:675-682). This method can include the steps of collecting a sample of cells from a patient, isolating nucleic acid (*e.g.*, genomic, mRNA or both) from the cells of the sample, contacting the nucleic acid sample with one or more primers which specifically hybridize to a Dkk or Dkk-related gene under conditions such
20 that hybridization and amplification of the Dkk or Dkk-related-gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. It is anticipated that PCR and/or LCR may be desirable to use as a preliminary amplification step in conjunction with any of the techniques used for detecting mutations described
25 herein.

Alternative amplification methods include: self sustained sequence replication (Guatelli, J.C. *et al.*, 1990, *Proc. Natl. Acad. Sci. USA* 87:1874-1878), transcriptional amplification system (Kwoh, D.Y. *et al.*, 1989, *Proc. Natl. Acad. Sci. USA* 86:1173-1177), Q-Beta Replicase (Lizardi, P.M. *et al.*, 1988, *Bio/Technology* 6:1197), or any
30 other nucleic acid amplification method, followed by the detection of the amplified molecules using techniques well known to those of skill in the art. These detection

- 99 -

schemes are especially useful for the detection of nucleic acid molecules if such molecules are present in very low numbers.

In an alternative embodiment, mutations in a Dkk or Dkk-related gene from a sample cell can be identified by alterations in restriction enzyme cleavage patterns. For example, sample and control DNA is isolated, amplified (optionally), digested with one or more restriction endonucleases, and fragment length sizes are determined by gel electrophoresis and compared. Differences in fragment length sizes between sample and control DNA indicates mutations in the sample DNA. Moreover, the use of sequence specific ribozymes (see, for example, U.S. Patent No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site.

In other embodiments, genetic mutations in a Dkk or Dkk-related gene can be identified by hybridizing a sample and control nucleic acids, *e.g.*, DNA or RNA, to high density arrays containing hundreds or thousands of oligonucleotides probes (Cronin, M.T. *et al.* (1996) *Human Mutation* 7: 244-255; Kozal, M.J. *et al.* (1996) *Nature Medicine* 2: 753-759). For example, genetic mutations in Dkk can be identified in two dimensional arrays containing light-generated DNA probes as described in Cronin, M.T. *et al. supra*. Briefly, a first hybridization array of probes can be used to scan through long stretches of DNA in a sample and control to identify base changes between the sequences by making linear arrays of sequential overlapping probes. This step allows the identification of point mutations. This step is followed by a second hybridization array that allows the characterization of specific mutations by using smaller, specialized probe arrays complementary to all variants or mutations detected. Each mutation array is composed of parallel probe sets, one complementary to the wild-type gene and the other complementary to the mutant gene.

In yet another embodiment, any of a variety of sequencing reactions known in the art can be used to directly sequence the Dkk or Dkk-related gene and detect mutations by comparing the sequence of the sample Dkk or Dkk-related sequence with the corresponding wild-type (control) sequence. Examples of sequencing reactions include those based on techniques developed by Maxim and Gilbert ((1977) *PNAS* 74:560) or Sanger ((1977) *PNAS* 74:5463). It is also contemplated that any of a variety

of automated sequencing procedures can be utilized when performing the diagnostic assays ((1995) *Biotechniques* 19:448), including sequencing by mass spectrometry (see, e.g., PCT International Publication No. WO 94/16101; Cohen *et al.* (1996) *Adv. Chromatogr.* 36:127-162; and Griffin *et al.* (1993) *Appl. Biochem. Biotechnol.* 38:147-
5 159).

Other methods for detecting mutations in the Dkk or Dkk-related gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA heteroduplexes (Myers *et al.* (1985) *Science* 230:1242). In general, the art technique of "mismatch cleavage" starts by providing heteroduplexes of
10 formed by hybridizing (labeled) RNA or DNA containing the wild-type Dkk or Dkk-related sequence with potentially mutant RNA or DNA obtained from a tissue sample. The double-stranded duplexes are treated with an agent which cleaves single-stranded regions of the duplex such as which will exist due to basepair mismatches between the control and sample strands. For instance, RNA/DNA duplexes can be treated with
15 RNase and DNA/DNA hybrids treated with S1 nuclease to enzymatically digesting the mismatched regions. In other embodiments, either DNA/DNA or RNA/DNA duplexes can be treated with hydroxylamine or osmium tetroxide and with piperidine in order to digest mismatched regions. After digestion of the mismatched regions, the resulting material is then separated by size on denaturing polyacrylamide gels to determine the
20 site of mutation. See, for example, Cotton *et al.* (1988) *Proc. Natl Acad Sci USA* 85:4397; Saleeba *et al.* (1992) *Methods Enzymol.* 217:286-295. In a preferred embodiment, the control DNA or RNA can be labeled for detection.

In still another embodiment, the mismatch cleavage reaction employs one or more proteins that recognize mismatched base pairs in double-stranded DNA (so called
25 "DNA mismatch repair" enzymes) in defined systems for detecting and mapping point mutations in Dkk cDNAs obtained from samples of cells. For example, the mutY enzyme of *E. coli* cleaves A at G/A mismatches and the thymidine DNA glycosylase from HeLa cells cleaves T at G/T mismatches (Hsu *et al.* (1994) *Carcinogenesis* 15:1657-1662). According to an exemplary embodiment, a probe based on a Dkk
30 sequence, e.g., a wild-type Dkk sequence, is hybridized to a cDNA or other DNA product from a test cell(s). The duplex is treated with a DNA mismatch repair enzyme,

and the cleavage products, if any, can be detected from electrophoresis protocols or the like. See, for example, U.S. Patent No. 5,459,039.

In other embodiments, alterations in electrophoretic mobility will be used to identify mutations in Dkk or Dkk-related genes. For example, single strand

- 5 conformation polymorphism (SSCP) may be used to detect differences in electrophoretic mobility between mutant and wild type nucleic acids (Orita *et al.* (1989) *Proc Natl. Acad. Sci USA*: 86:2766, see also Cotton (1993) *Mutat Res* 285:125-144; and Hayashi (1992) *Genet Anal Tech Appl* 9:73-79). Single-stranded DNA fragments of sample and control Dkk or Dkk-related nucleic acids will be denatured and allowed to renature. The
- 10 secondary structure of single-stranded nucleic acids varies according to sequence, the resulting alteration in electrophoretic mobility enables the detection of even a single base change. The DNA fragments may be labeled or detected with labeled probes. The sensitivity of the assay may be enhanced by using RNA (rather than DNA), in which the secondary structure is more sensitive to a change in sequence. In a preferred
- 15 embodiment, the subject method utilizes heteroduplex analysis to separate double stranded heteroduplex molecules on the basis of changes in electrophoretic mobility (Keen *et al.* (1991) *Trends Genet* 7:5).

In yet another embodiment the movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing

20 gradient gel electrophoresis (DGGE) (Myers *et al.* (1985) *Nature* 313:495). When DGGE is used as the method of analysis, DNA will be modified to insure that it does not completely denature, for example by adding a GC clamp of approximately 40 bp of high-melting GC-rich DNA by PCR. In a further embodiment, a temperature gradient is used in place of a denaturing gradient to identify differences in the mobility of control

25 and sample DNA (Rosenbaum and Reissner (1987) *Biophys Chem* 265:12753).

Examples of other techniques for detecting point mutations include, but are not limited to, selective oligonucleotide hybridization, selective amplification, or selective primer extension. For example, oligonucleotide primers may be prepared in which the known mutation is placed centrally and then hybridized to target DNA under conditions

30 which permit hybridization only if a perfect match is found (Saiki *et al.* (1986) *Nature* 324:163); Saiki *et al.* (1989) *Proc. Natl Acad. Sci USA* 86:6230). Such allele specific

oligonucleotides are hybridized to PCR amplified target DNA or a number of different mutations when the oligonucleotides are attached to the hybridizing membrane and hybridized with labeled target DNA.

Alternatively, allele specific amplification technology which depends on selective PCR amplification may be used in conjunction with the instant invention. Oligonucleotides used as primers for specific amplification may carry the mutation of interest in the center of the molecule (so that amplification depends on differential hybridization) (Gibbs *et al.* (1989) *Nucleic Acids Res.* 17:2437-2448) or at the extreme 3' end of one primer where, under appropriate conditions, mismatch can prevent, or reduce polymerase extension (Prossner (1993) *Tibtech* 11:238). In addition it may be desirable to introduce a novel restriction site in the region of the mutation to create cleavage-based detection (Gasparini *et al.* (1992) *Mol. Cell Probes* 6:1). It is anticipated that in certain embodiments amplification may also be performed using Taq ligase for amplification (Barany (1991) *Proc. Natl. Acad. Sci USA* 88:189). In such cases, ligation will occur only if there is a perfect match at the 3' end of the 5' sequence making it possible to detect the presence of a known mutation at a specific site by looking for the presence or absence of amplification.

The methods described herein may be performed, for example, by utilizing pre-packaged diagnostic kits comprising at least one probe nucleic acid or antibody reagent described herein, which may be conveniently used, *e.g.*, in clinical settings to diagnose patients exhibiting symptoms or family history of a disease or illness involving a Dkk gene.

Furthermore, any cell type or tissue in which Dkk or a Dkk-related sequence is expressed may be utilized in the prognostic assays described herein.

3. Monitoring of Effects During Clinical Trials

Monitoring the influence of agents (*e.g.*, drugs, compounds) on the expression or activity of Dkk or Dkk-related molecule (*e.g.*, modulation of cellular signal transduction, regulation of gene transcription in a cell involved in development or differentiation, regulation of cellular proliferation) can be applied not only in basic drug screening, but also in clinical trials. For example, the effectiveness of an agent determined by a

screening assay as described herein to increase Dkk or Dkk-related gene expression, protein levels, or upregulate Dkk or Dkk-related activity, can be monitored in clinical trials of subjects exhibiting decreased Dkk or Dkk-related gene expression, protein levels, or downregulated Dkk or Dkk-related activity. Alternatively, the effectiveness of an agent determined by a screening assay to decrease Dkk or Dkk-related gene expression, protein levels, or downregulate Dkk or Dkk-related activity, can be monitored in clinical trials of subjects exhibiting increased Dkk or Dkk-related gene expression, protein levels, or upregulated Dkk or Dkk-related activity. In such clinical trials, the expression or activity of Dkk or Dkk-related and, preferably, other genes that have been implicated in, for example, a proliferative disorder can be used as a "read out" or markers of the phenotype of a particular cell.

For example, and not by way of limitation, genes, including Dkk and Dkk-related genes, that are modulated in cells by treatment with an agent (*e.g.*, compound, drug or small molecule) which modulates Dkk or Dkk-related activity (*e.g.*, identified in a screening assay as described herein) can be identified. Thus, to study the effect of agents on proliferative disorders, developmental or differentiative disorder, hematopoietic disorder as well disorders characterized by abnormal cell differentiation and/or survival, an abnormal extracellular structure, or an abnormality in a defense mechanism, for example, in a clinical trial, cells can be isolated and RNA prepared and analyzed for the levels of expression of Dkk, Dkk-related, and other genes implicated in the proliferative disorder, developmental or differentiative disorder, hematopoietic disorder as well as disorders characterized by abnormal cell differentiation and/or survival, an abnormal extracellular structure, or an abnormality in a defense mechanism, respectively. The levels of gene expression (*i.e.*, a gene expression pattern) can be quantified by Northern blot analysis or RT-PCR, as described herein, or alternatively by measuring the amount of protein produced, by one of the methods as described herein, or by measuring the levels of activity of Dkk, Dkk-related, or other genes. In this way, the gene expression pattern can serve as a marker, indicative of the physiological response of the cells to the agent. Accordingly, this response state may be determined before, and at various points during treatment of the individual with the agent.

In a preferred embodiment, the present invention provides a method for monitoring the effectiveness of treatment of a subject with an agent (*e.g.*, an agonist, antagonist, peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate identified by the screening assays described herein) comprising the steps of (i) obtaining a pre-administration sample from a subject prior to administration of the agent; (ii) detecting the level of expression of a Dkk or Dkk-related protein, mRNA, or genomic DNA in the preadministration sample; (iii) obtaining one or more post-administration samples from the subject; (iv) detecting the level of expression or activity of the Dkk or Dkk-related protein, mRNA, or genomic DNA in the post-administration samples; (v) comparing the level of expression or activity of the Dkk or Dkk-related protein, mRNA, or genomic DNA in the pre-administration sample with the Dkk or Dkk-related protein, mRNA, or genomic DNA in the post administration sample or samples; and (vi) altering the administration of the agent to the subject accordingly. For example, increased administration of the agent may be desirable to increase the expression or activity of Dkk or Dkk-related nucleic acid or protein to higher levels than detected, *i.e.*, to increase the effectiveness of the agent. Alternatively, decreased administration of the agent may be desirable to decrease expression or activity of Dkk or Dkk-related nucleic acid or protein to lower levels than detected, *i.e.* to decrease the effectiveness of the agent. According to such an embodiment, Dkk or Dkk-related expression or activity may be used as an indicator of the effectiveness of an agent, even in the absence of an observable phenotypic response.

C. Methods of Treatment:

The present invention provides for both prophylactic and therapeutic methods of treating a subject at risk of (or susceptible to) a disorder or having a disorder associated with aberrant Dkk or Dkk-related expression or activity. With regards to both prophylactic and therapeutic methods of treatment, such treatments may be specifically tailored or modified, based on knowledge obtained from the field of pharmacogenomics. "Pharmacogenomics", as used herein, refers to the application of genomics technologies such as gene sequencing, statistical genetics, and gene expression analysis to drugs in clinical development and on the market. More specifically, the term refers the study of

how a patient's genes determine his or her response to a drug (*e.g.*, a patient's "drug response phenotype", or "drug response genotype".) Thus, another aspect of the invention provides methods for tailoring an individual's prophylactic or therapeutic treatment with either the Dkk or Dkk-related molecules of the present invention or Dkk or Dkk-related modulators according to that individual's drug response genotype. Pharmacogenomics allows a clinician or physician to target prophylactic or therapeutic treatments to patients who will most benefit from the treatment and to avoid treatment of patients who will experience toxic drug-related side effects.

1. Prophylactic Methods

In one aspect, the invention provides a method for preventing in a subject, a disease or condition associated with an aberrant Dkk or Dkk-related expression or activity, by administering to the subject an agent which modulates Dkk or Dkk-related expression or at least one Dkk or Dkk-related activity. Subjects at risk for a disease which is caused or contributed to by aberrant Dkk or Dkk-related expression or activity can be identified by, for example, any or a combination of diagnostic or prognostic assays as described herein. Administration of a prophylactic agent can occur prior to the manifestation of symptoms characteristic of the Dkk or Dkk-related aberrancy, such that a disease or disorder is prevented or, alternatively, delayed in its progression. Depending on the type of Dkk or Dkk-related aberrancy, for example, an agonist or antagonist agent can be used for treating the subject. The appropriate agent can be determined based on screening assays described herein. The prophylactic methods of the present invention are further discussed in the following subsections.

2. Therapeutic Methods

Another aspect of the invention pertains to methods of modulating Dkk or Dkk-related expression or activity for therapeutic purposes. The modulatory method of the invention involves contacting a cell with an agent that modulates one or more of the activities of Dkk or Dkk-related protein activity associated with the cell. An agent that modulates Dkk or Dkk-related protein activity can be an agent as described herein, such

as a nucleic acid or a protein, a naturally-occurring target molecule of a Dkk or Dkk-related protein, a peptide, a Dkk or Dkk-related peptidomimetic, or other small molecule. In one embodiment, the agent stimulates one or more Dkk or Dkk-related protein activity. Examples of such stimulatory agents include active Dkk or Dkk-related
5 protein and a nucleic acid molecule encoding Dkk or Dkk-related that has been introduced into the cell. In another embodiment, the agent inhibits one or more Dkk or Dkk-related protein activity. Examples of such inhibitory agents include antisense Dkk or Dkk-related nucleic acid molecules and antibodies. These modulatory methods can be performed *in vitro* (e.g., by culturing the cell with the agent) or, alternatively, *in vivo*
10 (e.g., by administering the agent to a subject). As such, the present invention provides methods of treating an individual afflicted with a disease or disorder characterized by aberrant expression or activity of a Dkk or Dkk-related protein or nucleic acid molecule. The present invention also provides methods of modulating the function, morphology, proliferation, and/or differentiation of cells in the tissues in which a Dkk or Dkk-related
15 protein or nucleic acid molecule is expressed. Alternatively, Dkk or Dkk-related polypeptides, nucleic acids, and modulators thereof, can be used to treat disorders associated with abnormal or aberrant metabolism or function of cells in the tissues in which the Dkk or Dkk-related protein or nucleic acid molecule is expressed.

For example, tissues in which Dkk-3 is expressed include embryonic eye, bone,
20 and cartilage, fetal brain, lung, and kidney, and adult heart (in particular, atrioventricular valves and atrial myocytes), eye (in particular, the integrating bipolar and ganglion cells of the retina, the ciliary body, and lens epithelium), brain (in particular, neurons of the cortex and hippocampus), placenta, lung, and skeletal muscle. Accordingly, Dkk-3 polypeptides, nucleic acids, or modulators thereof, can be used to treat cardiovascular
25 disorders, such as ischemic heart disease (e.g., angina pectoris, myocardial infarction, and chronic ischemic heart disease), hypertensive heart disease, pulmonary heart disease, valvular heart disease (e.g., rheumatic fever and rheumatic heart disease, endocarditis, mitral valve prolapse, and aortic valve stenosis), congenital heart disease (e.g., valvular and vascular obstructive lesions, atrial or ventricular septal defect, and
30 patent ductus arteriosus), or myocardial disease (e.g., myocarditis, congestive cardiomyopathy, and hypertrophic cardiomyopathy).

In another embodiment, Dkk-3 polypeptides, nucleic acids, or modulators thereof, can be used to treat optic disorders such as diseases associated with amaurosis (*e.g.*, a. fugax and a. albuminuric) diseases associated with amblyopia, glaucoma, optic neuropathy (*e.g.*, ischemic neuropathy, optic neuritis, and infiltrative neuropathy),
5 ophthalmia (*e.g.*, o. catarrhal, trachoma, o. neuroparalytic, and conjunctiva), visual disorders resulting from systemic disease or disorders of other tissues (*e.g.*, diabetes mellitus, hyperthyroidism, and vitamin A or riboflavin deficiency), or tumors, neoplasms, and metastases.

In another embodiment, Dkk-3 polypeptides, nucleic acids, or modulators
10 thereof, can be used to treat disorders of the brain, such as cerebral edema, senile dementia of the Alzheimer type, epilepsy, amnesia, hydrocephalus, brain herniations, iatrogenic disease (due to, *e.g.*, infection, toxins, or drugs), inflammations (*e.g.*, bacterial and viral meningitis, encephalitis, and cerebral toxoplasmosis), cerebrovascular diseases (*e.g.*, hypoxia, ischemia, and infarction, intracranial hemorrhage and vascular
15 malformations, and hypertensive encephalopathy), and tumors (*e.g.*, neuroglial tumors, neuronal tumors, tumors of pineal cells, meningeal tumors, primary and secondary lymphomas, intracranial tumors, and medulloblastoma), and to treat injury or trauma to the brain.

In another embodiment, Dkk-3 polypeptides, nucleic acids, or modulators
20 thereof, can be used to treat placental disorders, such as toxemia of pregnancy (*e.g.*, preeclampsia and eclampsia), placentitis, or spontaneous abortion.

In another embodiment, Dkk-3 polypeptides, nucleic acids, or modulators thereof, can be used to treat pulmonary disorders, such as atelectasis, pulmonary congestion or edema, chronic obstructive airway disease (*e.g.*, emphysema, chronic
25 bronchitis, bronchial asthma, and bronchiectasis), diffuse interstitial diseases (*e.g.*, sarcoidosis, pneumoconiosis, hypersensitivity pneumonitis, Goodpasture's syndrome, idiopathic pulmonary hemosiderosis, pulmonary alveolar proteinosis, desquamative interstitial pneumonitis, chronic interstitial pneumonia, fibrosing alveolitis, hamman-rich syndrome, pulmonary eosinophilia, diffuse interstitial fibrosis, Wegener's
30 granulomatosis, lymphomatoid granulomatosis, and lipid pneumonia), or tumors (*e.g.*, bronchogenic carcinoma, bronchioloalveolar carcinoma, bronchial carcinoid,

hamartoma, and mesenchymal tumors).

In another embodiment, Dkk-3 polypeptides, nucleic acids, or modulators thereof, can be used to treat disorders of skeletal muscle, such as muscular atrophy (due to, *e.g.*, denervation, malnutrition, loss of blood supply, or neuromuscular disease, *e.g.*, amyotonia congenita, amyotrophic lateral sclerosis of Charcot, and progressive muscular atrophy of Aran-Duchenne), myositis (due to, *e.g.*, bacterial, viral, fungal or parasitic infection), muscular dystrophies (*e.g.*, Duchenne type, Becker type, facioscapulohumeral, limb-girdle, myotonic dystrophy, and ocular myopathy), myasthenia gravis, or tumors and tumor-like lesions of muscles (*e.g.*, traumatic myositis ossificans, desmoids, musculoaponeurotic fibromatosis, Dupuytren's contracture, nodular (pseudosarcomatous) fasciitis, rhabdomyoma, rhabdomyosarcoma, and granular cell myoblastomas).

Tissues in which Dkk-4 is expressed include cerebellum, activated human T-lymphocytes, lung, and esophagus. Accordingly, in one embodiment, Dkk-4 polypeptides, nucleic acids, or modulators thereof, can be used to treat disorders of the cerebellum, such as disturbances of synergy (*e.g.*, asynergia or limb ataxia, dysmetria, decomposition of movement, hypermetria, hypometria, dysdiadochokinesia, hypotonia, tremor, dysarthria, nystagmus), disturbances of equilibrium (due to, *e.g.*, a lesion involving the vestibulocerebellum), disturbances of gait stance, or tone (due to, *e.g.*, a lesion or degeneration of the spinocerebellum), or tumors (*e.g.*, astrocytoma and medulloblastoma).

In another embodiment, Dkk-4 polypeptides, nucleic acids, or modulators thereof, can be used to treat lymphocytic disorders, such as lymphopenia, lymphocytosis, acute and chronic lymphadenitis, malignant lymphomas (*e.g.*, Non-Hodgkin's lymphomas, Hodgkin's lymphomas, leukemias, multiple myeloma, histiocytoses, and angioimmunoblastic lymphadenopathy).

In another embodiment, Dkk-4 polypeptides, nucleic acids, or modulators thereof, can be used to treat pulmonary disorders, such as atelectasis, pulmonary congestion or edema, chronic obstructive airway disease (*e.g.*, emphysema, chronic bronchitis, bronchial asthma, and bronchiectasis), diffuse interstitial diseases (*e.g.*, sarcoidosis, pneumoconiosis, hypersensitivity pneumonitis, Goodpasture's syndrome,

idiopathic pulmonary hemosiderosis, pulmonary alveolar proteinosis, desquamative interstitial pneumonitis, chronic interstitial pneumonia, fibrosing alveolitis, hamman-rich syndrome, pulmonary eosinophilia, diffuse interstitial fibrosis, Wegener's granulomatosis, lymphomatoid granulomatosis, and lipid pneumonia), or tumors (*e.g.*,
5 bronchogenic carcinoma, bronchioloalveolar carcinoma, bronchial carcinoid, hamartoma, and mesenchymal tumors).

In another embodiment, Dkk-4 polypeptides, nucleic acids, or modulators thereof, can be used to treat esophageal disorders, such as neuromuscular disturbances (*e.g.*, achalasia, annular narrowings, Schatzki's rings, hiatal hernia, Mallory-Weiss
10 syndrome), esophagitis (due to *e.g.*, bacteremia, viremia, fungal infections, uremia, graft-versus-host disease, chemotherapy, radiation, and prolonged gastric intubation), diverticula (*e.g.*, Zenker's diverticulum), systemic sclerosis, varices (due to, *e.g.*, portal hypertension, systemic amyloidosis and sarcoidosis), or tumors or neoplasms (*e.g.*, leiomyoma, fibromas, lipomas, hemangiomas, lymphangiomas, squamous papillomas,
15 adenocarcinomas and undifferentiated carcinomas, and sarcomas).

Dkk-1 is highly expressed, for example, in placenta. Accordingly, Dkk-1 polypeptides, nucleic acids, or modulators thereof, can be used to treat placental disorders, such as toxemia of pregnancy (*e.g.*, preeclampsia and eclampsia), placentitis, or spontaneous abortion.

20 Tissues in which Dkk-2 is expressed include, for example, heart, brain, placenta, lung, and skeletal muscle. Accordingly, Dkk-2 polypeptides, nucleic acids, or modulators thereof, can be used to treat cardiovascular disorders, such as ischemic heart disease (*e.g.*, angina pectoris, myocardial infarction, and chronic ischemic heart disease), hypertensive heart disease, pulmonary heart disease, valvular heart disease (*e.g.*,
25 rheumatic fever and rheumatic heart disease, endocarditis, mitral valve prolapse, and aortic valve stenosis), congenital heart disease (*e.g.*, valvular and vascular obstructive lesions, atrial or ventricular septal defect, and patent ductus arteriosus), or myocardial disease (*e.g.*, myocarditis, congestive cardiomyopathy, and hypertrophic cardiomyopathy).

In another embodiment, Dkk-2 polypeptides, nucleic acids, or modulators thereof, can be used to treat disorders of the brain, such as cerebral edema, senile dementia of the Alzheimer type, epilepsy, amnesia, hydrocephalus, brain herniations, iatrogenic disease (due to, *e.g.*, infection, toxins, or drugs), inflammations (*e.g.*, bacterial
5 and viral meningitis, encephalitis, and cerebral toxoplasmosis), cerebrovascular diseases (*e.g.*, hypoxia, ischemia, and infarction, intracranial hemorrhage and vascular malformations, and hypertensive encephalopathy), and tumors (*e.g.*, neuroglial tumors, neuronal tumors, tumors of pineal cells, meningeal tumors, primary and secondary lymphomas, intracranial tumors, and medulloblastoma), and to treat injury or trauma to
10 the brain.

In another embodiment, Dkk-2 polypeptides, nucleic acids, or modulators thereof, can be used to treat placental disorders, such as toxemia of pregnancy (*e.g.*, preeclampsia and eclampsia), placentitis, or spontaneous abortion.

In another embodiment, Dkk-2 polypeptides, nucleic acids, or modulators
15 thereof, can be used to treat pulmonary disorders, such as atelectasis, pulmonary congestion or edema, chronic obstructive airway disease (*e.g.*, emphysema, chronic bronchitis, bronchial asthma, and bronchiectasis), diffuse interstitial diseases (*e.g.*, sarcoidosis, pneumoconiosis, hypersensitivity pneumonitis, Goodpasture's syndrome, idiopathic pulmonary hemosiderosis, pulmonary alveolar proteinosis, desquamative
20 interstitial pneumonitis, chronic interstitial pneumonia, fibrosing alveolitis, hamman-rich syndrome, pulmonary eosinophilia, diffuse interstitial fibrosis, Wegener's granulomatosis, lymphomatoid granulomatosis, and lipid pneumonia), or tumors (*e.g.*, bronchogenic carcinoma, bronchioloalveolar carcinoma, bronchial carcinoid, hamartoma, and mesenchymal tumors).

25 In another embodiment, Dkk-2 polypeptides, nucleic acids, or modulators thereof, can be used to treat disorders of skeletal muscle, such as muscular atrophy (due to, *e.g.*, denervation, malnutrition, loss of blood supply, or neuromuscular disease, *e.g.*, amyotonia congenita, amyotrophic lateral sclerosis of Charcot, and progressive muscular atrophy of Aran-Duchenne), myositis (due to, *e.g.*, bacterial, viral, fungal or parasitic
30 infection), muscular dystrophies (*e.g.*, Duchenne type, Becker type, facioscapulohumeral, limb-girdle, myotonic dystrophy, and ocular myopathy),

myasthenia gravis, or tumors and tumor-like lesions of muscles (*e.g.*, traumatic myositis ossificans, desmoids, musculoaponeurotic fibromatosis, Dupuytren's contracture, nodular (pseudosarcomatous) fasciitis, rhabdomyoma, rhabdomyosarcoma, and granular cell myoblastomas).

5 Soggy-1 is expressed in, for example, testis (*e.g.*, spermatogenic epithelium of the seminiferous tubules, spermatogonia) and in embryonic developing dorsal root ganglia, cartilage primordium of the nasal septum, and the eye. Accordingly, Soggy-1 polypeptides, nucleic acids, or modulators thereof, can be used to treat testicular disorders, such as unilateral testicular enlargement (*e.g.*, nontuberculous, granulomatous
10 orchitis), inflammatory diseases resulting in testicular dysfunction (*e.g.*, gonorrhea and mumps), and tumors (*e.g.*, germ cell tumors, interstitial cell tumors, androblastoma, testicular lymphoma and adenomatoid tumors). In another embodiment, Soggy-1 polypeptides, nucleic acids, or modulators thereof, can be used to treat infertility due to, for example, spermatogenetic failure.

15 In one aspect, the above-described methods involve administering an agent (*e.g.*, an agent identified by a screening assay described herein), or combination of agents that modulates (*e.g.*, upregulates or downregulates) Dkk or Dkk-related expression or activity. In another embodiment, the method involves administering a Dkk or Dkk-related protein or nucleic acid molecule as therapy to compensate for reduced or aberrant
20 Dkk or Dkk-related expression or activity.

A preferred embodiment of the present invention involves a method for treatment of a disease or disorder associated with a Dkk or Dkk-related protein which includes the step of administering a therapeutically effective amount of an antibody to a Dkk or Dkk-related protein to a subject. As defined herein, a therapeutically effective amount of
25 antibody (*i.e.*, an effective dosage) ranges from about 0.001 to 30 mg/kg body weight, preferably about 0.01 to 25 mg/kg body weight, more preferably about 0.1 to 20 mg/kg body weight, and even more preferably about 1 to 10 mg/kg, 2 to 9 mg/kg, 3 to 8 mg/kg, 4 to 7 mg/kg, or 5 to 6 mg/kg body weight. The skilled artisan will appreciate that certain factors may influence the dosage required to effectively treat a subject, including
30 but not limited to the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and other diseases present. Moreover, treatment of a

- 112 -

subject with a therapeutically effective amount of an antibody can include a single treatment or, preferably, can include a series of treatments. In a preferred example, a subject is treated with antibody in the range of between about 0.1 to 20 mg/kg body weight, one time per week for between about 1 to 10 weeks, preferably between 2 to 8 weeks, more preferably between about 3 to 7 weeks, and even more preferably for about 4, 5, or 6 weeks. It will also be appreciated that the effective dosage of antibody used for treatment may increase or decrease over the course of a particular treatment. Changes in dosage may result from the results of diagnostic assays as described herein.

Stimulation of Dkk or Dkk-related activity is desirable in situations in which Dkk or Dkk-related activity is abnormally downregulated and/or in which increased Dkk or Dkk-related activity is likely to have a beneficial effect. Likewise, inhibition of Dkk or Dkk-related activity is desirable in situations in which Dkk or Dkk-related activity is abnormally upregulated and/or in which decreased Dkk or Dkk-related activity is likely to have a beneficial effect. One example of such a situation is where a subject has a disorder characterized by aberrant development or cellular differentiation. Another example of such a situation is where the subject has a proliferative disease (*e.g.*, cancer) or a neurogenerative disorder. Yet another example of such a situation is where it is desirable to achieve tissue regeneration in a subject (*e.g.*, where a subject has undergone brain or spinal cord injury and it is desirable to regenerate neuronal tissue in a regulated manner.)

Accordingly, in one embodiment, the disease is a disease characterized by an abnormal cell proliferation, differentiation, and/or survival. For example, the disease can be a hyper- or hypoproliferative disease. The invention also provides methods for treating diseases characterized by an abnormal cell proliferation, differentiation, and/or survival in a subject, which are not characterized by an abnormal Dkk or Dkk-related activity (*e.g.*, hDkk-3 activity). In fact, since Dkk is likely to be capable of modulating the proliferative state of a cell (*i.e.*, state of proliferation, differentiation, and or survival of a cell), Dkk can regulate disease wherein the abnormal proliferative state of a cell results from a defect other than an abnormal Dkk activity.

Hyperproliferative diseases can be treated with Dkk or Dkk-related (*e.g.*, hDkk-3) therapeutics include neoplastic and hyperplastic diseases, such as various forms of cancers and leukemias, and fibroproliferative disorders. Other hyperproliferative diseases that can be treated or prevented with the subject Dkk or Dkk-related

5 therapeutics (*e.g.* hDkk-3 therapeutics) include malignant conditions, premalignant conditions, and benign conditions. The condition to be treated or prevented can be a solid tumor, such as a tumor arising in an epithelial tissue. Accordingly, treatment of such a cancer could comprise administration to the subject of a Dkk or Dkk-related therapeutic decreasing the interaction of Dkk with a Dkk receptor. Other cancers that
10 can be treated or prevented with a Dkk or Dkk-related protein include cancers of the epithelia (*e.g.*, carcinomas of the pancreas, kidney, stomach, colon, esophagus liver, secretory glands (*e.g.*, adenocarcinoma) bladder, lung, breast, skin (*e.g.*, malignant melanoma, seminoma squamous adenocarcinoma), reproductive tract including prostate gland, testis, ovary, cervix and uterus); cancers of the hematopoietic and immune system
15 (*e.g.*, leukemias and lymphomas); cancers of the central nervous, brain system and eye (*e.g.*, malignant astrocytoma, gliomas, neuroblastoma and retinoblastoma); and cancers of connective tissues, bone, heart, muscles and vasculature (*e.g.*, sarcomas, for example, osteosarcoma). Additional solid tumors within the scope of the invention include those that can be found in a medical textbook.

20 The condition to be treated or prevented can also be a soluble tumor, such as leukemia, either chronic or acute, including chronic or acute myelogenous leukemia, chronic or acute lymphocytic leukemia, promyelocytic leukemia, monocytic leukemia, myelomonocytic leukemia, and erythroleukemia. Yet other proliferative disorders that can be treated with a Dkk or Dkk-related therapeutic of the invention include heavy
25 chain disease, multiple myeloma, lymphoma, *e.g.*, Hodgkin's lymphoma and non-Hodgkin's lymphoma, and Waldenstroem's macroglobulemia.

Diseases or conditions characterized by a solid or soluble tumor can be treated by administering a Dkk or Dkk-related therapeutic either locally or systemically, such that aberrant cell proliferation is inhibited or decreased. Methods for administering the
30 compounds of the invention are further described below.

The invention also provides methods for preventing the formation and/or development of tumors. For example, the development of a tumor can be preceded by the presence of a specific lesion, such as a pre-neoplastic lesion, *e.g.*, hyperplasia, metaplasia, and dysplasia, which can be detected, *e.g.*, by cytologic methods. Such lesions can be found, *e.g.*, in epithelial tissue. Thus, the invention provides a method for inhibiting progression of such a lesion into a neoplastic lesion, comprising administering to the subject having a preneoplastic lesion an amount of a Dkk or Dkk-related therapeutic sufficient to inhibit progression of the preneoplastic lesion into a neoplastic lesion.

The invention also provides for methods for treating or preventing diseases or conditions in which proliferation of cells is desired. For example, Dkk or Dkk-related therapeutics can be used to stimulate tissue repair or wound healing, such as after surgery or to stimulate tissue healing from burns. Other diseases in which proliferation of cells is desired are hypoproliferative diseases, *i.e.*, diseases characterized by an abnormally low proliferation of certain cells.

In yet another embodiment, the invention provides a method for treating or preventing diseases or conditions characterized by aberrant cell differentiation. Accordingly, the invention provides methods for stimulating cellular differentiation in conditions characterized by an inhibition of normal cell differentiation which may or may not be accompanied by excessive proliferation. Alternatively, Dkk or Dkk-related therapeutics can be used to inhibit differentiation of specific cells.

In a preferred method, the aberrantly proliferating and/or differentiating cell is a cell present in the nervous system. A role for Dkk in the nervous system is suggested at least in part from the fact that human Dkk-3 is expressed in human fetal brain.

Accordingly, the invention provides methods for treating diseases or conditions associated with a central or peripheral nervous system. For example, the invention provides methods for treating lesions of the nervous system associated with an aberrant proliferation, differentiation or survival of any of the following cells: cells of the central nervous system including neurons and glial cells (*e.g.*, astrocytes and oligodendrocytes) and supporting cells of peripheral neurons (*e.g.*, Schwann cells and satellite cells).

Disorders of the nervous system include, but are not limited to: spinal cord injuries,

brain injuries, brain tumors (*e.g.*, astrocytic tumors, for example, astrocytomas and glioblastomas), lesions associated with surgery, ischemic lesions, malignant lesions, infectious lesions, degenerative lesions (*e.g.*, Parkinson's disease, Alzheimer's disease, Huntington's chorea, amyotrophic lateral sclerosis), demyelinating diseases (*e.g.*,
5 multiple sclerosis, human immunodeficiency associated myelopathy, transverse myelopathy, progressive multifocal leukoencephalopathy, pontine myelinolysis), motor neuron injuries, progressive spinal muscular atrophy, progressive bulbar palsy, primary lateral sclerosis, infantile and juvenile muscular atrophy, progressive bulbar paralysis of childhood (*i.e.*, Fazio-Londe syndrome), poliomyelitis, and hereditary motorsensory
10 neuropathy (*i.e.*, Charcot-Marie-Tooth disease).

In another embodiment, the invention provides a method for enhancing the survival and/or stimulating proliferation and/or differentiation of cells and tissues *in vitro*. In a preferred embodiment, Dkk or Dkk-related therapeutics are used to promote tissue regeneration and/or repair (*e.g.*, to treat nerve injury). For example, tissues from a
15 subject can be obtained and grown *in vitro* in the presence of a Dkk or Dkk-related therapeutic, such that the tissue cells are stimulated to proliferate and/or differentiate. The tissue can then be readministered to the subject.

Among the approaches which may be used to ameliorate disease symptoms involving an aberrant Dkk or Dkk-related activity and/or an abnormal cell proliferation, differentiation, and/or survival, are, for example, antisense, ribozyme, and triple helix
20 molecules described above. Examples of suitable compounds include the antagonists, agonists or homologues described in detail above.

Yet other Dkk or Dkk-related therapeutics consist of a first peptide comprising a Dkk or Dkk-related peptide capable of binding to a Dkk receptor, and a second peptide
25 which is cytotoxic. Such therapeutics can be used to specifically target and lyse cells expressing or overexpressing a receptor for Dkk.

3. Pharmacogenomics

The Dkk or Dkk-related molecules of the present invention, as well as agents, or
30 modulators which have a stimulatory or inhibitory effect on Dkk or Dkk-related activity (*e.g.*, Dkk or Dkk-related gene expression) as identified by a screening assay described

herein can be administered to individuals to treat (prophylactically or therapeutically) disorders (*e.g.*, proliferative or developmental disorders) associated with aberrant Dkk or Dkk-related activity. In conjunction with such treatment, pharmacogenomics (*i.e.*, the study of the relationship between an individual's genotype and that individual's response to a foreign compound or drug) may be considered. Differences in metabolism of therapeutics can lead to severe toxicity or therapeutic failure by altering the relation between dose and blood concentration of the pharmacologically active drug. Thus, a physician or clinician may consider applying knowledge obtained in relevant pharmacogenomics studies in determining whether to administer a Dkk or Dkk-related molecule or Dkk or Dkk-related modulator as well as tailoring the dosage and/or therapeutic regimen of treatment with a Dkk or Dkk-related molecule or Dkk or Dkk-related modulator.

Pharmacogenomics deals with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See *e.g.*, Eichelbaum, M., *Clin Exp Pharmacol Physiol*, 1996, 23(10-11) :983-985 and Linder, M.W., *Clin Chem*, 1997, 43(2):254-266. In general, two types of pharmacogenetic conditions can be differentiated. Genetic conditions transmitted as a single factor altering the way drugs act on the body (altered drug action) or genetic conditions transmitted as single factors altering the way the body acts on drugs (altered drug metabolism). These pharmacogenetic conditions can occur either as rare genetic defects or as naturally-occurring polymorphisms. For example, glucose-6-phosphate dehydrogenase deficiency (G6PD) is a common inherited enzymopathy in which the main clinical complication is haemolysis after ingestion of oxidant drugs (anti-malarials, sulfonamides, analgesics, nitrofurans) and consumption of fava beans.

One pharmacogenomics approach to identifying genes that predict drug response, known as "a genome-wide association", relies primarily on a high-resolution map of the human genome consisting of already known gene-related markers (*e.g.*, a "bi-allelic" gene marker map which consists of 60,000-100,000 polymorphic or variable sites on the human genome, each of which has two variants.) Such a high-resolution genetic map can be compared to a map of the genome of each of a statistically significant number of patients taking part in a Phase II/III drug trial to identify markers

associated with a particular observed drug response or side effect. Alternatively, such a high resolution map can be generated from a combination of some ten-million known single nucleotide polymorphisms (SNPs) in the human genome. As used herein, a "SNP" is a common alteration that occurs in a single nucleotide base in a stretch of
5 DNA. For example, a SNP may occur once per every 1000 bases of DNA. A SNP may be involved in a disease process, however, the vast majority may not be disease-associated. Given a genetic map based on the occurrence of such SNPs, individuals can be grouped into genetic categories depending on a particular pattern of SNPs in their individual genome. In such a manner, treatment regimens can be tailored to groups of
10 genetically similar individuals, taking into account traits that may be common among such genetically similar individuals.

Alternatively, a method termed the "candidate gene approach", can be utilized to identify genes that predict drug response. According to this method, if a gene that encodes a drug's target is known (*e.g.*, a Dkk protein or Dkk receptor of the present
15 invention), all common variants of that gene can be fairly easily identified in the population and it can be determined if having one version of the gene versus another is associated with a particular drug response.

As an illustrative embodiment, the activity of drug metabolizing enzymes is a major determinant of both the intensity and duration of drug action. The discovery of
20 genetic polymorphisms of drug metabolizing enzymes (*e.g.*, N-acetyltransferase 2 (NAT 2) and cytochrome P450 enzymes CYP2D6 and CYP2C19) has provided an explanation as to why some patients do not obtain the expected drug effects or show exaggerated drug response and serious toxicity after taking the standard and safe dose of a drug. These polymorphisms are expressed in two phenotypes in the population, the extensive
25 metabolizer (EM) and poor metabolizer (PM). The prevalence of PM is different among different populations. For example, the gene coding for CYP2D6 is highly polymorphic and several mutations have been identified in PM, which all lead to the absence of functional CYP2D6. Poor metabolizers of CYP2D6 and CYP2C19 quite frequently experience exaggerated drug response and side effects when they receive standard doses.
30 If a metabolite is the active therapeutic moiety, PM show no therapeutic response, as demonstrated for the analgesic effect of codeine mediated by its CYP2D6-formed

- 118 -

metabolite morphine. The other extreme are the so called ultra-rapid metabolizers who do not respond to standard doses. Recently, the molecular basis of ultra-rapid metabolism has been identified to be due to CYP2D6 gene amplification.

Alternatively, a method termed the "gene expression profiling", can be utilized to
5 identify genes that predict drug response. For example, the gene expression of an animal dosed with a drug (*e.g.*, a Dkk molecule or Dkk modulator of the present invention) can give an indication whether gene pathways related to toxicity have been turned on.

Information generated from more than one of the above pharmacogenomics
10 approaches can be used to determine appropriate dosage and treatment regimens for prophylactic or therapeutic treatment an individual. This knowledge, when applied to dosing or drug selection, can avoid adverse reactions or therapeutic failure and thus enhance therapeutic or prophylactic efficiency when treating a subject with a Dkk molecule or Dkk or Dkk-related modulator, such as a modulator identified by one of the
15 exemplary screening assays described herein.

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application are hereby incorporated by reference.

20

EXAMPLES

The invention is based, at least in part, on the discovery of a family of genes encoding human cysteine-rich secreted proteins which are related to *Xenopus* Dickkopf (Dkk) proteins. This family includes hDkk-1, hDkk-2, hDkk-3, and hDkk-4. hDkks 1-4
25 contain two highly conserved cysteine-rich domains (CRDs), the most C-terminal of which demonstrates similarity to the colipase protein family. The invention is based also in part on the discovery of a family of Dkk-related proteins, referred to as Soggy proteins, as well as the genes encoding Soggy proteins. Soggy-1 is a novel secreted protein which is related to the N-terminal region of Dkk-3 but lacks CRDs. The
30 following examples illustrate the structure and function of each of these novel human secreted proteins.

Example 1: Isolation And Characterization of Human hDkk-3 cDNA

In this example, the isolation and characterization of the gene encoding human Dkk-3 (also referred to as "hDkk-3", "Cysteine Rich Secreted Protein-1", "CRSP-1"
5 "CRISPY-1" or "TANGO 59") is described.

Isolation of a Human Dkk-3 cDNA

The invention is based at least in part on the discovery of a human gene encoding a secreted protein, referred to herein as human Dickkopf-3 (hDkk-3). A partial cDNA
10 was isolated using a Signal Sequence Trap method. This methodology takes advantage of the fact that molecules such as Dkk have an amino terminal signal sequence which directs certain secreted and membrane-bound proteins through the cellular secretory apparatus.

Briefly, a randomly primed cDNA library using mRNA prepared from human
15 fetal brain tissue (Clontech, Palo Alto CA) was made by using the Stratagene-ZAP-cDNA Synthesis™ kit, (catalog #20041). The cDNA was ligated into the mammalian expression vector pTrap adjacent to a cDNA encoding placental alkaline phosphatase lacking a secretory signal. The plasmids were transformed into *E. coli* and DNA was prepared using the Wizard™ DNA purification kit (Promega). DNA was transfected into
20 COS-7 cells with lipofectamine™ (Gibco-BRL). After 48 hours incubation the COS cell supernatants were assayed for alkaline phosphatase on a Wallac Micro-Beta scintillation counter using the Phospha-Light™ kit (Tropix Inc. Catalog #BP300). The individual plasmid DNAs scoring positive in the COS cell Alkaline Phosphatase secretion assay were further analyzed by DNA sequencing using standard procedures.

25 Using a partial cDNA isolated by the above-described method (clone Amhb3c2), a full length cDNA encoding human Dkk-3 was isolated from a lambda Ziplox™ human fetal brain cDNA library using conventional hybridization techniques (Sambrook *et al.*, *supra*). The nucleotide sequence encoding the full length human Dkk-3 protein is shown in Figure 1 and is set forth as SEQ ID NO: 1. The full length protein encoded by
30 this nucleic acid is comprised of about 350 amino acids and has the amino acid sequence

- 120 -

shown in Figure 1 and set forth as SEQ ID NO:2. The coding portion (open reading frame) of SEQ ID NO:1 is set forth as SEQ ID NO:3. DNA for the clone Fmhb059 was deposited with the ATCC as Accession No. 98452.

5 Analysis of Human hDkk-3

Determination of the hydrophobicity profile of human Dkk-3 having the amino acid sequence set forth in SEQ ID NO:2 indicated the presence of a hydrophobic region from about amino acid 1 to about amino acid 22 of SEQ ID NO:2. Further analysis of the amino acid sequence SEQ ID NO:2 using a signal peptide prediction program
10 predicted the presence of a signal peptide from about amino acid 1 to about amino acid 22 of SEQ ID NO:2. Accordingly, the mature hDkk-3 protein includes about 328 amino acids spanning from about amino acid 23 to about amino acid 350 of SEQ ID NO:2. The presence of the signal sequence, in addition to the fact that hDkk-3 has been identified using a Signal Sequence Trap system, indicates that hDkk-3 is a secreted
15 protein. Furthermore, the prediction of such a signal peptide and signal peptide cleavage site can be made, for example, utilizing the computer algorithm SIGNALP (Nielsen, *et al.*, (1997) *Protein Engineering* 10:1-6).

Examination of the cDNA sequence depicted in Figure 1 shows that human Dkk-3 is particularly rich in cysteine residues. As shown in Figure 1, hDkk-3 contains 20
20 cysteine residues located between amino acid 147 and amino acid 284 of SEQ ID NO: 2. This region has been termed the cysteine-rich region. These cysteine residues can form 10 disulfide bridges.

A BLAST search (Altschul *et al.*, (1990) *J. Mol. Biol.* 215:403) of the nucleotide and the amino acid sequences of hDkk-3 has revealed that hDkk-3 is similar to a chicken
25 cDNA encoding a protein of unknown function having GenBank Accession No. D26311. This cDNA was isolated from a chicken lens cDNA library and was shown to be expressed in lens fibers and lens epithelium, but not in neural retina nor in liver cells. (Sawada *et al.*, (1996) *Int. J. Dev. Biol.* 40:531). hDkk-3 and the chicken protein have 56% amino acid sequence identity and 72% amino acid sequence similarity. The amino
30 acid sequence similarity between the chicken protein and human Dkk-3 is particularly high in the cysteine-rich domain of hDkk-3 which is located between amino acids 147

- 121 -

and 284 of SEQ ID NO: 2. In particular, the 20 cysteine residues of hDkk-3 located in this region are present in the chicken protein.

Two genes recently identified in a screen for suppressors of glioblastoma formation (Ligon *et al.* (1997) *Oncogene* 14:1075-1081) also show homology to hDkk-3. These genes, RIG ("Regulated In Glioblastoma") and RIG-like 7-1 (GenBank Accession Nos. U32331 and AF034208, respectively) were identified in a differential screen for mRNAs regulated by the introduction of a normal copy of chromosome 10 into a glioblastoma cell line harboring a deletion in chromosome 10 that promotes tumorigenesis. A schematic diagram summarizing the relationship between the sequences of the hDkk-3 and the RIG genes is presented as Figure 12. The indicated region of identity between hDkk-3 and RIG comprises a short portion of the 3' UTR of the human Dkk-3 mRNA (*e.g.*, RIG mRNA is ~100% identical to residues 2479 to 2153 of SEQ ID NO:1). RIG-like 7-1 is homologous to hDkk-3 across a longer region (*e.g.*, 97% identical from about nucleotides 316 to 2438 of SEQ ID NO:1) although the encoded RIG-like 7-1 protein lacks the Dkk N-terminal signal sequence and is not therefore predicted to be a secreted protein. These data associate hDkk-3 with human glioblastoma and suggest that hDkk-3 may be important in the suppression of the tumorigenic phenotype. A role in glioblastoma is also consistent with the high level of hDkk-3 mRNA expression observed in human brain tissue. In addition, the co-localization of the hDkk-3, RIG and RIG-like genes to a region of chromosome 11 (11p15.1) implicated in the development of human malignant astrocytoma (Ligon *et al.*, *supra*) further indicates a role for these genes in tumorigenesis.

Human hDkk-3 protein has also some amino acid sequence similarity to metallothionein, particularly in the cyteine-rich domain.

Tissue Distribution of hDkk-3 mRNA

For Northern blots, all hybridizations were to Clontech Multiple Tissue Northern Blots and were performed in ExpressHyb solution (Clontech) for 1-20 hours. All probes were prepared by random primed radiolabelling (Prime-It, Stratagene). Blots were washed sequentially to a final stringency of 0.2x SSC/0.2% SDS and exposed to autoradiographic film. Hybridizations of a control β -actin cDNA probe consistently

- 122 -

demonstrated even loading of the Northern blots. The results of hybridization of the probe to various mRNA samples are described below.

Hybridization of a Clontech Fetal Multiple Tissue Northern (MTN) blot (Clontech, LaJolla, CA) containing RNA from fetal brain, lung, liver, and kidney

5 indicated the presence of high levels of hDkk-3 mRNA (~2.5kb) in fetal brain, lung, and slightly lower levels of hDkk-3 mRNA in fetal kidney. However, no significant level of hDkk-3 mRNA was found in fetal liver.

Hybridization of a Clontech human Multiple Tissue Northern (MTN) blot (Clontech, LaJolla, CA) containing RNA from adult heart, brain, placenta, lung, liver,

10 skeletal muscle, kidney, and pancreas with a human Dkk-3 probe indicated the presence of high levels of hDkk-3 mRNA in heart, slightly lower levels in brain, and much lower levels in placenta and lung. Some hDkk-3 mRNA was also found in adult skeletal muscle. However, no significant levels of hDkk-3 mRNA was observed in adult liver, kidney, or pancreas. Interestingly, the chicken gene which is homologous to hDkk-3
15 was not expressed at detectable levels in liver either (Sawada *et al.*, (1996) *Int. J. Dev. Biol.* 40:531).

Further hybridization of a Clontech human Multiple Tissue Northern (MTN) blot (Clontech, LaJolla, CA) including RNA from bone marrow, adrenal gland, trachea,

lymph node, spinal cord, thyroid, and stomach revealed high levels of expression of
20 hDkk-3 in mRNA isolated from adult spinal cord, and lower level expression in adrenal gland, trachea, thyroid, and stomach.

Thus, hDkk-3 is expressed in a tissue specific manner, with the strongest expression observed in brain, heart, and spinal cord.

25 **Example 2: Isolation And Characterization of mDkk-3 cDNA**

In this example, the isolation and characterization of the gene encoding murine Dkk-3 (also referred to as "mDkk-3", "murine Cysteine Rich Secreted Protein-1", "murine CRSP-1" or "murine CRISPY-1") is described.

Identification of a Murine Dkk-3 cDNA

A full length mDkk-3 cDNA was identified by comparison of the hDkk-3 sequence to a proprietary EST Database using the BLAST-X algorithm. A single clone identified in a adult mouse brain cDNA library was obtained and sequenced fully. DNA
5 for the clone Fmmb059s was deposited with the ATCC as Accession No. 98634. mDkk-3 is predicted to have a signal peptide from residues 1 to 23 of SEQ ID NO:17, cleavage of which results in a mature protein having 326 amino acids in length corresponding to amino acids 24 to 349 of SEQ ID NO:17.

10 Tissue Distribution of mDkk-3 mRNA

To determine the expression pattern of mDkk-3, *in situ* hybridization was performed as follows. Normal mouse embryos and adult mouse tissues were collected from C57BL/6 mice, embedded in TissueTek™ O.C.T Compound (Sakura Finetek U.S.A., Inc., Torrance, CA), frozen on dry ice, and stored at -80°C. Cryostat serial
15 sections (8µm) were thaw mounted on Superfrost Plus™ slides (VWR Scientific, West Chester, PA.) and air dried on a slide warmer at 40°C for 20 minutes. Sections were then fixed with 4% formaldehyde in DEPC treated 0.1 M phosphate-buffered saline (PBS, pH 7.5) at room temperature for 10 minutes and rinsed twice in DEPC-PBS. Sections were rinsed in 0.1 M triethanolamine-HCl (TEA, pH 8.0), incubated in 0.25%
20 acetic anhydride-TEA for 10 minutes and rinsed in DEPC-2X SSC (standard sodium citrate). Sections were dehydrated through a series of graded ethanols, incubated in 100% chloroform for 5 minutes, rinsed in 100% and 95% ethanol for 1 minutes and air dried.

Antisense and sense RNA transcripts were prepared by *in vitro* transcription
25 (Riboprobe Gemini System™, Promega) of PCR amplified cDNA templates. Template amplification primers were as follows;

mDkk-3 forward 5'-CAGTGAGTGCTGTGGAGACC-3' (SEQ ID NO:30), and

reverse 5'-TCTTCAGTCAGGCTCCTCTC-3' (SEQ ID NO:31).

Probes were labeled with ³⁵S-UTP (NEN) and purified on G-25 spin columns
30 (Pharmacia). The hybridization cocktail contained: 50% formamide, 10% dextran

- 124 -

sulfate, 0.1% sodium dodecyl sulfate (SDS), 0.1% sodium thiosulfate, 1X Denhardt's solution, 0.6 M NaCl, 10 mM Tris (pH 7.5), 1 mM EDTA, 100 mM dithiothreitol (DTT), 0.1 mg/ml sheared salmon sperm, 50 µg/ml yeast tRNA, 0.5 mg/ml yeast total RNA, and ³⁵S-UTP labeled probe at a concentration of 5 X 10⁷ c.p.m./100µl of final hybridization solution; 100 µl of hybridization solution was put on each section. The sections were then covered with a glass coverslip and incubated in a humidified chamber at 55°C for 18 h. After hybridization, slides were washed with 2 X SSC. Sections were then sequentially incubated at 37°C in TNE (a solution containing 10 mM Tris-HCl (pH 7.6), 500 mM NaCl, and 1 mM EDTA), for 10 minutes, in TNE with 10µg/ml RNase A for 30 minutes, and finally in TNE for 10 minutes. Slides were then rinsed with 2 X SSC at room temperature, washed in 2 X SSC at 50°C for 1 h, 0.2 X SSC at 55°C for 1h, and 0.2 X SSC at 60°C for 1 h. Sections were dehydrated with a series of graded concentrations of ethanol 0.3 M ammonium acetate, air dried and exposed to Kodak Biomax MR™ scientific imaging film for 6 days at room temperature.

mDkk-3 expression in the brain was found to be highly localized to the cortex and hippocampus but was not observed in the dentate gyrus. Higher power magnification confirmed the mDkk-3 mRNA was localized to neurons within these structures. In the adult eye, mDkk-3 mRNA was found to be highly expressed in the retina, ciliary body, and lens epithelium. Expression in the retina was localized to the integrating bipolar and ganglion cells. In adult heart, mDkk-3 was detected in the atrioventricular valves and also in myocytes of the atria. Expression was highly restricted to the atria and noticeably absent from ventricular tissue. High level expression of mDkk3 mRNA was also observed in developing eye, bone and cartilage in day 14 embryos. These findings corroborate and extend the northern analysis of hDkk-3 mRNA expression in human tissues and also suggest that Dkk-3 may play a role in bone and ocular physiology in addition to functions in neural and cardiac tissues.

Example 3: Secretion and Post-Translational Modification of Dkk-3

This example describes the secretion and post-translational modification (*e.g.*, glycosylation and processing) of hDkk-3 as well as methods for small and large scale purification of hDkk-3.

5

hDkk-3 Expression Constructs

Expression constructs for two forms of hDkk-3 were prepared using the mammalian expression vector pMET-stop. Form-1 comprised a cDNA incorporating the complete 350aa hDkk-3 protein coding sequence (hDkk-3flag.long) and form-2
10 comprised the entire hDkk-3 protein coding sequence except for the final 18 amino acids (hDkk-3flag.short). A C-terminal sequence encoding the FLAG epitope (DYKDDDDK) (SEQ ID NO:19) was added to both hDkk-3 forms for ease of detection and purification. hDkk-3flag cDNAs were generated by PCR from a full length hDkk-3 cDNA template and ligated into pMET-stop using EcoR1 and Sal1 restriction sites.

15

Trial Transfection - Small Scale Expression

Expression constructs for hDkk-3flag.long and hDkk-3flag.short were transfected into 293T cells using 10 µl of lipofectamine (GIBCO/BRL) and 2 µg of DNA per well of a 6-well plate of cells which were 70-80% confluent. After 5 hours at
20 37°C, cells were fed with 1ml of 20%FCS/DMEM. After incubation overnight at 37°C, cells were conditioned in 1ml OptiMEM for 48 hours at 37°C. Samples of supernatant and cell pellets were solubilized in boiling SDS-PAGE gel buffer, run out on a 4-20% SDS-PAGE gel, transferred to a nylon membrane and probed with the anti-FLAG monoclonal antibody M2. Samples from both supernatant and pellet samples showed
25 significant immunoreactivity within a molecular weight range of 40-65 kDa on autoradiographic film using a HRP conjugated secondary antibody and ECL detection reagents. Thus, both forms of hDkk-3 tested are secreted from 293T cells thereby confirming experimentally that hDkk-3 is a secreted protein. It should be noted that the molecular weights of both forms of hDkk-3 tested are greater than predicted from the
30 amino acid sequence, suggesting that the hDkk-3 proteins secreted by 293T cells may be glycosylated. This is consistent with the presence of four potential sites for N-linked

- 126 -

glycosylation in the hDkk-3 protein (*e.g.*, at about amino acids 96-99, 106-109, 121-124, and 204-207 of SEQ ID NO:2).

Deglycosylation of hDkk-3

5 Given the heterogenous nature of secreted human Dkk-3, the effect of N-Glycanase treatment on the mobility of secreted flag-tagged hDkk-3 was studied. Briefly, 1mL samples of 293T cell supernatants collected 72 hours after transfection with the appropriate constructs were incubated with 50μL anti-flag M2 agarose beads (Sigma) for 16hrs at 4°C. Beads were washed with PBS (pH7.4) containing,
10 sequentially, 0.1%, 0.05% and 0.01% Triton X-100. The beads were resuspended in 20μL of 20mM sodium phosphate, pH 7.5, 50mM EDTA, 0.02% sodium azide, (incubation buffer) together with 0.5% SDS, 5% 2-mercaptoethanol and boiled for 2 minutes. The supernatant was split into equal 10μL aliquots which were diluted with 10μL incubation
15 enzyme buffer (20mM Tris-HCl, 1mM EDTA, 50mM NaCl, 0.02% sodium azide pH 7.5) or with enzyme buffer alone as control. After 18 hours at 37°C, samples were boiled in equal volumes of SDS-PAGE buffer and analyzed by SDS-PAGE and Western blotting. For western analysis, samples were electroblotted onto PVDF (Novex) after SDS-PAGE on 4-20% gradient gels, probed with M2 anti-flag antibody (1:500, Sigma)
20 followed by HRP conjugated sheep anti-mouse IgG (1:5000, Amersham), developed with chemiluminescent reagents (Renaissance, Dupont) and exposed to autoradiography film (Biomax MR2 film, Kodak).

Utilizing the above-described methodology, it was determined that hDkk-3 protein displayed a significant increase in mobility following N-Glycanase treatment.
25 The major 45-65 kD form of soluble hDkk-3 was observed as two species of 45-55 and 40 kD following deglycosylation. This finding is consistent with the presence of multiple potential sites of N-linked glycosylation in the hDkk-3 protein. The reason for the heterogeneity of deglycosylated hDkk-3 reflects either proteolytic processing or incomplete removal of carbohydrate from one or more attachment sites. A 30 kD hDkk3
30 species was also observed in these experiments, the mobility of which was unaltered by

N-Glycanase treatment. This form was only observed after overnight incubation of the samples and may be a non-specific degradation product.

Large Scale hDkk-3 Protein Production

5 For scale-up of hDkk-3flag.long protein expression, 30 x 150mm plates of 293T cells at 70-80% confluence were transfected with 27 µg DNA, 100 µl lipofectamine in 18ml OptiMEM for 5 hours at 37°C. 18 ml of 10%FCS/DMEM was added to each plate and incubated overnight at 37°C. 24 hours after the start of transfection, transfection supernatant was aspirated and 35 mls OptiMEM was added to each plate and the plates
10 incubated at 37°C for 72 hours. Conditioned medium was harvested, spun at 4000 rpm for 30 min. at 4°C, and filtered through a 0.45 micron filter unit. 1100 ml was passed over a 1.6 x 10 cm anti-FLAG M2 affinity column pre-equilibrated in PBS pH7.4 buffer at a flow rate of 2.0 ml per minute. After washing with 200 ml of PBS pH 7.4 buffer, bound material was eluted by a step of 200 mM Glycine pH 3.0 buffer and 0.5 ml
15 fractions collected. Upon elution, a significant protein peak was detected by absorbance at 280nm. Samples corresponding to conditioned medium, flow through and eluted fractions were analyzed by Coomassie blue and silver stained SDS-PAGE and by western blot analysis as described above. Significant immunoreactivity within a molecular weight range of 40-65 kDa was detected in conditioned medium and eluted
20 fractions but not in the flow through sample, indicating that the secreted hDkk-3flag.long protein bound to the affinity column specifically and was eluted efficiently by the described conditions. Coomassie blue staining of SDS-PAGE gels suggested that the predominant immunoreactive protein constituted >90% of the protein present in the bound and eluted protein peak. Peak fractions of eluted protein were pooled and
25 dialysed against Phosphate Buffered Saline resulting in a 4 ml volume of recombinant hDkk-3flag.long protein at a concentration of approximately 1mg/ml.

Example 4: Isolation and Characterization of hDkk-4

30 In this example, the isolation and characterization of the gene encoding human Dkk-4 (also referred to as "hDkk-4", "Cysteine Rich Secreted Protein-2", "CRSP-2" or "CRISPY-2") is described.

Isolation and Analysis of a Human Dkk-4 cDNA

To identify novel proteins related to hDkk-3, the human Dkk-3 amino acid sequence was used to search the dbEST database using TBLASTN (WashUversion, 2.0, BLOSUM62 search matrix). A dbEST clone with accession number AA565546 was identified as having homology to a portion of the hDkk-3 cDNA. This clone was obtained from the IMAGE consortium and sequenced fully to define the entire hDkk-4 sequence depicted in Figure 2.

Determination of the hydrophobicity profile of human Dkk-4 having the amino acid sequence set forth in SEQ ID NO:5 indicated the presence of a hydrophobic region from about amino acid 1 to about amino acid 19 of SEQ ID NO:5. Further analysis of the amino acid sequence SEQ ID NO:5 using a signal peptide prediction program predicted the presence of a signal peptide from about amino acid 1 to about amino acid 19 of SEQ ID NO:5. Accordingly, the mature hDkk-4 protein includes about 205 amino acids spanning from about amino acid 20 to about amino acid 224 of SEQ ID NO:5.

Tissue Distribution of hDkk-4

hDkk-4 mRNA was undetectable by Northern analysis in all adult and fetal human tissues examined. Accordingly, a survey was performed of a cDNA library panel by PCR with hDkk-4 specific PCR primers. Using such primers, products were identified in libraries prepared from cerebellum, activated human T-lymphocytes, lung and esophagus.

Secretion and Post-Translational Modification of human Dkk-4

Flag epitope-tagged human Dkk-4 protein was transiently overexpressed in 293T cells and analyzed as described previously for hDkk-3. Soluble hDkk-4 was consistently detected as three major immunoreactive species of approximately 40 kD [form (i)], 30-32 kD [form (ii)] and 15-17 kD [form (iii)]. Neither form (i), (ii) or (iii) was significantly affected by N-glycanase treatment, consistent with the absence of N-glycosylation sites from the protein.

To determine the possible cause of heterogeneity in the size of secreted hDkk-4, Edman N-terminal sequencing of anti-flag affinity purified material corresponding to bands (i), (ii) and (iii) was performed. Briefly, flag-tagged Dkk-4 protein was isolated by passing the conditioned media over an M2-biotin (Sigma)/streptavidin Poros column (2.1 X 30 mm, PE Biosystems); the column was then washed with PBS, pH 7.4 and flag-tagged protein eluted with 200 mM glycine, pH 3.0. Eluted fractions with 280nm absorbance greater than background were analyzed by SDS-PAGE and western blot. Purified Dkk-4 protein bound to PVDF membrane after SDS-PAGE and electroblotting was sequenced for N-terminal amino acid analysis on a PE Applied Biosystems Model 494 Procise instrument using Edman-based chemistry protein sequencing. The amino acid residues were analyzed by HPLC (Spherogel micro PTH 3-micron column) and determined by separation and peak height as compared to standards.

The N-terminal sequence of band (i) was found to be XVLDFNNIRS (SEQ ID NO:34) which corresponds exactly to the predicted signal peptide cleavage site (between Ala-18 and Leu-19). Because the same band is identified by anti-flag antibodies, which recognize the C-terminal epitope tag, band (i) was thus identified as the full length, mature hDkk-4 protein. The band (iii) N-terminal sequence was found to be SQGRKGQEGS (SEQ ID NO:35) which corresponds to CRD-2 cleaved at the dibasic site Lys132/Lys133 (*e.g.*, Lys113/Lys114 of the mature protein following cleavage of the a 19 amino acid signal sequence or Lys 114/Lys 115 following cleavage of a 18 amino acid signal sequence). These data obtained for bands (i) and (iii) indicate clearly that hDkk4 is proteolytically processed by 293T cells, resulting in the release of CRD-2 (a 91 amino acid biologically-active fragment) from the full length protein.

Moreover, the three major species migrated similarly on SDS-PAGE conducted under either reducing or non-reducing conditions. Thus, each of the major C-terminal (anti-flag immunoreactive) hDkk-4 species exist as independent proteolytic fragments that are not covalently linked via disulfide bonds to other subunits or complex components when secreted from 293T cells.

Example 5: Isolation and Characterization of hDkk-1

In this example, the isolation and characterization of the gene encoding human Dkk-1 (also referred to as "hDkk-1", "Cysteine Rich Secreted Protein-3", "CRSP-3" or "CRISPY-3") is described.

5

Identification of a Human Dkk-1 cDNA

Searching a proprietary database of EST information using the sequence of hDkk-3, an hDkk-1 partial sequence was found corresponding to a clone from a human fetal kidney cDNA library having the identification code jthKb075a10. This clone was
10 sequenced further and to define the entire hDkk-1 sequence depicted in Figure 3. DNA for the clone jthKb075a10 was deposited with the ATCC as Accession No. 98633. hDkk-1 has a predicted signal peptide from about amino acid residue 1 to 20 of SEQ ID NO:8, cleavage of which results in a mature protein having 246 amino acid residues in length and corresponding to amino acid residues 21 to 266 of SEQ ID NO:8.

15

Tissue Distribution of hDkk-1

Northern blot analysis of various tissues including heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas was performed as previously described using a probe specific for hDkk-1. A ~1.8 kb hDkk1 mRNA was detected at high levels
20 in human placenta, but not in other tissues tested.

Secretion and Post-Translational Modification of hDkk-1

Flag epitope-tagged human Dkk-1 protein was transiently overexpressed in 293T cells and analyzed as described previously. hDkk-1 was efficiently secreted from
25 mammalian cells and was readily detected in conditioned medium of transfected cells. Mature secreted hDkk-1 migrated with a molecular weight of approximately 42-50 kD. Treatment with N-Glycanase had no significant effect on the mobility of soluble hDkk-1. Although hDkk-1 contains one potential site of N-linked glycosylation at its extreme C-terminus (*e.g.*, at amino acids 256-259 of SEQ ID NO8), this site is not conserved in
30 *Xenopus* Dkk-1 (Glinka *et al.*, *supra*) and appears not to be a major site of carbohydrate addition in 293T cells.

Example 6: Isolation and Characterization of hDkk-2

In this example, the isolation and characterization of the gene encoding human Dkk-2 (also referred to as "hDkk-2", "Cysteine Rich Secreted Protein-4", "CRSP-4" or "CRISPY-4") is described.

5

Isolation of a Human Dkk-2 cDNA

Using the hDkk-3 sequence to query the dbEST database, a clone having similarity to a portion of hDkk-3 was identified having Accession No. W55979. This clone was subsequently obtained from the IMAGE consortium and sequenced to define a partial hDkk-2 sequence set forth as SEQ ID NO:10. This cDNA comprises a coding region from nucleotides 1-537, as well as 3' untranslated sequences (nucleotides 538 to 702). The coding region alone is set forth as SEQ ID NO:12. The predicted amino acid sequence corresponds to amino acids 1 to 179 of SEQ ID NO:11. A cDNA encoding full length hDkk-2 was isolated from a human fetal lung lambda Ziplox libraries by conventional plaque hybridization (Sambrook *et al.*, 1989) and fully sequenced. The full-length nucleotide sequence is set forth as SEQ ID NO:20 and the predicted amino acid sequence is set forth as SEQ ID NO:21. The coding region alone is set forth as SEQ ID NO:22. The predicted amino acid sequence corresponds to amino acids 1 to 259 of SEQ ID NO:21. DNA for the clone fthu133 was deposited with the ATCC as Accession No. _____. hDkk-2 has a predicted signal peptide from about amino acid residue 1 to 33 of SEQ ID NO:21, cleavage of which results in a mature protein having 226 amino acid residues in length and corresponding to amino acid residues 34 to 259 of SEQ ID NO:21.

25 **Tissue Distribution of hDkk-2**

Northern blot analysis of various tissues (*e.g.*, heart, brain, skeletal muscle, colon, thymus, spleen, kidney, liver, small intestine, placenta, lung, and peripheral blood leukocytes) was performed as previously described using a probe specific for hDkk-2. Of the tissues tested, hDkk-2 mRNA expression was highest in heart, brain, placenta, lung, and skeletal muscle. hDkk-2 transcripts of approximately 4.0 and 4.5 kb were observed.

30

Secretion and Post-translational Modification of hDkk-2

Flag epitope-tagged human Dkk-2 protein was transiently overexpressed in 293T cells and analyzed as described previously. Soluble hDkk-2 was detected as a major species of 15-17 kD, closely similar in size to form (iii) of hDkk-4. Additional minor forms of hDkk-2 were also observed in certain experiments in the range of 20-21kD. Deglycosylation of hDkk-2 was not studied since the protein sequence lacks potential N-glycosylation sites. By comparison with the data presented in Example 4 regarding the dibasic proteolytic cleavage site in the hDkk-4 protein sequences, it is predicted that the major 15-17 kD form of hDkk-2 detected in these experiments corresponds to CRD-2, as was the case for hDkk-4.

Example 7: Isolation of Soggy proteins

In this example, the isolation and characterization of the gene encoding human and murine Soggy-1 (also referred to as "Cysteine Rich Secreted Protein-N" or "CRISP-N") is described.

Identification of a Human and Murine Soggy-1 cDNAs

Human Soggy-1 was identified as a novel protein with similarity to the N-terminal domain of hDkk3. A human partial sequence was identified in the dbEST database for a clone having the accession number AA397836. This clone was obtained from the IMAGE collection and sequenced fully to define the entire human Soggy-1 sequence depicted in Figure 7. Two murine partial sequences were likewise identified in the dBEST database. The clones were obtained from the IMAGE consortium and sequenced. The entire murine Soggy-1 sequence is depicted in Figure 8. Human and murine Soggy cDNAs encode proteins of 242aa and 230aa, respectively, and are predicted to be secreted owing to the presence of N-terminal signal peptides. hSoggy-1 has a predicted signal peptide from about amino acid residue 1 to 30 of SEQ ID NO:14, cleavage of which results in a mature protein having 194 amino acid residues in length and corresponding to amino acid residues 31 to 224 of SEQ ID NO:14. mSoggy-1 has a predicted signal peptide from about amino acid residue 1 to 20 of SEQ ID NO:27,

- 133 -

cleavage of which results in a mature protein having 210 amino acid residues in length and corresponding to amino acid residues 21 to 230 of SEQ ID NO:27. Human and murine Soggy proteins display 59% overall identity although significant amino acid identities are seen beyond this domain that extend into the CRDs of Dkk-3 (Figure 10).

- 5 However, cysteine residues are not conserved within these domains and the residues shared by Soggy and Dkk-3 are poorly conserved in other Dkks indicating that the sequence relationship between these proteins is unique. Homology is most obvious within a 51 amino acid region in which 33% identity is observed between hSoggy, mSoggy, hDkk-3 and mDkk-3. This 51 amino acid domain is referred to herein as an
- 10 SGY domain. Human and mouse Soggy-1 proteins each possess 2 sites of potential N-linked glycosylation which are within the SGY domain and are also conserved with Dkk3. (e.g., NNLT, corresponding to amino acid residues 97-100 of SEQ ID NO:14 and NKTG corresponding to amino acid residues 112-115 of SEQ ID NO:14). In contrast to other Dkks, the C-terminal domain of Soggy-1 shows no similarity to other protein
- 15 sequences in the public databases nor does it contain any cysteine residues. Soggy was so named in view of its lack of CRDs compared to hDkk-3, which had been previously designated Cysteine Rich Secreted Protein-1 ("CRISPY-1").

Tissue Distribution of Soggy-1

- 20 To investigate Soggy-1 mRNA expression, a mouse cDNA probe was used on murine Northern blots. A 1kb mSoggy-1 mRNA was expressed at very high levels in testis and, interestingly, demonstrated transient expression during mouse embryogenesis. Soggy-1 mRNA, which was undetectable at day 7 of gestation, was transiently expressed at day 11 and day 15, after which the expression level declined to
- 25 undetectable levels. Thus, mSoggy-1 displays a developmentally regulated pattern of expression.

In situ analysis was performed as described in Example 1. For detection of murine Soggy-1, the following primers were used:

mSoggy forward 5'-ACCTGCAATGTGTCGACTGAG-3' (SEQ ID NO:32), and

- 30 *reverse* 5'-CACTTACAGCTGTTGGGATG-3' (SEQ ID NO:33).

Consistent with the Northern analysis, very high level expression of Soggy-1 mRNA was observed by *in situ* analysis in adult testis. Upon high magnification, Soggy-1 mRNA was found to be expressed at high levels in the spermatogenic epithelium of the seminiferous tubules and in the spermatogonia at various stages of development. A series of sagittal sections of mouse embryos from E13.5 - E 17.5 and post-natal day 1.5 pups were also analyzed. In E15.5 embryos, Soggy-1 mRNA transcripts were localized to the developing dorsal root ganglia (DRGs) and also found in the cartilage primordium of the nasal septum. Soggy-1 expression was also seen in the eye from E13.5 to E16.5, as observed for mDkk-3. Expression of Soggy-1 mRNA at various stages of development is consistent with the northern analysis described above and suggests that Soggy-1 may play a role in multiple stages of development.

Secretion and Post-Translational Modification of Soggy Proteins

Flag epitope-tagged human Soggy-1 protein was transiently overexpressed in 293T cells and analysed as previously described. hSoggy was efficiently secreted from transfected 293T cells and migrated with a molecular weight of approximately 40-50 kD. Given the heterogenous nature of secreted human Soggy-1, the effect of N-Glycanase treatment on the mobility of secreted flag-tagged hSoggy-1 was studied. hSoggy displayed a 5-10 kD decrease in apparent molecular weight after N-Glycanase treatment, consistent with the presence of 2 potential sites of N-glycosylation in the protein.

Example 8: Structure of the Dkk Family proteins and Dkk-Related Proteins

The amino acid and nucleotide homology between Dkk family members and Dkk-related proteins is set forth in the following tables. Where indicated, mDkk-1 and xDkk-1 correspond to a murine and *Xenopus* proteins set forth in Glinks *et al.*, *supra*, and having Accession Nos: AF030433 and AF030434, respectively. Likewise cDkk-3 has Accession No. D26311

Table II sets forth overall sequence identities as determined using the ALIGN program, (version 2.0) using a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4:

	hDkk-3	hDkk-4	hDkk-1	hDkk-2	mDkk-1	xDkk-1	CLFEST
hDkk-3	100	16.0	18.6	15.1	18.5	16.5	53.0
hDkk-4		100	33.7	35.2	32.6	33.7	16.2
hDkk-1			100	33.1	80.2	53.5	17.4
hDkk-2				100	30.5	33.7	12.5

5

Table III sets forth nucleic acid identities as determined using the using the Wilbur Lipman DNA alignment program, Ktuple: 3; Gap Penalty: 3; Window: 20:

	hDkk-3	hDkk-4	hDkk-1	hDkk-2	mDkk-1	xDkk-1	CLFEST
hDkk-3	100	30.0	37.2	34.7	31.5	45.4	58.8
hDkk-4		100	43.0	35.9	38.8	38.4	36.7
hDkk-1			100	59.3	66.4	53.7	32.1
hDkk-2				100	38.8	38.4	36.7

- 136 -

Table IV sets forth local amino acid identities as determined using the Smith-Waterman algorithm as implemented in the program Bestfit of the GCG package, with gap penalties of 8 for opening and 1 for extending:

hDkk-1								
mDkk-1	82							
xDkk-1	64	63						
hDkk-2	50	48	47					
hDkk-3	39	37	37	37				
mDkk-3	36	33	38	40	83			
cDkk-3	34	31	35	36	61	60		
hDkk-4	45	43	47	46	40	39	34	
	hDkk-1	mDkk-1	xDkk-1	hDkk-2	hDkk-3	mDkk-3	cDkk-3	hDkk-4

5

A multiple alignment of the amino acid sequences of hDkk-1, hDkk-2, hDkk-3, hDkk-4, mDkk-1, mDkk-3, xDkk-1, and cDkk-3 is shown in Figure 6. Predicted signal peptides are underlined, N-glycosylation sites are indicated by a thick bar, CRD-1 by an open box and CRD-2 by a shaded box. The proteolytic cleavage site within hDkk4 is indicated by a double asterisk. The domain structure of the full length human Dkk proteins of the present invention as well as human Soggy are schematically illustrated in Figure 9. Signal peptides (darkened boxes), Cysteine Rich Domain 1 ("CRD-1") (also referred to as the "amino-terminal cysteine-rich domain"), Cysteine Rich Domain 2 ("CRD-2") (also referred to as the "carboxy-terminal cysteine-rich domain"), the soggy domain (SGY) within hDkk-3 and hSoggy-1, and sites of N-glycosylation are indicated.

10

15

As demonstrated at least in Figures 6 and 9, human Dkks 1 through 4 each possess an N-terminal signal peptide and contain two conserved cysteine-rich domains (CRDs) separated by a linker or spacer region. Each CRD possesses 10 conserved cysteine residues. The second CRD has elsewhere been described as a colipase-like domain because the positions of the ten conserved cysteines in this domain have been shown to be closely similar to those in proteins of the colipase family (Aravind and Koonin, *supra*). Conservation of CRD-1 and CRD-2 suggests important functions for

20

- 137 -

these domains. In contrast to the CRDs, the linker or spacer region that joins CRD-1 and CRD-2 is highly variable between hDkks, being notably larger in hDkk-1, -2 and -4 (50-55aa) as compared to Dkk-3 (12aa). Four potential sites of N-linked glycosylation are present in hDkk3 and are conserved in chicken and mouse Dkk-3. These sites are not conserved in other Dkk family members. hDkk1 possesses one potential N-glycosylation site located close to the C-terminus of the protein which is conserved in murine Dkk-1 but not in *Xenopus* Dkk-1 (Fig. 6). In addition, each hDkk possesses several potential dibasic proteolytic cleavage sites, suggesting the proteins may be subject to post-translational processing. hDkk3 is the most divergent of the four human Dkks and possesses an extended N-terminal unique region which precedes CRD-1 and an extended C-terminal unique region which is highly acidic.

Example 9: Effects of hDkks and Soggy on Wnt-induced axis duplication in *Xenopus* embryos

This Example describes the functional activities of the hDkk and Soggy proteins of the present invention.

Xenopus embryo culture and RNA microinjections

Eggs were obtained from *Xenopus* females injected with 700 units of human chorionic gonadotropin, fertilized *in vitro* and cultured in 0.1 x MMR (Newport and Kirschner (1982) *Cell* 30:675-686). Embryonic stages were determined according to Nieuwkoop and Faber (1967) *Normal table of Xenopus laevis* (Daudin) Amsterdam: North Holland Publ. All cDNAs were subcloned into pCS2 vector (Rupp *et al.* (1984) *Genes & Development* 8:1311-1323), and capped mRNAs were synthesized *in vitro* as described (Krieg and Melton (1984) *Nucleic Acids Res.* 12:7057-7070, using the Message Machine kit (Ambion). The following plasmids were used as templates for mRNA synthesis: hDkk-1-pCS2, hDkk-2-pCS2, hDkk-3-pCS2, hDkk-4-pCS2, hSoggy-pCS2, Xwnt8 (Christian *et al.*, (1991) *Development* 111:1045-1055), Xwnt2B (Landesman and Sokol (1997) *Mech. Dev.* 61:1199-209), Xwnt3a (Wolda *et al.* (1993) *Dev. Biol.* 155:46-5), Xfz8-pXT7 (Itoh *et al.* (1998) *Mech. Devel.* 74:145-157), Xdsh-pXT7 (Sokol, *et al.* (1995) *Mech. Devel.* 74:145-157). Protein expression from all

- 138 -

pCS2-Dkk constructs was confirmed by *in vitro* transcription and translation (TNT, Promega). For secondary axis induction, a single ventral blastomere of 4- or 8-cell embryos was injected with 10nl of a solution containing 2-4pg of Xwnt8 mRNA, 2.5-5pg of Xwnt3a mRNA or 10pg of Xwnt2B mRNA as described (Itoh *et al.* (1995) *supra.*). The effect of Dkk RNAs was tested by coinjecting Wnt mRNAs with 2.5ng of hDkk mRNAs. For studies of Frizzled and Dhshevelled, 5ng Fz8 and 1ng Xdsh mRNAs were injected as indicated. After injections, embryos were cultured in 3% Ficoll 400 (Pharmacia), 0.5xMMR. Secondary axes were scored at stage 35 as complete, when they contained anterior neuroectodermal derivatives including pronounced cement gland and eyes, and as partial, when the secondary neural tube with melanocytes, but without head structures, was apparent.

Inhibition of Secondary Axis Induction by hDkk-1 and hDkk-4 in *Xenopus* Embryos

hDkk-1 or hDkk-2 mRNAs were coinjected with Xwnt8 mRNA into single ventral blastomeres of 4- or 8-cell embryos. Injected embryos were cultured for 2 days and secondary axes were scored based on external morphology. Xwnt8 injected embryos displayed complete axis duplication, which was inhibited by co-injection with mRNAs encoding hDkk-1 and hDkk-4. To determine whether hDkks interacted with specific Wnt ligands, several different Wnts were assayed in combination with hDkk-1 or hDkk-4 for secondary axis formation. hDkk-1 and hDkk-4 inhibited axis duplication in response to Xwnt3a and Xwnt2b in addition to Xwnt8. hDkk-1 consistently demonstrated stronger inhibition of Wnt signaling than hDkk-4. Thus, hDkk-1 and hDkk -4 do not show any clear selectivity for the Wnt ligands used in this study. This compares to the FRPs, which also show little specificity with respect to their ability to inhibit Wnts (Leyns *et al.* (1997) *supra*; Wang *et al.* (1997) *supra*; Salic *et al.* (1997) *supra*; Mayr *et al.* (1997) *supra*; Finch *et al.* (1997) *supra*).

To investigate the mechanism by which hDkk-1 and hDkk-4 inhibit Wnt signaling, Dkk mRNAs were coinjected with Xdsh, a downstream component of the Wnt signaling pathway (Itoh *et al.* (1998) *supra*). hDkks-1 and -4 did not block secondary axis formation by Xdsh, indicating that Dkks function upstream of, or parallel with, Xdsh signaling. Similar findings have been reported previously for xDkk-1 (Glinka *et*

al. (1998) *supra*). It was also determined whether hDkks could antagonize signaling by *Xenopus* Frizzled-8 (Xfz8), which can also induce a secondary axis through Wnt signaling (Itoh *et al.* (1998) *supra*). Neither hDkk-1 or hDkk-4 inhibited the axis-inducing activity of Xfz8 mRNA. This data, taken together with the fact that hDkk-1 and hDkk-4 are secreted, indicate that Dkks antagonize Wnt signaling at a point upstream of Wnt receptors.

Assay for Inhibition of Secondary Axis Induction by hDkk-2, hDkk-3 and hSoggy-1 in *Xenopus* Embryos

hDkk-2, hDkk-3 or Soggy mRNAs were coinjected with Xwnt8 mRNA into single ventral blastomeres of 4- or 8-cell embryos and secondary axes were scored after two days as described for hDkk-1 and hDkk-4. Injection of mRNAs encoding hDkk-2, hDkk-3 or hSoggy-1 had no effect on Xwnt8-induced axis duplication. The ability of hDkk-2, hDkk-3 and hSoggy-1 to interact with specific Wnt ligands was also determined as described previously. hDkk-2, hDkk-3 and hSoggy-1 were inactive against each of the three Wnts tested. The lack of activity of hDkk-2, hDkk-3 and hSoggy-1 suggests that these proteins antagonize other members of the Wnt superfamily not tested here, or that they perform functions distinct from Wnt inhibition.

Example 10: Preparation of Antibodies Specific for hDkk and hSoggy Proteins

This example describes the making of polyclonal antibodies specific for hDkk-1, hDkk-4, hDkk-1, hDkk-2, and hSoggy-1.

Peptides were synthesized using Fmoc solid phase methodology utilizing MAP resin technology which increases the antigenic response (Tarn (1988) *Proc. Natl. Acad. Sci. USA* 85:5409-5413. For each protein, the peptides used for immunization are listed below:

hDkk-3

peptide #44 FREVEELMEDTQHKL

peptide #46 GSFMEEVVRQELEDLE

hDkk-4

peptide #91 HAEGTTGHPVQENQP

- 140 -

hDkk-1

peptide #93 GNKYQTIDNYQPYPYPC

hDkk-2

peptide #56 GHYSNHDLGWQNLGR

5 hSoggy-1

peptide #58 LQAIRDGLRKGTHKD

Peptides were designed to meet at least the following criteria: (1) not included within the cysteine-rich domain; (2) not including an N-glycosylation site; and (3)

10 hydrophilic (*e.g.*, solvent exposed).

Antibodies were generated in New Zealand white rabbits over a 10-week period. The immunogen includes KLH-peptide emulsified by mixing with an equal volume of Freund's Adjuvant, and injected into three subcutaneous dorsal sites, for a total of 0.1mg peptide per immunization. Animals were bled from the articular artery.

15 The blood was allowed to clot and the serum collected by centrifugation. The serum is stored at -20°C.

For purification, peptide antigens were immobilized on an activated support. Antisera was passed through the sera column and then washed. Specific antibodies were eluted *via* a pH gradient, collected and stored in a borate buffer (0.125M total

20 borate) at ~0.25mg/ml. The anti-peptide titers were determined using ELISA methodology with free peptide bound in solid phase (1µg/well). Detection was obtained using biotinylated anti-rabbit IgG, HRP-SA conjugate, and ABTS.

All antibodies performed well in ELISA assays. Anti-peptide #44, #46, and #58 are particularly useful for detection of hDkk-3 and hSoggy-1, respectively, as

25 determined by western blotting of supernatants from hDkk-3- and hSoggy-1-transfected 293T cells.

The Dkk family comprises a novel family of secreted proteins which to date

30 includes hDkk-1, hDkk-2, hDkk-3, hDkk-4, xDkk-1, mDkk-1 and cDkk-1. Structurally, Dkks 1-4 are related by several conserved features. Firstly, all four

- 141 -

proteins are secreted proteins. Secondly, Dkks 1-4 each possess two distinct cysteine rich domains. Each domain contains 10 conserved cysteine residues, and these domains are highly conserved between family members. The C-terminal cysteine rich domain, referred to as CRD-2, bears significant similarity to proteins of the colipase
5 family and sequence conservation among the Dkks is greatest within CRD-2 (Aravind and Koonin, *supra*). This may reflect a need for Dkks to interact with lipids in order to regulate Wnt function, since Wnt proteins remain tightly associated with the cell surface.

Despite the similarities between Dkks 1-4, notable differences between these
10 family members appear with regard to their mRNA expression patterns. In adult human tissues hDkk-1 and hDkk-4 showed highly restricted mRNA expression patterns while hDkk-2 and hDkk-3 are more widely expressed. Murine Dkk-3 mRNA was found to be restricted to the myocytes of the atria in the heart, neurons of the cortex and hippocampus in the brain and also to the retinal neurons and lens
15 epithelium in the eye. Such specific expression patterns reflect localized action of the Dkks as regulators of Wnt activity and/or that of other signaling molecules. Different Wnt family members have been shown to have divergent patterns of mRNA expression in adult and embryonic mammalian tissues. For example, murine Wnts 4, 7a and 7b are expressed in brain and lung, whereas Wnt6 is highly expressed in testis
20 (Gavin *et al.*, 1990). Wnts 5b and 13 are more broadly expressed (Gavin *et al.* (1990) *supra*; Katoh *et al.* (1996) *supra*). Although Wnts have been studied mostly in the context of their roles in embryonic development and tumorigenesis, the expression of many family members in normal adult human and mouse tissues, together with their regulators such as the Dkks, suggests that these signaling proteins play important roles
25 in normal tissue homeostasis.

Marked differences in the post-translational processing of different human Dkk proteins was also observed. hDkk-3 is secreted from 293T cells as a heterogeneously glycosylated protein, whereas Dkk-1, 2 and 4 proteins show no evidence of glycosylation. This is consistent with sequence analysis that identifies 4 potential sites
30 of N-glycosylation in the hDkk-3 protein but no sites in either hDkk-2 or hDkk-4. A single putative site in hDkk-1 does not appear to be utilized in 293T cells and may

- 142 -

well not be a significant site for N-linked carbohydrate addition in view of its C-terminal location and lack of conservation with xDkk-1. C-terminal proteolysis of hDkk4 in 293T cells was also characterized. Dkk proteins contain multiple potential sites of proteolytic processing. Secreted hDkk-4 was consistently detected as three
5 major C-terminal fragments. N-terminal sequencing identified two of these as mature, full length hDkk4 and CRD-2, which was derived from the full length protein by a specific proteolytic event at lysines 132 and 133. Thus, the hDkk-4 CRD-2 is released from the full length protein upon expression in 293T cells. Similar processing of hDkk4 in COS cells has been observed.

10 Within the Dkk family, Dkks 1, 2 and 4 display closest similarity whereas Dkk-3 is more distantly related. For example, Dkk-3 contains a linker region connecting CRD-1 and CRD-2 which is shorter than in other Dkks. Dkk-3 also possesses extended N-and C-terminal regions compared to other Dkks. Within the Dkk-3 N-terminal unique region, a distinct domain has been identified that is also
15 found in Soggy (the SGY domain). The SGY domains of human and mouse Soggy-1 and Dkk-3 proteins contain two conserved sites of N-linked glycosylation. Protein expression studies confirm that, like hDkk3, hSoggy is secreted as a glycoprotein. Murine Soggy-1 is highly expressed in adult testis and also displays a highly restricted mRNA expression in E15 -E16 mouse embryos, being localized mainly to the cartilage
20 primordia within the developing vertebrae/developing dorsal root ganglia. Soggy mRNA was also detected at high levels in the developing eye, similar to Dkk-3. This developmentally regulated pattern of Soggy expression overlaps with that of Dkk-3 suggesting that Soggy may play a role in the regulation of Dkk-3 function.

25 **Equivalents**

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

What is claimed:

1. An isolated nucleic acid molecule selected from the group consisting of:
 - a) a nucleic acid molecule comprising a nucleotide sequence which
5 is at least 60% homologous to a nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:20, SEQ ID NO:22, the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide
10 sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a complement thereof;
 - b) a nucleic acid molecule comprising a fragment of at least 1000 nucleotides of a nucleic acid comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID
15 NO:9, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:20, SEQ ID NO:22, the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a complement thereof;
 - c) a nucleic acid molecule which encodes a polypeptide comprising
20 an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, or SEQ ID NO:21, the amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the amino acid
25 sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the amino acid sequence encoded by the DNA insert of the plasmid deposited with the ATCC as Accession Number _____;
 - d) a nucleic acid molecule which encodes a fragment of a
30 polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number

- 144 -

98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, wherein the fragment comprises at least 15 contiguous amino acid residues of the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with the ATCC as Accession Number _____; and

e) a nucleic acid molecule which encodes a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, wherein the nucleic acid molecule hybridizes to a nucleic acid molecule comprising SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:20, SEQ ID NO:22, the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the DNA insert of the plasmid deposited with Accession Number _____ under stringent conditions.

- 145 -

2. The isolated nucleic acid molecule of claim 1 which is selected from the group consisting of:

- 5 a) a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:20, SEQ ID NO:22, the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a complement thereof;
- 10 and
- b) a nucleic acid molecule which encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with the ATCC as Accession Number _____.
- 15

3. The nucleic acid molecule of claim 1 further comprising vector nucleic acid sequences.

20

4. The nucleic acid molecule of claim 1 further comprising nucleic acid sequences encoding a heterologous polypeptide.

25 5. A host cell which contains the nucleic acid molecule of claim 1.

6. The host cell of claim 5 which is a mammalian host cell.

7. A non-human mammalian host cell containing the nucleic acid molecule of claim 1.

30

- 146 -

8. An isolated polypeptide selected from the group consisting of:

a) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with the ATCC as Accession Number _____;

b) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule comprising SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:20, SEQ ID NO:22, the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the DNA insert of the plasmid deposited with the ATCC as Accession Number _____ under stringent conditions;

c) a polypeptide which is encoded by a nucleic acid molecule comprising a nucleotide sequence which is at least 60% homologous to a nucleic

acid comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:20, SEQ ID NO:22, the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the DNA insert of the plasmid deposited with ATCC as Accession Number _____; and

d) a polypeptide comprising an amino acid sequence which is at least 60% homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____.

9. The isolated polypeptide of claim 8 comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____.

10. The polypeptide of claim 8 further comprising heterologous amino acid sequences.

11. An antibody which selectively binds to a polypeptide of claim 8.

- 148 -

12. A method for producing a polypeptide selected from the group consisting of:

- 5 a) a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____;
- 10 b) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the
- 15 nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide
- 20 encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____; and
- 25 c) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:21, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, wherein the polypeptide is
- 30 encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule comprising SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID

- 149 -

NO:7, SEQ ID NO:9, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:20, SEQ ID NO:22, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the
5 nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____ under stringent conditions;

comprising culturing the host cell of claim 5 under conditions in which the nucleic acid molecule is expressed.

10

13. A method for detecting the presence of a polypeptide of claim 8 in a sample comprising:

- a) contacting the sample with a compound which selectively binds to the polypeptide; and
- 15 b) determining whether the compound binds to the polypeptide in the sample to thereby detect the presence of a polypeptide of claim 8 in the sample.

14. The method of claim 13, wherein the compound which binds to the
20 polypeptide is an antibody.

15. A kit comprising a compound which selectively binds to a polypeptide of claim 8 and instructions for use.

25 16. A method for detecting the presence of a nucleic acid molecule in claim 1 in a sample comprising:

- a) contacting the sample with a nucleic acid probe or primer which selectively hybridizes to the nucleic acid molecule; and
- 30 b) determining whether the nucleic acid probe or primer binds to a nucleic acid molecule in the sample to thereby detect the presence of a nucleic acid molecule of claim 1 in the sample.

- 150 -

17. The method of claim 16, wherein the sample comprises mRNA molecules and is contacted with a nucleic acid probe.

18. A kit comprising a compound which selectively hybridizes to a nucleic acid molecule of claim 1 and instructions for use.

19. A method for identifying a compound which binds to a polypeptide of claim 8 comprising:

- a) contacting the polypeptide, or a cell expressing the polypeptide with a test compound; and
- b) determining whether the polypeptide binds to the test compound.

20. The method of claim 19, wherein the binding of the test compound to the polypeptide is detected by a method selected from the group consisting of:

- a) detection of binding by direct detection of test compound/polypeptide binding;
- b) detection of binding using a competition binding assay; and
- c) detection of binding using an assay for Dkk activity.

21. A method of modulating the activity of a polypeptide of claim 8 comprising contacting the polypeptide or a cell expressing the polypeptide with a compound which binds to the polypeptide in a sufficient concentration to modulate the activity of the polypeptide.

22. A method for identifying a compound which modulates the activity of a polypeptide of claim 8 comprising:

- a) contacting a polypeptide of claim 8 with a test compound; and
- b) determining the effect of the test compound on the activity of the polypeptide to thereby identify a compound which modulates the activity of the polypeptide.

FIG. 1A-1

```

      M   Q   R   L   G   A   T   L   L   C
GGCACGAGGGGCGGCGGCTGCGGGCGCAGAGCGGAG ATG CAG CGG CTT GGG GCC ACC CTG CTG TGC
10
      67
L   L   A   A   A   V   P   T   A   P   A   P   T   A   T   S   A
CTG CTG GCG GCG GCG GTC CCC ACG GCC CCC GCG CCC GCT CCG ACG GCG ACC TCG GCT
30
      127
P   V   K   P   G   P   A   L   S   Y   P   Q   E   E   A   T   L   N   E   M
CCA GTC AAG CCC GGC CCG GCT CTC AGC TAC CCG CAG GAG GAG GCC ACC CTC AAT GAG ATG
50
      187
F   R   E   V   E   E   L   M   E   D   T   Q   H   K   L   R   S   A   V   E
TTC CGC GAG GTT GAG GAA CTG ATG GAG GAC ACG CAG CAG CAC AAA TTG CGC AGC GCG GTG GAA
70
      247
E   M   E   A   E   E   A   A   A   K   A   S   S   E   V   N   L   A   N   L
GAG ATG GAG GCA GAA GAA GCT GCT GCT AAA GCA TCA TCA GAA GTG AAC CTG GCA AAC TTA
90
      307
P   P   S   Y   H   N   E   T   N   T   D   T   K   V   G   N   N   T   I   H
CCT CCC AGC TAT CAC AAT GAG ACC AAC ACA GAC ACG AAC GAT GTT GGA AAT AAT ACC ATC CAT
110
      367
V   H   R   E   I   H   K   I   T   N   N   Q   T   G   Q   M   V   F   S   E
GTG CAC CGA GAA ATT CAC AAG ATA ACC AAC AAC CAG CAG ACT GGA CAA ATG GTC TTT TCA GAG
130
      427
T   V   I   T   S   V   G   D   E   E   G   R   R   S   H   E   C   I   I   D
ACA GTT ATC ACA TCT GTG GGA GAC GAA GAA GGC AGA AGG AGC CAC GAG TGC ATC ATC GAC
150
      487
E   D   C   G   P   S   M   Y   C   Q   F   A   S   F   Q   Q   Y   T   C   Q   P
GAG GAC TGT GGG CCC AGC ATG ATG TAC TGC CAG TTT GCC AGC TTC CAG TAC ACC TGC CAG CCA
170
      547
C   R   G   Q   R   M   L   C   T   R   D   S   E   C   C   G   D   Q   L   C
TGC CGG GGC CAG AGG ATG CTC TGC ACC CGG GAC AGT GAG TGC TGT GGA GAC CAG CTG TGT
190
      607

```

FIG. 1A-2

V	W	G	H	C	T	K	M	A	T	R	G	S	N	G	T	I	C	D	N	210
GTC	TGG	GGT	CAC	TGC	ACC	AAA	ATG	GCC	ACC	AGG	GGC	AGC	AAT	GGG	ACC	ATC	TGT	GAC	AAC	667
Q	R	D	C	Q	P	G	L	C	C	A	F	Q	R	G	L	L	F	P	V	230
CAG	AGG	GAC	TGC	CAG	CCG	GGG	CTG	TGC	TGT	GCC	TTC	CAG	AGA	GGC	CTG	CTG	TTC	CCT	GTG	727
C	T	P	L	P	V	E	G	E	L	C	H	D	P	A	S	R	L	L	D	250
TGC	ACA	CCC	CTG	CCC	GTG	GAG	GGC	GAG	CTT	TGC	CAT	GAC	CCC	GCC	AGC	CGG	CTT	CTG	GAC	787
L	I	T	W	E	L	E	P	D	G	A	L	D	R	C	P	C	A	S	G	270
CTC	ATC	ACC	TGG	GAG	CTA	GAG	CCT	GAT	GGA	GCC	TTG	GAC	CGA	TGC	CCT	TGT	GCC	AGT	GGC	847
L	L	C	Q	P	H	S	H	S	L	V	Y	V	C	K	P	T	F	V	G	290
CTC	CTC	TGC	CAG	CCC	CAC	AGC	CAC	AGC	CTG	GTG	TAT	GTG	TGC	AAG	CCG	ACC	TTC	GTG	GGG	907
S	R	D	Q	D	G	E	I	L	L	P	R	E	V	P	D	E	Y	E	V	310
AGC	CGT	GAC	CAA	GAT	GGG	GAG	ATC	CTG	CTG	CCC	AGA	GAG	GTC	CCC	GAT	GAG	TAT	GAA	GTT	967
G	S	F	M	E	E	V	R	Q	E	L	E	D	L	E	R	S	L	T	E	330
GGC	AGC	TTC	ATG	GAG	GAG	GTG	CGC	CAG	GAG	CTG	GAG	GAC	CTG	GAG	AGG	AGC	CTG	ACT	GAA	1027
E	M	A	L	G	E	P	A	A	A	A	A	A	L	L	G	G	E	E	I	350
GAG	ATG	GCG	CTG	GGG	GAG	CCT	GCG	GCT	GCC	GCC	GCT	GCA	CTG	CTG	GGA	GGG	GAA	GAG	ATT	1087
*																				351
TAG																				1090

FIG. 1B-1

ATCTGGACCAGGCTGTGGGTAGATGTGCAATAGAAATAGCTAATTTATTTCCCCANGTGTGTGCTTTAAAGCGTGGGCTG 1169

ACCAGGCTTCTTCCCTACATCTTCTTCCCAGTAAGTTTCCCCCTCTGGCTTGACAGCATGAGGTGTTGTGCAATTTGTTTCAG 1248

CTCCCCCAGGCTGTTCTCCAGGCTTTCACAGTCTGGTGCTTGGGAGAGTCAGGCAGGGTTAAACTGCAGGAGCAGTTTGC 1327

CACCCCTGTCCAGATTATTGGCTGCTTGGCTCTACCCAGTTGGCAGACAGCCGTTTGTTCATCATGGCTTTGATATAATTG 1406

TTTGAGGGGAGGAGATGGAACAATGTGGAGTCTCCCTCTGATTGGTTTGGGGAAATGTGGAGAAAGAGTGCCCCTGCTT 1485

TGCAAAACATCAACCTGGCAAAAATGCAACAAATGAATTTCCACGCAGTTCTTTCCATGGGCATAGGTAAGCTGTGCCT 1564

TCAGCTGTTGCAGATGAAATGTTCTGTTCACCCTGCATTACATGTGTGTTTATTTCATCCAGCAGTGTGTGCTCAGCTCCTAC 1643

CTCTGTGCCAGGCAGCATTTTCATATCCAAGATCAATTCCTCTCTCAGCACAGCCCTGGGGAGGGGTCATTGTTCTC 1722

CTCGTCCATCAGGGATTTTCAGAGGCTCAGAGACTGCAAGCTGCTTGCCCAAGTCACACAGCTAGTGAAGACCAGAGCAG 1801

TTTCATCTGGTTGTGACTCTAAGCTCAGTGCTCTCTCCACTACCCACACAGCCCTTGGTGCCACCAAAAGTGCTCCCC 1880

AAAAGGAAGGAGAAATGGGATTTTCTTTTGAGGCATGCACATCTGGAATTAAGGTCAAACTAATTCTCACATCCCCTCTA 1959

AAAGTAAACTACTGTTAGGAACAGCAGTGTTCACAGTGTGGGGCAGCCGTCCTTCTAATGAAGACAATGATATTGAC 2038

ACTGTCCCTCTTTGGCAGTTGCATTAGTAACTTTGAAAAGGTATATGACTGAGCGTAGCATACAGGTTAACCTGCAGAAA 2117

CAGTACTTAGGTAATTGTAGGGCGGAGGATTATAAATGAAATTTGCAAAATCACTTAGCAGCAACTGAAGACAATTATCA 2196

FIG. 1B-2

ACCACGTGGAGAAAATCAAAACCGAGCAGGGCTGTGTGAAACATGGTTGTAATATGCGACTGCCAACACTGAACTCTACG 2275

CCACTCCACAAATGATGTTTTTCAGGTGTCATGGACTGTGCCCACCATGTATTTCATCCAGAGTTCCTTAAAGTTTAAAGTT 2354

GCACATGATTGTATAAGCATGCTTTCTTTGAGTTTAAATTATGTATAAACATAAAGTTGCATTTAGAAATCAAGCATAA 2433

ATCACTTCAACTGCTAAAAAATAAAAAAAAAAAAAAAAAAAAAA 2479

5/29

FIG. 2A

GAATTGGCACGAGACGACGCTGCTGAGCTGCCAGCTTAGTGGAAGCTCTGCTCTGGGTGGAGAGCAGCCTCGCTTTG	79
GTGACGCACAGTGTGGGACCCCTCCAGAGCCCCGGGATTGAAGG ATG GTG GCG GCC GTC CTG CTG GGG	148
<div style="text-align: center;">M V A A V L L G</div>	8
L S W L C S P L G A L V L D F N N I R S	28
CTG AGC TGG CTC TGC TCT CCC CTG GGA GCT CTG GTC GAC TTC AAC AAC ATC AGG AGC	208
S A D L H G A R K G S Q C L S D T D C N	48
TCT GCT GAC CTG CAT GGG GCC CGG AAG GGC TCA CAG TGC CTG TCT GAC ACG GAC TGC AAT	268
T R K F C L Q P R D E K P F C A T C R G	68
ACC AGA AAG TTC TGC CTC CAG CCC CGC GAT GAG AAG CCG TTC TGT GCT ACA TGT CGT GGG	328
L R R R C Q R D A M C C P G T L C V N D	88
TTG CGG AGG AGG TGC CAG CGA GAT GCC ATG TGC TGC CCT GGG ACA CTC TGT GTG AAC GAT	388
V C T T M E D A T P I L E R Q L D E Q D	108
GTT TGT ACT ACG ATG GAA GAT GCA ACC CCA ATA TTA GAA AGG CAG CTT GAT GAG CAA GAT	448
G T H A E G T T G H P V Q E N Q P K R K	128
GGC ACA CAT GCA GAA GGA ACA ACT GGG CAC CCA GTC CAG GAA AAC CAA CCC AAA AGG AAG	508
P S I K K S Q G R K G Q E G E S C L R T	148
CCA AGT ATT AAG AAA TCA CAA GGC AGG AAG GGA GAA AGT TGT CTG AGA ACT	568

6/29

FIG. 2B

F	D	C	T	G	G	C	C	A	R	H	F	W	T	K	I	C	K	P	168	
TTT	GAC	TGT	GGC	CCT	GGA	CTT	TGC	TGT	GCT	CAT	TTT	TGG	ACG	AAA	ATT	TGT	AAG	CCA	628	
V	L	L	E	G	Q	V	C	S	R	R	G	H	K	D	T	A	Q	A	P	188
GTC	CTT	TTG	GAG	GGA	CAG	GTC	TGC	TCC	AGA	AGA	GGG	CAT	AAA	GAC	ACT	GCT	CAA	GCT	CCA	688
E	I	F	Q	R	C	D	C	G	P	G	L	L	C	R	S	Q	L	T	S	208
GAA	ATC	TTC	CAG	CGT	TGC	GAC	TGT	GGC	CCT	GGA	CTA	CTG	TGT	CGA	AGC	CAA	TTG	ACC	AGC	748
N	R	Q	H	A	R	L	R	V	C	Q	K	I	E	K	L	*				225
AAT	CGG	CAG	CAT	GCT	CGA	TTA	AGA	GTA	TGC	CAA	AAA	ATA	GAA	AAG	CTA	TAA				799
ATATTTCAAAAATAAAGAAGAAATCCACATTGCAAAAAAATAAAAAA																			848	

7/29

FIG. 3A

GTCGACCCACGCGTCCGCGACGCGTGGCGGCACGGTTTCGTGGGACCCAGGCTTGCAAAGTGACGGTCATTTC	79													
TTTCTTTCTCCTCTTGAGTCCTTCTGAG ATG ATG GCT CTG GGC GCA GCG GGA GCT ACC CGG GTC	144													
<table border="0"> <tr> <td>M</td><td>M</td><td>A</td><td>L</td><td>G</td><td>A</td><td>A</td><td>G</td><td>A</td><td>T</td><td>R</td><td>V</td> <td>12</td> </tr> </table>	M	M	A	L	G	A	A	G	A	T	R	V	12	
M	M	A	L	G	A	A	G	A	T	R	V	12		
F V A M V A A A L G G H P L L G V S A T	32													
TTT GTC GCG ATG GTA GCG GCT CTC GGC GGC CAC CCT CTG CTG GGA GTG AGC GCC ACC	204													
L N S V L N S N A I K N L P P L G G A	52													
TTG AAC TCG GTT CTC AAT TCC AAC GCT ATC AAG AAC CTG CCC CCA CCG CTG GGC GGT	264													
A G H P G S A V S A A P G I L Y P G G N	72													
GCG GGG CAC CCA GGC TCT GCA GTC AGC GCC CCG GGA ATC CTG TAC CCG GGC GSG AAT	324													
K Y Q T I D N Y Q P Y P C A E D E E C G	92													
AAG TAC CAG ACC ATT GAC AAC TAC CAG CCG TAC CCG TGC GCA GAG GAC GAG TGC GGC	384													
T D E Y C A S P T R G G G A G V Q I C L	112													
ACT GAT GAG TAC TGC GCT AGT CCC ACC CGC GGA GGC GAC GCA GGC GTG CAA ATC TGT CTC	444													
A C R K R R K R C M R H A M C C P G N Y	132													
GCC TGC AGG AAG CGC CGA AAA CGC TGC ATG CGT CAC GCT ATG TGC TGC CCC GGG AAT TAC	504													
C K N G I C V S S D Q N H F R G E I E E	152													
TGC AAA AAT GGA ATA TGT GTG TCT TCT TCT GAT CAA AAT CAT TTC CGA GGA GAA ATT GAG GAA	564													
T I T E S F G N D H S T L D G Y S R R T	172													
ACC ATC ACT GAA AGC TTT GGT AAT GAT CAT AGC ACC TTG GAT GGG TAT TCC AGA AGA ACC	624													

8/29

FIG. 3B

T L S S K M Y H T K G Q E G S V C L R S 192
 ACC TTG TCT TCA AAA ATG TAT CAC ACC AAA GGA CAA GAA GGT TCT GTT TGT CTC CGG TCA 684

 S D C A S G L C C A R H F W S K I C K P 212
 TCA GAC TGT GCC TCA GGA TTG TGT TGT GCT AGA CAC TTC TGG TCC AAG ATC TGT AAA CCT 744

 V L K E G Q V C T K H R R K G S H G L E 232
 GTC CTG AAA GAA GGT CAA GTG TGT ACC AAG CAT AGG AGA AAA GGC TCT CAT GGA CTA GAA 804

 I F Q R C Y C G E G L S C R I Q K D H H 252
 ATA TTC CAG CGT TGT TAC TGT GGA GAA GGT CTG TCT TGC CGG ATA CAG AAA GAT CAC CAT 864

 Q A S N S S R L H T C Q R H * 267
 CAA GCC AGT AAT TCT TCT AGG CTT CAC ACT TGT CAG AGA CAC TAA 909

 ACCAGCTATCCAAATGCAGTGAACCTCTTTTATATAATAGATGCTATGAAACCTTTTATGACCTTCATCAACTCAATC 988

 CTAAGGATATACAAAGTTCTGTGTTTCAGTTAAGCATTCCTCAATAACACCTTCCAAAAACCTGGAGTGTAAAGAGCTTTGT 1067

 TTCTTTATGGAACTCCCCCTGTGATTGCAGTAAATTACTGTATTGTAAATTTCTCAGTGTGGCACTTACCTGTAAATGCAA 1146

 TGAAACTTTTAAATTATTTTCTAAAGGTGCTGCACCTGCCTATTTTTCCTCTTGTATGTAAATTTTGTACACATTGAT 1225

 TGTATCTTGACTGCACAAATATTCTATATTGAACTGAAGTAAATCATTTTCAGCTTATAGTTCTTAAAAAGCATAAACCTT 1304

 TACCCCATTTAATTCTAGAGTCTAGAACGCAAGGATCTCTTGGAATGACAAATGATAGGTACCTAAAAATGTAACATGAA 1383

 AATACTAGCTTATTTTCTGAAAATGTACTATCTTAAATGCTTAAATATATTTCCCTTTAGGCTGTGATAGTTTTTGAAAT 1462

 AAAATTTAACATTTAATATCATGAAATGTTTATAAGTAGACATAAAAAAAAAGGGCGGCCGC 1536

FIG. 4A-1

GTCGACCCACGCTCCGGGGAGCCCGGGGAGCGTAGCGCAAGTCCGCTCCCTAGGCATCGCTGCGCTGCGACGCA 79
TTCGCTGTCTCTTGTGAGTCAGGGGACAAACGCTTCGGGGCAACTGTGAGTGGCGGTGTGGGGACCTCGATTCTCTTCA 158
GATCTCGAGGATTCCGGTCCGGGGACGCTCTCTGATCCCTTACTAAAGCGCCTGCTAACTTTGAAAAGGAGCACTGTGTC 237
CTGCAAAAGTTTGACACATAAAGGATAGGAAAAGAGAGAGAAAAGCAACTGAGTTGAAGGAGAGAGCTGATGCGG 316
GCCCTCCTGATCAATTAAAGAGGAGAGTTAAACCGCCGAGATCCCGGGGGACCAAGGAGGTGCGGGGCAAGAAGGAACGG 395
AAGCGGTCCGATCCACAGGGCTGGGTTTTCTTGCACCTTGGGTACGCCCTCCTTGGCGAGAAAGCGCCTCGCATTTGAT 474
TGCTTCCAGTTATTGCAGAACTTCCTGTCTCCTGGTGAGAAAGCGGTCTCGCTTGGGTTCCGCTAATTTCTGTCTGTGAGG 553
CGTGAGACTGAGTTCATAGGGTCCCTGGGTCCCCGAACCAAGGAGTTGAGGGAACACAAATCTGCAAGCCCCCGCGACC 632
CAAGTGAGGGCCCCGTGTGGGGTCCCTCCCTCCCTTTCATTTCCACCCCTCCGGGCTTTGCGTCTTCTCTGGGGACCC 711
M A A L M R S K D S S C C L L L L L 17
CCTCGCCGGGAG ATG GCC GCG TTG ATG CGG AGC AAG GAT TCG TCC TCG TGC TGC CTC CTC CTA CTG 774
A A V L M V E S S Q I G S S R A K L N S 37
GCC GCG GTG CTG ATG GTG GAG AGC TCA CAG ATC GGC AGT TCG CGG GCC AAA CTC AAC TCC 834
I K S S L G G E T P G Q A A N R S A G M 57
ATC AAG TCC TCT CTG GGC GGG GAG ACG CCT GGT CAG GCC AAT CGA TCT GCG GGC ATG 894

10/29

FIG. 4A-2

Y	Q	G	L	A	F	G	G	S	K	K	G	K	N	L	G	Q	A	Y	P	77
TAC	CAA	GGA	CTG	GCA	TTC	GGC	GGC	AGT	AAG	AAG	GGC	AAA	AAC	CTG	GGG	CAG	GCC	TAC	CCT	954
C	S	S	D	K	E	C	E	V	G	R	Y	C	H	S	P	H	Q	G	S	97
TGT	AGC	AGT	GAT	AAG	GAG	TGT	GAA	GTT	GGG	AGG	TAT	TGC	CAC	AGT	CCC	CAC	CAA	GGA	TCA	1014
S	A	C	M	V	C	R	R	K	K	K	R	C	H	R	D	G	M	C	C	117
TCG	GCC	TGC	ATG	GTG	TGT	CGG	AGA	AAA	AAG	AAG	CGC	TGC	CAC	CGA	GAT	GGC	ATG	TGC	TGC	1074
P	S	T	R	C	N	N	G	I	C	I	P	V	T	E	S	I	L	T	P	137
CCC	AGT	ACC	CGC	TGC	AAT	AAT	GGC	ATC	TGT	ATC	CCA	GTT	ACT	GAA	AGC	ATC	TTA	ACC	CCT	1134
H	I	P	A	L	D	G	T	R	H	R	D	R	N	H	G	H	Y	S	N	157
CAC	ATC	CCG	GCT	CTG	GAT	GGT	ACT	CGG	CAC	AGA	GAT	CGA	AAC	CAC	GGT	CAT	TAC	TCA	AAC	1194
H	D	L	G	W	Q	N	L	G	R	P	H	T	K	M	S	H	I	K	G	177
CAT	GAC	TTG	GGA	TGG	CAG	AAT	CTA	GGA	AGA	CCA	CAC	ACT	AAG	ATG	TCA	CAT	ATA	AAA	GGG	1254
H	E	G	D	P	C	L	R	S	S	D	C	I	E	G	F	C	C	A	R	197
CAT	GAA	GGA	GAC	CCC	TGC	CTA	CGA	TCA	TCA	GAC	TGC	ATT	GAA	GGG	TTT	TGC	TGT	GCT	CGT	1314
H	F	W	T	K	I	C	K	P	V	L	H	Q	G	E	V	C	T	K	Q	217
CAT	TTC	TGG	ACC	AAA	ATC	TGC	AAA	CCA	GTG	CTC	CAT	CAG	GGG	GAA	GTC	TGT	ACC	AAA	CAA	1374
R	K	K	G	S	H	G	L	E	I	F	Q	R	C	D	C	A	K	G	L	237
CGC	AAG	AAG	GGT	TCT	CAT	GGG	CTG	GAA	ATT	TTC	CAG	CGT	TGC	GAC	TGT	GCG	AAG	GGC	CTG	1434
S	C	K	V	W	K	D	A	T	Y	S	S	K	A	R	L	H	V	C	Q	257
TCT	TGC	AAA	GTA	TGG	AAA	GAT	GCC	ACC	TAC	TCC	TCC	AAA	GCC	AGA	CTC	CAT	GTG	TGT	CAG	1494

11/29

FIG. 4A-3

K I * 260
AAA ATT TGA 1503

TCACCATTGAGGAACATCATCAATTGCAGACTGTGAAGTTGTGTATTTAATGCATTATAGCATGGTGGAATAAGGTT 1582

CAGATGCAGAAAGAA TGGCTAAAA TAAGAAACGGTGATAAGAATATAGATGATCACAAAAAGGGAGAGAAAACATGAAC 1661

TGAATAGATTAGAAATGGGTGACAAATGCAGTGCAGCCAGTGTTTCCATTATGCAACTTGTCTATGTAAATAATGTACAC 1740

12/29

FIG. 4B-1

ATTGTGGAAAATGCTATTATTAAAGAGAACAAAGCACACAGTGGAATTAAGTAGCATGTGACTTTCCAAAGAGT 1819
TTAGGTTGTGCTGGAGGAGAGGTTTCCTTCAGATTGCTGATTGCTTATACAAATAACCTACATGCCAGATTTCTATTCA 1898
ACGTTAGAGTTTAAACAAAATACTCCTAGAAATAACTTGTTATACAAATAGGTTCTAAAAATAAAATTGCTAAACAAGAAAT 1977
GAAACATGGAGCATTGTTAATTACAAACAGAAAAATTACCTTTTGATTGTGAACACTACTTCTGCTGTTCAATCAAGAG 2056
TCTTGGTAGATAAGAAAAAATCAGTCAATATTTCCAAAATAATTGCCAAAATAATGGCCAGTTGTTTAGGAAGGCCCTTTA 2135
GGAGACAAATAAATAACAAACAAACAGCCACAAATACTTTTTTTTCCAAAATTTTAGTTTTACCTGTAAATAAAGAA 2214
CTGATACAGACAAAAACAGTTCCTTCAGATTCTACGGAAATGACAGTATATCTCTTTATCCTATGTGTGATTCCCTGCTC 2293
TGAATGCATTATATTTCCAAAAGTATACCCATAAAATTTGTGACTAGTAAAAATACTTACACAGAGCAGAAATTTTCACAGAT 2372
GGCAAAAAAATTTAAAGATGTCCAATATATGTGGGAAAAGAGCTAACAGAGAGATCATATTCTTAAAGATTGGCCAT 2451
AACCCTGTATTTTGATAGAAATTAGATTGGTAAATACATGTATTATACATACTCTGTGGTAATAGAGACTTGAGCTGGAT 2530
CTGTACTGCACCTGGAGTAAGCAAGAAAATTTGGGAAAACCTTTTTCGTTTGTTCAGGTTTGGCAACACATAGATCATATG 2609
TC TGAGGCACAAGTTGGCTGTTTCATCTTTGAAACCAGGGGATGCACAGTCTAAATGAATATCTGCATGGGATTGCTAT 2688
CATAATNTTTCCTATGCNGNTGAATTCNGTGTGAGGTCCCTGTGTCCGTCCTATCCCTCAAATTAATTTATTTATAGTGCT 2767
GAGATCCCTCAAATAATCTCAATTTCCGAGGTTTCACAAAAATGGACTCCTGAAGTAGACAGAGTAGTGAGGTTTCATTGC 2846

FIG. 4B-2

CCTCTATAAGCTTCTGACTAGCCAAATGGCAATCATCCAAATTTTCTTCCCAAACCTCTGCAGCATCTGCTTTATTTGCCAAA 2925
GGGCTAGTTTCGGTTTTCTGCCCAGCCATTGCGGGTTAAAAAATATAAGTAGGATAACTTGTAACCTGCATATTGCTAA 3004
TCTATAGACACCACAGTTTCTAAATTCTTTGAAACCACCTTTACTACTTTTTTAAACTTAACCTCAGTTCTAAATACTTT 3083
GTCTGGAGCACAAAACAATAAAGGTTATCTTATAGTTGTGACTTTTAAACCTTTGTAGACCACAATTCACCTTTTGTAGTT 3162
TTCTTTTACTTAAATCCCATCTGCAGTCTCAAAATTTAAGTTCTCCCAGTAGAGATTGAGTTTGAGCCCTGTATATCTATT 3241
AAAAATTCAACTTCCCACATATATTACTAAGATGATTAAGACTTACATTTTCTGCACAGGCTCTGCAAAAAACAAAAAT 3320
TATAAACTAGTCCATCCAAGAACCAAGTTTGATATAACAGGTTGCTATAAGCTTGGTGAAATGAAAAATGGAACATTTC 3399
AATCAAAACATTTCCCTATATAAACAATTATATATTACAAATTTGGTTTCTGCAATATTTTCTTATGTCCACCCCTTTTAA 3478
AAATTATTATTGAAGTAATTTATTTACAGGAAATGTTAATGAGATGTATTTTCTTATAGAGATATTTCTTACAGAAAG 3557
CTTTGTAGCAGAAATATATTTCAGCTATTGACTTTGTAAATTTAGGAAAAAATGTATAATAAGATAAAATCTATAAAATTT 3636
TTCTCCTCTAAAAACTGAAAAAATAAAAAAAGGGCGCGC 3687

14/29

FIG. 5A-1

FGTCACCCACGGCTCCGCTGTGGCAGCCAGCTACCGGTCCGTGACCATCCAGCTTGACGCTCAGCTTTGTTCATTC	79												
GAATTGGCGGGCCAGCGCGGAACAAAC ATG CAG CGG CTC GGG GGT ATT TTG CTG TGT ACA CTG	12 145												
<table border="0"> <tr> <td>M</td><td>Q</td><td>R</td><td>L</td><td>G</td><td>G</td><td>I</td><td>L</td><td>L</td><td>C</td><td>T</td><td>L</td> </tr> </table>	M	Q	R	L	G	G	I	L	L	C	T	L	
M	Q	R	L	G	G	I	L	L	C	T	L		
L A A A V P T A P A P S P T V T W T P A	32												
CTG GCG GCG GCG CTC CCC ACT GCT CCT GCT CCT TCC CCG ACG GTC ACT TGG ACT CCG GCG	205												
E P G P A L N Y P Q E E A T L N E M F R	52												
GAG CCG GGC CCA GCT CTC AAC TAC CCT CAG GAG GAA GCT ACG CTC AAT GAG ATG TTT CGA	265												
E V E E L M E D T Q H K L R S A V E E M	72												
GAG GTG GAG GAG CTG ATG GAA GAC ACT CAG CAC AAA CTG CGC AGT GCC GTG GAG GAG ATG	325												
E A E E A A A K T S S E V N L A S L P P	92												
GAG GCG GAA GAA GCA GCT GCT AAA ACG TCC TCT GAG GTG AAC CTG GCA AGC TTA CCT CCC	385												
N Y H N E T S T E T R R V G N N T V H V H	112												
AAC TAT CAC AAT GAG ACC AGC ACG GAG ACC AGG GTG GGA AAT AAC ACA GTC CAT GTG CAC	445												
Q E V H K I T N N Q S G Q V V F S E T V	132												
CAG GAA GTT CAC AAG ATA ACC AAC AAC CAG AGT GGA CAG GTG GTC TTT TCT GAG ACA GTC	505												
I T S V G D E E G K R S H E C I I D E D	152												
ATT ACA TCT GTA GGG GAT GAA GAA GGC AAG AGG AGC CAT GAA TGT ATC ATT GAT GAA GAC	565												
C G P T R Y C Q F S S F K Y T C Q P C R	172												
TGT GGG CCC ACC AGG TAC TGC CAG TTC TCC AGC TTC AAG TAC ACC TGC CAG CCA TGC CGG	625												

FIG. 5A-2

D	Q	Q	M	L	C	T	R	D	S	E	C	C	G	D	Q	L	C	A	W	192
GAC	CAG	CAG	ATG	CTA	TGC	ACC	CGA	GAC	AGT	GAG	TGC	TGT	GGA	GAC	CAG	CTG	TGT	GCC	TGG	685
G	H	C	T	Q	K	A	T	K	G	G	N	G	T	I	C	D	N	Q	R	212
GGT	CAC	TGC	ACC	CAA	AAG	GCC	ACC	AAA	GGT	GGC	AAT	GGG	ACC	ATC	TGT	GAC	AAC	CAG	AGG	745
D	C	Q	P	G	L	C	C	A	F	Q	R	G	L	L	F	P	V	C	T	232
GAT	TGC	CAG	CCT	GGC	CTG	TGT	TGT	GCC	TTC	CAA	AGA	GGC	CTG	CTG	TTC	CCC	GTG	TGC	ACA	805
P	L	P	V	E	G	E	L	C	H	D	P	T	S	Q	L	L	D	L	I	252
CCC	CTG	CCC	GTG	GAG	GGA	GAG	CTC	TGC	CAT	GAC	CCC	ACC	AGC	CAG	CTG	CTG	GAT	CTC	ATC	865
T	W	E	L	E	P	E	G	A	L	D	R	C	P	C	A	S	G	L	L	272
ACC	TGG	GAA	CTG	GAG	CCT	GAA	GGA	GCT	TTG	GAC	CGA	TGC	CCC	TGC	GCC	AGT	GGC	CTC	CTA	925
C	Q	P	H	S	H	S	L	V	Y	M	C	K	P	A	F	V	G	S	H	292
TGC	CAG	CCA	CAC	AGC	CAC	AGT	CTG	GTG	TAC	ATG	TGC	AAG	CCA	GCC	TTC	GTG	GGC	AGC	CAT	985
D	H	S	E	E	S	Q	L	P	R	E	A	P	D	E	Y	E	D	V	G	312
GAC	CAC	AGT	GAG	GAG	AGC	CAG	CTG	CCC	AGG	GAG	GCC	CCG	GAT	GAG	TAC	GAA	GAT	GTT	GGC	1045
F	I	G	E	V	R	Q	E	L	E	D	L	E	R	S	L	A	Q	E	M	332
TTC	ATA	GGG	GAA	GTG	CGC	CAG	GAG	CTG	GAA	GAC	CTG	GAG	CGG	AGC	CTA	GCC	CAG	GAG	ATG	1105

FIG. 5B-1

A F E G P A P V E S L G G E E I * 350
GCA TTT GAG GGG CCT GCC CCT GTG GAG TCA CTA GGC GGA GAG GAG GAG ATT TAG 1159
GCCCAGACCCAGCTGAGTCACTGGTAGATGTGCAATAGAAATGGCTAATTATTATTTCCAGGAGTGTCCCCAAGTGTGG 1238
AATGGCCGAGCTCCTTCCAGTAGCTTTTCCCTCTGGCTTGACAAGGTACAGTGCAGTACATTTCTTCCAGCCGCCCTG 1317
CTTCTCTGACTTGGGAAAGACAGGCATGGCGGGTAAGGGCAGCGGTGAGTCGTCCCTCGCTGTTGCTAGAAACGCTGTC 1396
TTGTTCTTCATGGATGGAAAGATTGTTTGAAGGAGAGGATGGGAAGGGGTGAAGTCTGCTCATGATGGATTTGGGGGA 1475
TACAGGGAGGAGGATGCCCTTGCCAGACGCTGGACTTGGCAAAATGTAAACCTTTGCTTTTGTCTTTGCGCCGCTCCCAT 1554
GGGCTGAGGCAGTGGCTACACAAGAGCTATGCTGCTCTGTGGCCTCCCACATATTCAATCCCTGTGTGTTTCAGCTCCTACC 1633
TCACTGTCAGCACAGCCCTTCATAGCCACGCCCCCTCTTGCTCACCACAGCCCTAGGAGGGGACCAGAGGGGACTTCTCT 1712
CAGAGCCCCATGCTCTCTCTCAACCCCATACCAGCCTCTGTGCCAGCGACAGTCCCTTCCAAATGGAGGGAGTGAAAT 1791
CCTTTGGTTTAATTATTTCTCTTCAAGGCACGCCCTGCCACTAAGGTCAGGCTGACTTGCAATGTCCCTCTAAACGTTTCG 1870
TAGCAGTGTGGTGGACACTGTCTTCCACCGACTGCTTCAATACCTCTGAAAGCCAGTGCTCGGAGTGCAGTTCGTGTAA 1949
ATTAAATTTGCAGGAAGTATACTTGGCTAATTGTAGGGCTAGGATTGTGAATGAAATTTGCAAAGTCGCTTAGCAACAAT 2028
GGAAAGCCTTTCTCAGTCACACCGAGAAGTCACAACCAAGCCAGGTTGTGTAGAGTACAGCTGTGACATACAGACAGAA 2107

FIG. 5B-2

GAAGGCTGGGCTGGATGTCAGGCCCTCAGATGACGGTTTCAGGTGCCAGGAACCTATTACCAATTCTGTATCTATCCAGAGT 2186

TATTAAATTTGAAAAGTTGCACACATTTGTATAAGCATGCCTTTCTCCTGAGTTTAAATTATATGTATACACAAACATG 2265

TGGCCCTCAAAGATCATGCACAAACCACACTCTTTTGCTAATTCTTTGGACTTTTCTCTTTGATTTTCAATAAATACAAA 2344

TCCCCCTTCATGCAAAAAAAAAAAAAAGGGGGCCGC 2381

FIG. 6A

[illegible]

19/29

FIG. 6B

hdkk-1	240	VQICLACRKR	RKRCMRHAMC	CPGNYCKNGI	CVS..SDQNH	F..RGEIET	ITESFGN.DH
mdkk-1		VQICLACRKR	RKRCMTHAMC	CPGNYCKNGI	OMP..SDHSH	FP.RGEIEES	I IENLGN.DH
xdkk-1		.LVCLACRKR	RKRCLRDAMC	CTGNYCSNGI	CVPVEQDER	FHQGYLEET	I LENYNNADH
hdkk-2		...CMVCRRK	KRCHRDGMC	CPSTRCNGI	CIPV.TESIL	TPHIPALDGT	RHRD.RNHGH
hdkk-3		QYTCQPCRQ	RMLCTRDSEC	CGDQLCVWGH	CTKMAT....
mdkk-3		KYTCQPCRQ	QMLCTRDSEC	CGDQLCAWGH	CTQKAT....
cdkk-3		EYKCQPKTQ	HTHCSRDEVC	CGDQLCVWGE	CRKATS....
hdkk-4		..FCATCRGL	RRRCQRDAMC	CPGTLCVNDV	CTTME.DATP	ILERQLDEQD	GTHAEGTTGH
hdkk-1	241	STL..DGYSR	RTTLSSKMYH	TKQEGSVCL	RSSDCASGLC	CA..RHFWSK	ICKPVLKEGQ
mdkk-1		NAAAGGYPR	RTTLTSKIYH	TKQEGSVCL	RSSDCAAGLC	CA..RHFWSK	ICKPVLKEGQ
xdkk-1		ATM..DTHSK	LTTSPSGMQP	FKGRDGDVCL	RSTDCAPGLC	CA..RHFWSK	ICKPVLDEGQ
hdkk-2		YSNHDLGWQN	LGRPHTKMSH	IKGHEGDVCL	RSSDCIEGFC	CA..RHFWTK	ICKPVLHQGE
hdkk-3	RGSNGTICD	NQDRCQPGLC	CAFQRGLLFP	VCTPLPVEGE
mdkk-3	KGGNGTICD	NQDRCQPGLC	CAFQRGLLFP	VCTPLPVEGE
cdkk-3	RGNGTICE	NQHDCNPGTC	CAFQKELLFP	VCTPLPEEGE
hdkk-4		PV..QENQPK	RKPSIKKSQG	RKGQEGESCL	RTFDCGPGLC	CA..RHFWTK	ICKPVLLEGQ
hdkk-1	301	VC...TKHRR	KG.SHGLE..	.IFQRCYCGE	GLSCRIQK.D	HHQASNSSRL	HTCQRH~~~~
mdkk-1		VC...TKHRR	KG.SHGLE..	.IFQRCYCGE	GLACRIQK.D	HHQASNSSRL	HTCQRH~~~~
xdkk-1		VC...TKHRR	KG.SHGLE..	.IFQRCYCGA	GLSCRLQKGE	FTTVPKTSRL	HTCQRH~~~~
hdkk-2		VC...TKQRK	KG.SHGLE..	.IFQRCDCAK	GLSCKVWKD.	.ATYSSKARL	HVCQKI~~~~
hdkk-3		LCHDPASRL	LLDLITWELEPD	GALDRCPCAS	GLLCQPH.SH	SLVYVCKPTF	VGSRDQDGE.
mdkk-3		LCHDPTSQ	LLDLITWELEPE	GALDRCPCAS	GLLCQPH.SH	SLVYMCKPAF	VGSHDHSEE.
cdkk-3		PHCDPSNR	LLDLITWELEPD	GVLERCPCAS	GLICQPQSSH	STTSVCELSS	NETRKNKED
hdkk-4		VC...SRRGH	KDTAQAPE..	.IFQRCDCGP	GLLCRSQ LTS	NRQH...ARL	RVCQKIEKL~

FIG. 6C

361	424
hdkk-1	hdkk-1
mdkk-1	mdkk-1
xdkk-1	xdkk-1
hdkk-2	hdkk-2
hdkk-3	hdkk-3
mdkk-3	mdkk-3
cdkk-3	cdkk-3
hdkk-4	hdkk-4

FIG. 7A

	M	1
CTCGAGGCCAAAATTTCGGACGAGGCCGGCTGTGGTCTAGCATAAAGCGGAGCCAGAAAGGGCGGGGT	ATG	77
G E A S P P A P A R R H L L V L L L L L L	L L L	21
GGA GAA GCC TCC CCA CCT GCC CCC GCA AGG CGG CAT CTG CTG GTC CTG CTG CTC CTC	CTC	137
S T L V I P S A A A P I H D A D A Q E S	S	41
TCT ACC CTG GTG ATC CCC TCC GCT GCA GCT CCT ATC CAT GAT GCT GAC GCC CAA GAG AGC	AGC	197
S L G L T G L Q S L L Q G F S R L F L K	K	61
TCC TTG GGT CTC ACA GGC CTC CAG AGC CTA CTC CAA GGC TTC AGC CGA CTT TTC CTG AAA	AAA	257
G N L L R G I D S L F S A P M D F R G L	L	81
GGT AAC CTG CTT CGG GGC ATA GAC AGC TTA TTC TCT GCC CCC ATG GAC TTC CGG GGC CTC	CTC	317
P G N Y H K E E N Q E H Q L G N N T L S	S	101
CCT GGG AAC TAC CAC AAA GAG GAG AAC CAG GAG CAC CAG CTG GGG AAC AAC ACC CTC TCC	TCC	377
S H L Q I D K M T D N K T G E V L I S E	E	121
AGC CAC CTC CAG ATC GAC AAG ATG ACC GAC AAC AAG ACA GGA GAG GTG CTG ATC TCC GAG	GAG	437
N V V A S I Q P A E G S F E G D L K V P	P	141
AAT GTG GTG GCA TCC ATT CAA CCA CCG GAG GGG AGC TTC GAG GGT GAT TTG AAG GTA CCC	CCC	497
R M E E K E A L V P I Q K A T D S F H T	T	161
AGG ATG GAG GAG AAG GAG GCC CTG GTA CCC ATC CAG AAG GCC ACG GAC AGC TTC CAC ACA	ACA	557
E L H P R V A F W I I K L P R R S H Q	Q	181
GAA CTC CAT CCC CGG GTG GCC TTC TGG ATC ATT AAG CTG CCA CGG CGG AGG TCC CAC CAG	CAG	617

TACATCTAAAAAATAAAAAAATAATGGCGCGCGC

	M	C	R	L	R	5
GAAATTCGCACGAGCGCAGAAGCGCGAATGAAGGCAAGCTCCACCACCTGCA ATG TGT CGA CTG AGG						71
V L L L L L L P L A F V S S S A L P I H D						25
GTC TTG CTG CTG CTG CCC TTG GCC TTC GTG TCC TCC TCT GCT CTC CCC ATC CAT GAT						131
V D S Q Q N T S G F L G L Q R L L Q S F						45
GTC GAC TCT CAG CAG AAC ACC TCC GGG TTC CTG GGC CTT CAG AGG CTT CTC CAA AGC TTT						161
S R L F L K N D L L R D L D N F F S S P						65
AGT CGA CTG TTC CTA AAA AAT GAC CTG CTA CGA GAC CTG GAC AAC TTC TTC TCC TCC CCC						251
M D F R D L P R N F H Q E E N Q E H R M						85
AATG GAC TTC CGA GAC CTT CCT AGG AAC TTC CAT CAG GAA GAG AAC CAG GAG CAC AGA ATG						311
G N H T L S S H L Q I D K V T D N Q T G						105
GGC AAC CAT ACC CTC TCC AGC CAC CTA CAG ATA GAC AAG GTG ACT GAC AAC CAG ACA GGG						371
E V H I S E K V E A S I E P E R N P E G						125
GAG GTG CAC ATC TCG GAG AAA GTC GAG GCC TCC ATT GAG CCA GAA CGG AAC CCG GAA GGG						431
D W K V P K V E A K E P P V P V Q K V T						145
GAC TGG AAG GTT CCC AAA GTA GAA GCA AAA GAG CCC CCG GTG CCT GTG CAG AAG GTC ACC						491
D S L H P E P R Q V A F W I M K M P R R						165
GAC AGC TTG CAC CCA GAG CCC CGG CAG GTG GCT TTC TGG ATC ATG AAG ATG CCA AGG CGG						551

FIG. 8B

```

R   T   Q   P   D   V   Q   D   G   G   R   W   L   I   E   K   R   H   R   M   185
AGG ACC CAG CCC GAT GTC CAG GAT GGA GGC CGC TGG CTC ATA GAA AAG CGA CAT CGC ATG 611

Q   A   I   R   D   G   L   R   G   G   A   R   E   D   S   L   E   D   G   V   205
CAG GCC ATC CGG GAT GGG CTC CGT GGA GGC GCC GCG CGT GAG GAC AGC CTG GAG GAT GGG GTC 671

H   I   P   Q   H   A   K   L   P   V   R   R   K   T   H   F   L   Y   I   L   R   225
CAT ATC CCC CAA CAC GCC AAG CTG CCT GTC AGA AAG ACA CAC CAC TTT CTC TAC ATC CTC AGG 731

P   S   Q   Q   L   *
CCA TCC CAA CAG CTG TAA GTGGGACCAGATGTCCACACCCCTACCCCAACCATATGGAAATAAAGGTTTC 231
TTACATCTAAAAAAAAAAAAAAAAAAAAA 805
TTACATCTAAAAAAAAAAAAAAAAAAAAA 835

```

FIG. 9




FIG. 10A

1	60
hsoggy	MGEASPPAPA RRHL.LVLLL LLSTLVIPSA AAPIHDAQAQ ESSLG..... LTGLQSL
msoggy	~~~~~ MCRL.RVLLL LLPLAFVSSS ALPIHDVDSQ QNTSG..... FLGLQRL
hdkk-3	~~~~~ MQRLGATLLC LLAAAVPTA PAPAPTATSA PVKPGPALSY PQEEATLNEM
mdkk-3	~~~~~ MQRLLGGILLC TLLAAAVPTA PAPSPVTWT PAEPGPALNY PQEEATLNEM *

61	120
hsoggy	LQFSRLE.. LKGNLLRGID SL..... FSAPMDFRGL PGNYHKEENQ EHQLGNNTLS
msoggy	LQFSRLE.. LKNLLRLDLD NF..... FSSPMDFRDL PRNFHQEENQ EHRMGNHTLS
hdkk-3	FREVEELMED TQHKLRSAVE EMEAEAAAK ASSEVNLANL PPSYHNETNT DTKVGNNTIH
mdkk-3	FREVEELMED TQHKLRSAVE EMEAEAAAK TSSEVNLASL PPNYHNETST ETRVGNNTVH *


121	180
hsoggy	SHLQIDKMTD NKTGEVLISE NVVASIQPAE GSFEGDLKVP RMEKEALVP IQKATDSFHT
msoggy	SHLQIDKMTD NKTGEVLISE KVEASIEP.E RNPEGDWKVP KVEAKEPPVP VQKVTDLSHP
hdkk-3	VHREIHKITN NQTGQMFSE TVITSVGDEE GR..... RSHECII. DEDCGP
mdkk-3	VHREIHKITN NQTGQMFSE TVITSVGDEE GK..... RSHECII. DEDCGP *

181	240
hsoggy	ELHPR.VAFW IIKLPRRRSH ... QDALEG GHWLSEKRHR LQAIRDGLRK .. GTHKD..
msoggy	E.. PRQVAFW IMKMPRRRTQ PDVQDG GRWLIEKRHR MQAIRDGLRG .. GARED..
hdkk-3	SMYCOFASFQ YTCQPCRGQR MLCTRDSECC GDQLCVWGHC TKMATRGSNG TICTDNQRDCQ
mdkk-3	TRYCQFSSFK YTCQPCRDQQ MLCTRDSECC GDQLCAWGHC TQKATKGNG TICTDNQRDCQ *

FIG. 10B

hsoggy	241VLEEG TESSHSRLS PRKTHLLYIL RPSRQL	300
msoggy	SLEDG VHIPQHAKLP VRKTHFLYIL RPSQQQL	
hdkk-3		PGLCCAFQRG LLFPVCTPLP V.EGELCH.. DPASRLLDLI TWELEPDGAL DRCPCASGLL	
mdkk-3		PGLCCAFQRG LLFPVCTPLP V.EGELCH.. DPTSQQLDLI TWELEPEGAL DRCPCASGLL	
		★	★
hdkk-3	301	CQPHSHSLVY VCKPTFVGSR DQDGEILLPR EVPDEYEVGS FMEEVVRQELE DLESLITEEM	360
mdkk-3		CQPHSHSLVY MCKPAFVGSH DHSEESQLPR EAPDEYEDVG FIGEVVRQELE DLESLAQEM	
hdkk-3	361	ALGEPAAAAA ALLGGEEI~	379
mdkk-3		AFEGPAPVES ..LGEEIEI	

FIG. 12



- 1 -

SEQUENCE LISTING

<110> MILLENNIUM PHARMACEUTICALS, INC.

<120> HUMAN DICKKOPF-RELATED PROTEIN AND NUCLEIC ACID
MOLECULES AND USES THEREFOR

<130> MNI-108CP2PC

<140> 09/263,022

<141> 1999-03-05

<160> 38

<170> PatentIn Ver. 2.0

<210> 1

<211> 2479

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (38)..(1087)

<220>

<223> 'n' at position 1146 may be any nucleotide

<400> 1

```

ggcacgaggg ggcggcggct gcgggcgcag agcggag atg cag cgg ctt ggg gcc 55
                                   Met Gln Arg Leu Gly Ala
                                   1           5

acc ctg ctg tgc ctg ctg ctg gcg gcg gcg gtc ccc acg gcc ccc gcg 103
Thr Leu Leu Cys Leu Leu Leu Ala Ala Ala Val Pro Thr Ala Pro Ala
      10           15           20

ccc gct ccg acg gcg acc tcg gct cca gtc aag ccc ggc ccg gct ctc 151
Pro Ala Pro Thr Ala Thr Ser Ala Pro Val Lys Pro Gly Pro Ala Leu
      25           30           35

agc tac ccg cag gag gag gcc acc ctc aat gag atg ttc cgc gag gtt 199
Ser Tyr Pro Gln Glu Glu Ala Thr Leu Asn Glu Met Phe Arg Glu Val
      40           45           50

gag gaa ctg atg gag gac acg cag cac aaa ttg cgc agc gcg gtg gaa 247
Glu Glu Leu Met Glu Asp Thr Gln His Lys Leu Arg Ser Ala Val Glu
      55           60           65           70

gag atg gag gca gaa gaa gct gct gct aaa gca tca tca gaa gtg aac 295
Glu Met Glu Ala Glu Glu Ala Ala Ala Lys Ala Ser Ser Glu Val Asn
      75           80           85

ctg gca aac tta cct ccc agc tat cac aat gag acc aac aca gac acg 343
Leu Ala Asn Leu Pro Pro Ser Tyr His Asn Glu Thr Asn Thr Asp Thr
      90           95           100

aac gtt gga aat aat acc atc cat gtg cac cga gaa att cac aag ata 391

```

- 2 -

Asn Val Gly Asn Asn Thr Ile His Val His Arg Glu Ile His Lys Ile	
105 110 115	
acc aac aac cag act gga caa atg gtc ttt tca gag aca gtt atc aca	439
Thr Asn Asn Gln Thr Gly Gln Met Val Phe Ser Glu Thr Val Ile Thr	
120 125 130	
tct gtg gga gac gaa gaa ggc aga agg agc cac gag tgc atc atc gac	487
Ser Val Gly Asp Glu Glu Gly Arg Arg Ser His Glu Cys Ile Ile Asp	
135 140 145 150	
gag gac tgt ggg ccc agc atg tac tgc cag ttt gcc agc ttc cag tac	535
Glu Asp Cys Gly Pro Ser Met Tyr Cys Gln Phe Ala Ser Phe Gln Tyr	
155 160 165	
acc tgc cag cca tgc cgg ggc cag agg atg ctc tgc acc cgg gac agt	583
Thr Cys Gln Pro Cys Arg Gly Gln Arg Met Leu Cys Thr Arg Asp Ser	
170 175 180	
gag tgc tgt gga gac cag ctg tgt gtc tgg ggt cac tgc acc aaa atg	631
Glu Cys Cys Gly Asp Gln Leu Cys Val Trp Gly His Cys Thr Lys Met	
185 190 195	
gcc acc agg ggc agc aat ggg acc atc tgt gac aac cag agg gac tgc	679
Ala Thr Arg Gly Ser Asn Gly Thr Ile Cys Asp Asn Gln Arg Asp Cys	
200 205 210	
cag ccg ggg ctg tgc tgt gcc ttc cag aga ggc ctg ctg ttc cct gtg	727
Gln Pro Gly Leu Cys Cys Ala Phe Gln Arg Gly Leu Leu Phe Pro Val	
215 220 225 230	
tgc aca ccc ctg ccc gtg gag ggc gag ctt tgc cat gac ccc gcc agc	775
Cys Thr Pro Leu Pro Val Glu Gly Glu Leu Cys His Asp Pro Ala Ser	
235 240 245	
cgg ctt ctg gac ctc atc acc tgg gag cta gag cct gat gga gcc ttg	823
Arg Leu Leu Asp Leu Ile Thr Trp Glu Leu Glu Pro Asp Gly Ala Leu	
250 255 260	
gac cga tgc cct tgt gcc agt ggc ctc ctc tgc cag ccc cac agc cac	871
Asp Arg Cys Pro Cys Ala Ser Gly Leu Leu Cys Gln Pro His Ser His	
265 270 275	
agc ctg gtg tat gtg tgc aag ccg acc ttc gtg ggg agc cgt gac caa	919
Ser Leu Val Tyr Val Cys Lys Pro Thr Phe Val Gly Ser Arg Asp Gln	
280 285 290	
gat ggg gag atc ctg ctg ccc aga gag gtc ccc gat gag tat gaa gtt	967
Asp Gly Glu Ile Leu Leu Pro Arg Glu Val Pro Asp Glu Tyr Glu Val	
295 300 305 310	
ggc agc ttc atg gag gag gtg cgc cag gag ctg gag gac ctg gag agg	1015
Gly Ser Phe Met Glu Glu Val Arg Gln Glu Leu Glu Asp Leu Glu Arg	
315 320 325	
agc ctg act gaa gag atg gcg ctg agg gag cct gcg gct gcc gcc gct	1063
Ser Leu Thr Glu Glu Met Ala Leu Arg Glu Pro Ala Ala Ala Ala Ala	
330 335 340	

gca ctg ctg gga agg gaa gag att tagatctgga ccaggctgtg ggtagatgtg 1117
Ala Leu Leu Gly Arg Glu Glu Ile
345 350

```
<210> 2
<211> 350
<212> PRT
<213> Homo sapiens
```

<400> 2
Met Gln Arg Leu Gly Ala Thr Leu Leu Cys Leu Leu Leu Ala Ala Ala
1 5 10 15

- 4 -

Val Pro Thr Ala Pro Ala Pro Ala Pro Thr Ala Thr Ser Ala Pro Val
 20 25 30
 Lys Pro Gly Pro Ala Leu Ser Tyr Pro Gln Glu Glu Ala Thr Leu Asn
 35 40 45
 Glu Met Phe Arg Glu Val Glu Glu Leu Met Glu Asp Thr Gln His Lys
 50 55 60
 Leu Arg Ser Ala Val Glu Glu Met Glu Ala Glu Glu Ala Ala Ala Lys
 65 70 75 80
 Ala Ser Ser Glu Val Asn Leu Ala Asn Leu Pro Pro Ser Tyr His Asn
 85 90 95
 Glu Thr Asn Thr Asp Thr Asn Val Gly Asn Asn Thr Ile His Val His
 100 105 110
 Arg Glu Ile His Lys Ile Thr Asn Asn Gln Thr Gly Gln Met Val Phe
 115 120 125
 Ser Glu Thr Val Ile Thr Ser Val Gly Asp Glu Glu Gly Arg Arg Ser
 130 135 140
 His Glu Cys Ile Ile Asp Glu Asp Cys Gly Pro Ser Met Tyr Cys Gln
 145 150 155 160
 Phe Ala Ser Phe Gln Tyr Thr Cys Gln Pro Cys Arg Gly Gln Arg Met
 165 170 175
 Leu Cys Thr Arg Asp Ser Glu Cys Cys Gly Asp Gln Leu Cys Val Trp
 180 185 190
 Gly His Cys Thr Lys Met Ala Thr Arg Gly Ser Asn Gly Thr Ile Cys
 195 200 205
 Asp Asn Gln Arg Asp Cys Gln Pro Gly Leu Cys Cys Ala Phe Gln Arg
 210 215 220
 Gly Leu Leu Phe Pro Val Cys Thr Pro Leu Pro Val Glu Gly Glu Leu
 225 230 235 240
 Cys His Asp Pro Ala Ser Arg Leu Leu Asp Leu Ile Thr Trp Glu Leu
 245 250 255
 Glu Pro Asp Gly Ala Leu Asp Arg Cys Pro Cys Ala Ser Gly Leu Leu
 260 265 270
 Cys Gln Pro His Ser His Ser Leu Val Tyr Val Cys Lys Pro Thr Phe
 275 280 285
 Val Gly Ser Arg Asp Gln Asp Gly Glu Ile Leu Leu Pro Arg Glu Val
 290 295 300
 Pro Asp Glu Tyr Glu Val Gly Ser Phe Met Glu Glu Val Arg Gln Glu
 305 310 315 320
 Leu Glu Asp Leu Glu Arg Ser Leu Thr Glu Glu Met Ala Leu Arg Glu

- 5 -

325	330	335
Pro Ala Ala Ala Ala Ala Ala Leu Leu Gly Arg Glu Glu Ile		
340	345	350

<210> 3
 <211> 1050
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> CDS
 <222> (1)..(1050)

 <400> 3

atg cag cgg ctt ggg gcc acc ctg ctg tgc ctg ctg ctg gcg gcg gcg	48
Met Gln Arg Leu Gly Ala Thr Leu Leu Cys Leu Leu Leu Ala Ala Ala	
1 5 10 15	
gtc ccc acg gcc ccc gcg ccc gct ccg acg gcg acc tcg gct cca gtc	96
Val Pro Thr Ala Pro Ala Pro Ala Pro Thr Ala Thr Ser Ala Pro Val	
20 25 30	
aag ccc ggc ccg gct ctc agc tac ccg cag gag gag gcc acc ctc aat	144
Lys Pro Gly Pro Ala Leu Ser Tyr Pro Gln Glu Glu Ala Thr Leu Asn	
35 40 45	
gag atg ttc cgc gag gtt gag gaa ctg atg gag gac acg cag cac aaa	192
Glu Met Phe Arg Glu Val Glu Glu Leu Met Glu Asp Thr Gln His Lys	
50 55 60	
ttg cgc agc gcg gtg gaa gag atg gag gca gaa gaa gct gct gct aaa	240
Leu Arg Ser Ala Val Glu Glu Met Glu Ala Glu Glu Ala Ala Ala Lys	
65 70 75 80	
gca tca tca gaa gtg aac ctg gca aac tta cct ccc agc tat cac aat	288
Ala Ser Ser Glu Val Asn Leu Ala Asn Leu Pro Pro Ser Tyr His Asn	
85 90 95	
gag acc aac aca gac acg aac gtt gga aat aat acc atc cat gtg cac	336
Glu Thr Asn Thr Asp Thr Asn Val Gly Asn Asn Thr Ile His Val His	
100 105 110	
cga gaa att cac aag ata acc aac aac cag act gga caa atg gtc ttt	384
Arg Glu Ile His Lys Ile Thr Asn Asn Gln Thr Gly Gln Met Val Phe	
115 120 125	
tca gag aca gtt atc aca tct gtg gga gac gaa gaa ggc aga agg agc	432
Ser Glu Thr Val Ile Thr Ser Val Gly Asp Glu Glu Gly Arg Arg Ser	
130 135 140	
cac gag tgc atc atc gac gag gac tgt ggg ccc agc atg tac tgc cag	480
His Glu Cys Ile Ile Asp Glu Asp Cys Gly Pro Ser Met Tyr Cys Gln	
145 150 155 160	
ttt gcc agc ttc cag tac acc tgc cag cca tgc cgg ggc cag agg atg	528
Phe Ala Ser Phe Gln Tyr Thr Cys Gln Pro Cys Arg Gly Gln Arg Met	
165 170 175	

- 6 -

```

ctc tgc acc cgg gac agt gag tgc tgt gga gac cag ctg tgt gtc tgg 576
Leu Cys Thr Arg Asp Ser Glu Cys Cys Gly Asp Gln Leu Cys Val Trp
180 185 190

ggt cac tgc acc aaa atg gcc acc agg ggc agc aat ggg acc atc tgt 624
Gly His Cys Thr Lys Met Ala Thr Arg Gly Ser Asn Gly Thr Ile Cys
195 200 205

gac aac cag agg gac tgc cag ccg ggg ctg tgc tgt gcc ttc cag aga 672
Asp Asn Gln Arg Asp Cys Gln Pro Gly Leu Cys Cys Ala Phe Gln Arg
210 215 220

ggc ctg ctg ttc cct gtg tgc aca ccc ctg ccc gtg gag ggc gag ctt 720
Gly Leu Leu Phe Pro Val Cys Thr Pro Leu Pro Val Glu Gly Glu Leu
225 230 235 240

tgc cat gac ccc gcc agc cgg ctt ctg gac ctc atc acc tgg gag cta 768
Cys His Asp Pro Ala Ser Arg Leu Leu Asp Leu Ile Thr Trp Glu Leu
245 250 255

gag cct gat gga gcc ttg gac cga tgc cct tgt gcc agt ggc ctc ctc 816
Glu Pro Asp Gly Ala Leu Asp Arg Cys Pro Cys Ala Ser Gly Leu Leu
260 265 270

tgc cag ccc cac agc cac agc ctg gtg tat gtg tgc aag ccg acc ttc 864
Cys Gln Pro His Ser His Ser Leu Val Tyr Val Cys Lys Pro Thr Phe
275 280 285

gtg ggg agc cgt gac caa gat ggg gag atc ctg ctg ccc aga gag gtc 912
Val Gly Ser Arg Asp Gln Asp Gly Glu Ile Leu Leu Pro Arg Glu Val
290 295 300

ccc gat gag tat gaa gtt ggc agc ttc atg gag gag gtg cgc cag gag 960
Pro Asp Glu Tyr Glu Val Gly Ser Phe Met Glu Glu Val Arg Gln Glu
305 310 315 320

ctg gag gac ctg gag agg agc ctg act gaa gag atg gcg ctg agg gag 1008
Leu Glu Asp Leu Glu Arg Ser Leu Thr Glu Glu Met Ala Leu Arg Glu
325 330 335

cct gcg gct gcc gcc gct gca ctg ctg gga agg gaa gag att 1050
Pro Ala Ala Ala Ala Ala Ala Leu Leu Gly Arg Glu Glu Ile
340 345 350

```

<210> 4

<211> 848

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (125)..(796)

<400> 4

gaattcggca cgagagacga cgtgctgagc tgccagctta gtggaagctc tgctctgggt 60

ggagagcagc ctcgctttgg tgacgcacag tgctgggacc ctccaggagc cccgggattg 120

- 7 -

```

aagg atg gtg gcg gcc gtc ctg ctg ggg ctg agc tgg ctc tgc tct ccc 169
    Met Val Ala Ala Val Leu Leu Gly Leu Ser Trp Leu Cys Ser Pro
      1             5             10             15

ctg gga gct ctg gtc ctg gac ttc aac aac atc agg agc tct gct gac 217
Leu Gly Ala Leu Val Leu Asp Phe Asn Asn Ile Arg Ser Ser Ala Asp
      20             25             30

ctg cat ggg gcc cgg aag ggc tca cag tgc ctg tct gac acg gac tgc 265
Leu His Gly Ala Arg Lys Gly Ser Gln Cys Leu Ser Asp Thr Asp Cys
      35             40             45

aat acc aga aag ttc tgc ctc cag ccc cgc gat gag aag ccg ttc tgt 313
Asn Thr Arg Lys Phe Cys Leu Gln Pro Arg Asp Glu Lys Pro Phe Cys
      50             55             60

gct aca tgt cgt ggg ttg cgg agg agg tgc cag cga gat gcc atg tgc 361
Ala Thr Cys Arg Gly Leu Arg Arg Arg Cys Gln Arg Asp Ala Met Cys
      65             70             75

tgc cct ggg aca ctc tgt gtg aac gat gtt tgt act acg atg gaa gat 409
Cys Pro Gly Thr Leu Cys Val Asn Asp Val Cys Thr Thr Met Glu Asp
      80             85             90             95

gca acc cca ata tta gaa agg cag ctt gat gag caa gat ggc aca cat 457
Ala Thr Pro Ile Leu Glu Arg Gln Leu Asp Glu Gln Asp Gly Thr His
      100            105            110

gca gaa gga aca act ggg cac cca gtc cag gaa aac caa ccc aaa agg 505
Ala Glu Gly Thr Thr Gly His Pro Val Gln Glu Asn Gln Pro Lys Arg
      115            120            125

aag cca agt att aag aaa tca caa ggc agg aag gga caa gag gga gaa 553
Lys Pro Ser Ile Lys Lys Ser Gln Gly Arg Lys Gly Gln Glu Gly Glu
      130            135            140

agt tgt ctg aga act ttt gac tgt ggc cct gga ctt tgc tgt gct cgt 601
Ser Cys Leu Arg Thr Phe Asp Cys Gly Pro Gly Leu Cys Cys Ala Arg
      145            150            155

cat ttt tgg acg aaa att tgt aag cca gtc ctt ttg gag gga cag gtc 649
His Phe Trp Thr Lys Ile Cys Lys Pro Val Leu Leu Glu Gly Gln Val
      160            165            170            175

tgc tcc aga aga ggg cat aaa gac act gct caa gct cca gaa atc ttc 697
Cys Ser Arg Arg Gly His Lys Asp Thr Ala Gln Ala Pro Glu Ile Phe
      180            185            190

cag cgt tgc gac tgt ggc cct gga cta ctg tgt cga agc caa ttg acc 745
Gln Arg Cys Asp Cys Gly Pro Gly Leu Leu Cys Arg Ser Gln Leu Thr
      195            200            205

agc aat cgg cag cat gct cga tta aga gta tgc caa aaa ata gaa aag 793
Ser Asn Arg Gln His Ala Arg Leu Arg Val Cys Gln Lys Ile Glu Lys
      210            215            220

cta taaatatttc aaaataaaga agaatccaca ttgcaaaaaa aaaaaaaaaa aa 848
Leu

```

- 8 -

<210> 5
 <211> 224
 <212> PRT
 <213> Homo sapiens

<400> 5
 Met Val Ala Ala Val Leu Leu Gly Leu Ser Trp Leu Cys Ser Pro Leu
 1 5 10 15
 Gly Ala Leu Val Leu Asp Phe Asn Asn Ile Arg Ser Ser Ala Asp Leu
 20 25 30
 His Gly Ala Arg Lys Gly Ser Gln Cys Leu Ser Asp Thr Asp Cys Asn
 35 40 45
 Thr Arg Lys Phe Cys Leu Gln Pro Arg Asp Glu Lys Pro Phe Cys Ala
 50 55 60
 Thr Cys Arg Gly Leu Arg Arg Arg Cys Gln Arg Asp Ala Met Cys Cys
 65 70 75 80
 Pro Gly Thr Leu Cys Val Asn Asp Val Cys Thr Thr Met Glu Asp Ala
 85 90 95
 Thr Pro Ile Leu Glu Arg Gln Leu Asp Glu Gln Asp Gly Thr His Ala
 100 105 110
 Glu Gly Thr Thr Gly His Pro Val Gln Glu Asn Gln Pro Lys Arg Lys
 115 120 125
 Pro Ser Ile Lys Lys Ser Gln Gly Arg Lys Gly Gln Glu Gly Glu Ser
 130 135 140
 Cys Leu Arg Thr Phe Asp Cys Gly Pro Gly Leu Cys Cys Ala Arg His
 145 150 155 160
 Phe Trp Thr Lys Ile Cys Lys Pro Val Leu Leu Glu Gly Gln Val Cys
 165 170 175
 Ser Arg Arg Gly His Lys Asp Thr Ala Gln Ala Pro Glu Ile Phe Gln
 180 185 190
 Arg Cys Asp Cys Gly Pro Gly Leu Leu Cys Arg Ser Gln Leu Thr Ser
 195 200 205
 Asn Arg Gln His Ala Arg Leu Arg Val Cys Gln Lys Ile Glu Lys Leu
 210 215 220

<210> 6
 <211> 672
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(672)

- 9 -

<400> 6
 atg gtg gcg gcc gtc ctg ctg ggg ctg agc tgg ctc tgc tct ccc ctg 48
 Met Val Ala Ala Val Leu Leu Gly Leu Ser Trp Leu Cys Ser Pro Leu
 1 5 10 15

gga gct ctg gtc ctg gac ttc aac aac atc agg agc tct gct gac ctg 96
 Gly Ala Leu Val Leu Asp Phe Asn Asn Ile Arg Ser Ser Ala Asp Leu
 20 25 30

cat ggg gcc cgg aag ggc tca cag tgc ctg tct gac acg gac tgc aat 144
 His Gly Ala Arg Lys Gly Ser Gln Cys Leu Ser Asp Thr Asp Cys Asn
 35 40 45

acc aga aag ttc tgc ctc cag ccc cgc gat gag aag ccg ttc tgt gct 192
 Thr Arg Lys Phe Cys Leu Gln Pro Arg Asp Glu Lys Pro Phe Cys Ala
 50 55 60

aca tgt cgt ggg ttg cgg agg agg tgc cag cga gat gcc atg tgc tgc 240
 Thr Cys Arg Gly Leu Arg Arg Arg Cys Gln Arg Asp Ala Met Cys Cys
 65 70 75 80

cct ggg aca ctc tgt gtg aac gat gtt tgt act acg atg gaa gat gca 288
 Pro Gly Thr Leu Cys Val Asn Asp Val Cys Thr Thr Met Glu Asp Ala
 85 90 95

acc cca ata tta gaa agg cag ctt gat gag caa gat ggc aca cat gca 336
 Thr Pro Ile Leu Glu Arg Gln Leu Asp Glu Gln Asp Gly Thr His Ala
 100 105 110

gaa gga aca act ggg cac cca gtc cag gaa aac caa ccc aaa agg aag 384
 Glu Gly Thr Thr Gly His Pro Val Gln Glu Asn Gln Pro Lys Arg Lys
 115 120 125

cca agt att aag aaa tca caa ggc agg aag gga caa gag gga gaa agt 432
 Pro Ser Ile Lys Lys Ser Gln Gly Arg Lys Gly Gln Glu Gly Glu Ser
 130 135 140

tgt ctg aga act ttt gac tgt ggc cct gga ctt tgc tgt gct cgt cat 480
 Cys Leu Arg Thr Phe Asp Cys Gly Pro Gly Leu Cys Cys Ala Arg His
 145 150 155 160

ttt tgg acg aaa att tgt aag cca gtc ctt ttg gag gga cag gtc tgc 528
 Phe Trp Thr Lys Ile Cys Lys Pro Val Leu Leu Glu Gly Gln Val Cys
 165 170 175

tcc aga aga ggg cat aaa gac act gct caa gct cca gaa atc ttc cag 576
 Ser Arg Arg Gly His Lys Asp Thr Ala Gln Ala Pro Glu Ile Phe Gln
 180 185 190

cgt tgc gac tgt ggc cct gga cta ctg tgt cga agc caa ttg acc agc 624
 Arg Cys Asp Cys Gly Pro Gly Leu Leu Cys Arg Ser Gln Leu Thr Ser
 195 200 205

aat cgg cag cat gct cga tta aga gta tgc caa aaa ata gaa aag cta 672
 Asn Arg Gln His Ala Arg Leu Arg Val Cys Gln Lys Ile Glu Lys Leu
 210 215 220

- 10 -

```

<210> 7
<211> 1536
<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (109)..(906)

<400> 7
gtcgacccac gcgtccgcgg acgcgtgggc ggcacggttt cgtggggacc caggcttgca 60

aagtgacggt cattttctct ttctttctcc ctcttgagtc cttctgag atg atg gct 117
                                         Met Met Ala
                                         1

ctg ggc gca gcg gga gct acc cgg gtc ttt gtc gcg atg gta gcg gcg 165
Leu Gly Ala Ala Gly Ala Thr Arg Val Phe Val Ala Met Val Ala Ala
      5                      10                      15

gct ctc ggc ggc cac cct ctg ctg gga gtg agc gcc acc ttg aac tcg 213
Ala Leu Gly Gly His Pro Leu Leu Gly Val Ser Ala Thr Leu Asn Ser
      20                      25                      30                      35

gtt ctc aat tcc aac gct atc aag aac ctg ccc cca ccg ctg ggc ggc 261
Val Leu Asn Ser Asn Ala Ile Lys Asn Leu Pro Pro Pro Leu Gly Gly
                      40                      45                      50

gct gcg ggg cac cca ggc tct gca gtc agc gcc gcg ccg gga atc ctg 309
Ala Ala Gly His Pro Gly Ser Ala Val Ser Ala Ala Pro Gly Ile Leu
                      55                      60                      65

tac ccg ggc ggc aat aag tac cag acc att gac aac tac cag ccg tac 357
Tyr Pro Gly Gly Asn Lys Tyr Gln Thr Ile Asp Asn Tyr Gln Pro Tyr
                      70                      75                      80

ccg tgc gca gag gac gag gag tgc ggc act gat gag tac tgc gct agt 405
Pro Cys Ala Glu Asp Glu Glu Cys Gly Thr Asp Glu Tyr Cys Ala Ser
                      85                      90                      95

ccc acc cgc gga ggg gac gca ggc gtg caa atc tgt ctc gcc tgc agg 453
Pro Thr Arg Gly Gly Asp Ala Gly Val Gln Ile Cys Leu Ala Cys Arg
100                      105                      110                      115

aag cgc cga aaa cgc tgc atg cgt cac gct atg tgc tgc ccc ggg aat 501
Lys Arg Arg Lys Arg Cys Met Arg His Ala Met Cys Cys Pro Gly Asn
                      120                      125                      130

tac tgc aaa aat gga ata tgt gtg tct tct gat caa aat cat ttc cga 549
Tyr Cys Lys Asn Gly Ile Cys Val Ser Ser Asp Gln Asn His Phe Arg
                      135                      140                      145

gga gaa att gag gaa acc atc act gaa agc ttt ggt aat gat cat agc 597
Gly Glu Ile Glu Glu Thr Ile Thr Glu Ser Phe Gly Asn Asp His Ser
                      150                      155                      160

acc ttg gat ggg tat tcc aga aga acc acc ttg tct tca aaa atg tat 645
Thr Leu Asp Gly Tyr Ser Arg Arg Thr Thr Leu Ser Ser Lys Met Tyr
165                      170                      175

```

- 11 -

```

cac acc aaa gga caa gaa ggt tct gtt tgt ctc cgg tca tca gac tgt      693
His Thr Lys Gly Gln Glu Gly Ser Val Cys Leu Arg Ser Ser Asp Cys
180                      185                      190                      195

gcc tca gga ttg tgt tgt gct aga cac ttc tgg tcc aag atc tgt aaa      741
Ala Ser Gly Leu Cys Cys Ala Arg His Phe Trp Ser Lys Ile Cys Lys
                200                      205                      210

cct gtc ctg aaa gaa ggt caa gtg tgt acc aag cat agg aga aaa ggc      789
Pro Val Leu Lys Glu Gly Gln Val Cys Thr Lys His Arg Arg Lys Gly
                215                      220                      225

tct cat gga cta gaa ata ttc cag cgt tgt tac tgt gga gaa ggt ctg      837
Ser His Gly Leu Glu Ile Phe Gln Arg Cys Tyr Cys Gly Glu Gly Leu
                230                      235                      240

tct tgc cgg ata cag aaa gat cac cat caa gcc agt aat tct tct agg      885
Ser Cys Arg Ile Gln Lys Asp His His Gln Ala Ser Asn Ser Ser Arg
                245                      250                      255

ctt cac act tgt cag aga cac taaaccagct atccaaatgc agtgaactcc      936
Leu His Thr Cys Gln Arg His
260                      265

ttttatataa tagatgctat gaaaaccttt tatgaccttc atcaactcaa tcctaaggat 996

atacaagttc tgtggtttca gttaagcatt ccaataacac cttccaaaaa cctggagtg 1056

aagagctttg tttctttatg gaactcccct gtgattgcag taaattactg tattgtaaat 1116

tctcagtgtg gcacttacct gtaaagcaaa tgaaactttt aattattttt ctaaagggtgc 1176

tgcactgcct atttttcctc ttgttatgta aatttttgta cacattgatt gttatcttga 1236

ctgacaaaata ttctatattg aactgaagta aatcatttca gcttatagtt cttaaaagca 1296

taacccttta ccccatTTaa ttctagagtc tagaacgcaa ggatctcttg gaatgacaaa 1356

tgataggtac ctaaaatgta acatgaaaat actagcttat tttctgaaat gtactatctt 1416

aatgcttaaa ttatatttcc ctttaggctg tgatagtttt tgaaataaaa tttaacattt 1476

aatatcatga aatgttataa gtagacataa aaaaaaaaaa aaaaaaaaaa gggcgggccgc 1536

<210> 8
<211> 266
<212> PRT
<213> Homo sapiens

<400> 8
Met Met Ala Leu Gly Ala Ala Gly Ala Thr Arg Val Phe Val Ala Met
 1              5              10              15

Val Ala Ala Ala Leu Gly Gly His Pro Leu Leu Gly Val Ser Ala Thr
                20              25              30

Leu Asn Ser Val Leu Asn Ser Asn Ala Ile Lys Asn Leu Pro Pro Pro

```

- 12 -

35					40					45					
Leu	Gly	Gly	Ala	Ala	Gly	His	Pro	Gly	Ser	Ala	Val	Ser	Ala	Ala	Pro
50						55					60				
Gly	Ile	Leu	Tyr	Pro	Gly	Gly	Asn	Lys	Tyr	Gln	Thr	Ile	Asp	Asn	Tyr
65					70					75					80
Gln	Pro	Tyr	Pro	Cys	Ala	Glu	Asp	Glu	Glu	Cys	Gly	Thr	Asp	Glu	Tyr
				85					90					95	
Cys	Ala	Ser	Pro	Thr	Arg	Gly	Gly	Asp	Ala	Gly	Val	Gln	Ile	Cys	Leu
			100					105					110		
Ala	Cys	Arg	Lys	Arg	Arg	Lys	Arg	Cys	Met	Arg	His	Ala	Met	Cys	Cys
			115				120					125			
Pro	Gly	Asn	Tyr	Cys	Lys	Asn	Gly	Ile	Cys	Val	Ser	Ser	Asp	Gln	Asn
	130					135					140				
His	Phe	Arg	Gly	Glu	Ile	Glu	Glu	Thr	Ile	Thr	Glu	Ser	Phe	Gly	Asn
145					150					155					160
Asp	His	Ser	Thr	Leu	Asp	Gly	Tyr	Ser	Arg	Arg	Thr	Thr	Leu	Ser	Ser
				165					170					175	
Lys	Met	Tyr	His	Thr	Lys	Gly	Gln	Glu	Gly	Ser	Val	Cys	Leu	Arg	Ser
			180					185					190		
Ser	Asp	Cys	Ala	Ser	Gly	Leu	Cys	Cys	Ala	Arg	His	Phe	Trp	Ser	Lys
		195					200					205			
Ile	Cys	Lys	Pro	Val	Leu	Lys	Glu	Gly	Gln	Val	Cys	Thr	Lys	His	Arg
	210					215					220				
Arg	Lys	Gly	Ser	His	Gly	Leu	Glu	Ile	Phe	Gln	Arg	Cys	Tyr	Cys	Gly
225					230					235					240
Glu	Gly	Leu	Ser	Cys	Arg	Ile	Gln	Lys	Asp	His	His	Gln	Ala	Ser	Asn
				245					250					255	
Ser	Ser	Arg	Leu	His	Thr	Cys	Gln	Arg	His						
			260					265							

<210> 9

<211> 798

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(798)

<400> 9

atg	atg	gct	ctg	ggc	gca	gcg	gga	gct	acc	cgg	gtc	ttt	gtc	gcg	atg	48
Met	Met	Ala	Leu	Gly	Ala	Ala	Gly	Ala	Thr	Arg	Val	Phe	Val	Ala	Met	
1				5				10					15			

- 13 -

gta gcg gcg gct ctc ggc ggc cac cct ctg ctg gga gtg agc gcc acc	96
Val Ala Ala Ala Leu Gly Gly His Pro Leu Leu Gly Val Ser Ala Thr	
20 25 30	
ttg aac tcg gtt ctc aat tcc aac gct atc aag aac ctg ccc cca ccg	144
Leu Asn Ser Val Leu Asn Ser Asn Ala Ile Lys Asn Leu Pro Pro Pro	
35 40 45	
ctg ggc ggc gct gcg ggg cac cca ggc tct gca gtc agc gcc gcg ccg	192
Leu Gly Gly Ala Ala Gly His Pro Gly Ser Ala Val Ser Ala Ala Pro	
50 55 60	
gga atc ctg tac ccg ggc ggg aat aag tac cag acc att gac aac tac	240
Gly Ile Leu Tyr Pro Gly Gly Asn Lys Tyr Gln Thr Ile Asp Asn Tyr	
65 70 75 80	
cag ccg tac ccg tgc gca gag gac gag gag tgc ggc act gat gag tac	288
Gln Pro Tyr Pro Cys Ala Glu Asp Glu Glu Cys Gly Thr Asp Glu Tyr	
85 90 95	
tgc gct agt ccc acc cgc gga ggg gac gca ggc gtg caa atc tgt ctc	336
Cys Ala Ser Pro Thr Arg Gly Gly Asp Ala Gly Val Gln Ile Cys Leu	
100 105 110	
gcc tgc agg aag cgc cga aaa cgc tgc atg cgt cac gct atg tgc tgc	384
Ala Cys Arg Lys Arg Arg Lys Arg Cys Met Arg His Ala Met Cys Cys	
115 120 125	
ccc ggg aat tac tgc aaa aat gga ata tgt gtg tct tct gat caa aat	432
Pro Gly Asn Tyr Cys Lys Asn Gly Ile Cys Val Ser Ser Asp Gln Asn	
130 135 140	
cat ttc cga gga gaa att gag gaa acc atc act gaa agc ttt ggt aat	480
His Phe Arg Gly Glu Ile Glu Glu Thr Ile Thr Glu Ser Phe Gly Asn	
145 150 155 160	
gat cat agc acc ttg gat ggg tat tcc aga aga acc acc ttg tct tca	528
Asp His Ser Thr Leu Asp Gly Tyr Ser Arg Arg Thr Thr Leu Ser Ser	
165 170 175	
aaa atg tat cac acc aaa gga caa gaa ggt tct gtt tgt ctc cgg tca	576
Lys Met Tyr His Thr Lys Gly Gln Glu Gly Ser Val Cys Leu Arg Ser	
180 185 190	
tca gac tgt gcc tca gga ttg tgt tgt gct aga cac ttc tgg tcc aag	624
Ser Asp Cys Ala Ser Gly Leu Cys Cys Ala Arg His Phe Trp Ser Lys	
195 200 205	
atc tgt aaa cct gtc ctg aaa gaa ggt caa gtg tgt acc aag cat agg	672
Ile Cys Lys Pro Val Leu Lys Glu Gly Gln Val Cys Thr Lys His Arg	
210 215 220	
aga aaa ggc tct cat gga cta gaa ata ttc cag cgt tgt tac tgt gga	720
Arg Lys Gly Ser His Gly Leu Glu Ile Phe Gln Arg Cys Tyr Cys Gly	
225 230 235 240	
gaa ggt ctg tct tgc cgg ata cag aaa gat cac cat caa gcc agt aat	768
Glu Gly Leu Ser Cys Arg Ile Gln Lys Asp His His Gln Ala Ser Asn	
245 250 255	

- 14 -

tct tct agg ctt cac act tgt cag aga cac 798
 Ser Ser Arg Leu His Thr Cys Gln Arg His
 260 265

<210> 10
 <211> 702
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(537)

<400> 10
 gaa ttc ggc acg agg gtt ggg agg tat tgc cac agt ccc cac caa gga 48
 Glu Phe Gly Thr Arg Val Gly Arg Tyr Cys His Ser Pro His Gln Gly
 1 5 10 15
 tca tcg gcc tgc atg gtg tgt cgg aga aaa aag aag cgc tgc cac cga 96
 Ser Ser Ala Cys Met Val Cys Arg Arg Lys Lys Lys Arg Cys His Arg
 20 25 30
 gat ggc atg tgc tgc ccc agt acc cgc tgc aat aat ggc atc tgt atc 144
 Asp Gly Met Cys Cys Pro Ser Thr Arg Cys Asn Asn Gly Ile Cys Ile
 35 40 45
 cca gtt act gaa agc atc tta acc cct cac atc ccg gct ctg gat ggt 192
 Pro Val Thr Glu Ser Ile Leu Thr Pro His Ile Pro Ala Leu Asp Gly
 50 55 60
 act cgg cac aga gat cga aac cac ggt cat tac tca aac cat gac ttg 240
 Thr Arg His Arg Asp Arg Asn His Gly His Tyr Ser Asn His Asp Leu
 65 70 75 80
 gga tgg cag aat cta gga aga cca cac act aag atg tca cat ata aaa 288
 Gly Trp Gln Asn Leu Gly Arg Pro His Thr Lys Met Ser His Ile Lys
 85 90 95
 ggg cat gaa gga gac ccc tgc cta cga tca tca gac tgc att gaa ggg 336
 Gly His Glu Gly Asp Pro Cys Leu Arg Ser Ser Asp Cys Ile Glu Gly
 100 105 110
 ttt tgc tgt gct cgt cat ttc tgg acc aaa atc tgc aaa cca gtg ctc 384
 Phe Cys Cys Ala Arg His Phe Trp Thr Lys Ile Cys Lys Pro Val Leu
 115 120 125
 cat cag ggg gaa gtc tgt acc aaa caa cgc aag aag ggt tct cat ggg 432
 His Gln Gly Glu Val Cys Thr Lys Gln Arg Lys Lys Gly Ser His Gly
 130 135 140
 ctg gaa att ttc cag cgt tgc gac tgt gcg aag ggc ctg tct tgc aaa 480
 Leu Glu Ile Phe Gln Arg Cys Asp Cys Ala Lys Gly Leu Ser Cys Lys
 145 150 155 160
 gta tgg aaa gat gcc acc tac tcc tcc aaa gcc aga ctc cat gtg tgt 528
 Val Trp Lys Asp Ala Thr Tyr Ser Ser Lys Ala Arg Leu His Val Cys
 165 170 175

- 15 -

cag aaa att tgatcacat tgaggaacat catcaattgc agactgtgaa 577
 Gln Lys Ile
 gttgtgtatt taatgcatta tagcatggtg gaaaataagg ttcagatgca gaagaatggc 637
 taaaataaga aacgtgataa gaatatagat gatcacaaaa aaaaaaaaaa aaaagatgcg 697
 gccgc 702

<210> 11
 <211> 179
 <212> PRT
 <213> Homo sapiens

<400> 11
 Glu Phe Gly Thr Arg Val Gly Arg Tyr Cys His Ser Pro His Gln Gly
 1 5 10 15
 Ser Ser Ala Cys Met Val Cys Arg Arg Lys Lys Lys Arg Cys His Arg
 20 25 30
 Asp Gly Met Cys Cys Pro Ser Thr Arg Cys Asn Asn Gly Ile Cys Ile
 35 40 45
 Pro Val Thr Glu Ser Ile Leu Thr Pro His Ile Pro Ala Leu Asp Gly
 50 55 60
 Thr Arg His Arg Asp Arg Asn His Gly His Tyr Ser Asn His Asp Leu
 65 70 75 80
 Gly Trp Gln Asn Leu Gly Arg Pro His Thr Lys Met Ser His Ile Lys
 85 90 95
 Gly His Glu Gly Asp Pro Cys Leu Arg Ser Ser Asp Cys Ile Glu Gly
 100 105 110
 Phe Cys Cys Ala Arg His Phe Trp Thr Lys Ile Cys Lys Pro Val Leu
 115 120 125
 His Gln Gly Glu Val Cys Thr Lys Gln Arg Lys Lys Gly Ser His Gly
 130 135 140
 Leu Glu Ile Phe Gln Arg Cys Asp Cys Ala Lys Gly Leu Ser Cys Lys
 145 150 155 160
 Val Trp Lys Asp Ala Thr Tyr Ser Ser Lys Ala Arg Leu His Val Cys
 165 170 175
 Gln Lys Ile

<210> 12
 <211> 537
 <212> DNA
 <213> Homo sapiens

- 16 -

<220>

<221> CDS

<222> (1)..(537)

<400> 12

gaa ttc ggc acg agg gtt ggg agg tat tgc cac agt ccc cac caa gga	48
Glu Phe Gly Thr Arg Val Gly Arg Tyr Cys His Ser Pro His Gln Gly	
1 5 10 15	
tca tcg gcc tgc atg gtg tgt cgg aga aaa aag aag cgc tgc cac cga	96
Ser Ser Ala Cys Met Val Cys Arg Arg Lys Lys Lys Arg Cys His Arg	
20 25 30	
gat ggc atg tgc tgc ccc agt acc cgc tgc aat aat ggc atc tgt atc	144
Asp Gly Met Cys Cys Pro Ser Thr Arg Cys Asn Asn Gly Ile Cys Ile	
35 40 45	
cca gtt act gaa agc atc tta acc cct cac atc ccg gct ctg gat ggt	192
Pro Val Thr Glu Ser Ile Leu Thr Pro His Ile Pro Ala Leu Asp Gly	
50 55 60	
act cgg cac aga gat cga aac cac ggt cat tac tca aac cat gac ttg	240
Thr Arg His Arg Asp Arg Asn His Gly His Tyr Ser Asn His Asp Leu	
65 70 75 80	
gga tgg cag aat cta gga aga cca cac act aag atg tca cat ata aaa	288
Gly Trp Gln Asn Leu Gly Arg Pro His Thr Lys Met Ser His Ile Lys	
85 90 95	
ggg cat gaa gga gac ccc tgc cta cga tca tca gac tgc att gaa ggg	336
Gly His Glu Gly Asp Pro Cys Leu Arg Ser Ser Asp Cys Ile Glu Gly	
100 105 110	
ttt tgc tgt gct cgt cat ttc tgg acc aaa atc tgc aaa cca gtg ctc	384
Phe Cys Cys Ala Arg His Phe Trp Thr Lys Ile Cys Lys Pro Val Leu	
115 120 125	
cat cag ggg gaa gtc tgt acc aaa caa cgc aag aag ggt tct cat ggg	432
His Gln Gly Glu Val Cys Thr Lys Gln Arg Lys Lys Gly Ser His Gly	
130 135 140	
ctg gaa att ttc cag cgt tgc gac tgt gcg aag ggc ctg tct tgc aaa	480
Leu Glu Ile Phe Gln Arg Cys Asp Cys Ala Lys Gly Leu Ser Cys Lys	
145 150 155 160	
gta tgg aaa gat gcc acc tac tcc tcc aaa gcc aga ctc cat gtg tgt	528
Val Trp Lys Asp Ala Thr Tyr Ser Ser Lys Ala Arg Leu His Val Cys	
165 170 175	
cag aaa att	537
Gln Lys Ile	

<210> 13

<211> 928

<212> DNA

<213> Homo sapiens

<220>

- 17 -

<221> CDS

<222> (75)..(800)

<400> 13

```

ctcgaggcca aaattcggca cgaggccggg ctgtggtcta gcataaaggc ggagcccaga 60

agaaggggcg ggggt atg gga gaa gcc tcc cca cct gcc ccc gca agg cgg 110
      Met Gly Glu Ala Ser Pro Pro Ala Pro Ala Arg Arg
              1                      5                      10

cat ctg ctg gtc ctg ctg ctg ctc ctc tct acc ctg gtg atc ccc tcc 158
His Leu Leu Val Leu Leu Leu Leu Leu Ser Thr Leu Val Ile Pro Ser
      15                      20                      25

gct gca gct cct atc cat gat gct gac gcc caa gag agc tcc ttg ggt 206
Ala Ala Ala Pro Ile His Asp Ala Asp Ala Gln Glu Ser Ser Leu Gly
      30                      35                      40

ctc aca ggc ctc cag agc cta ctc caa ggc ttc agc cga ctt ttc ctg 254
Leu Thr Gly Leu Gln Ser Leu Leu Gln Gly Phe Ser Arg Leu Phe Leu
      45                      50                      55                      60

aaa ggt aac ctg ctt cgg ggc ata gac agc tta ttc tct gcc ccc atg 302
Lys Gly Asn Leu Leu Arg Gly Ile Asp Ser Leu Phe Ser Ala Pro Met
              65                      70                      75

gac ttc cgg ggc ctc cct ggg aac tac cac aaa gag gag aac cag gag 350
Asp Phe Arg Gly Leu Pro Gly Asn Tyr His Lys Glu Glu Asn Gln Glu
              80                      85                      90

cac cag ctg ggg aac aac acc ctc tcc agc cac ctc cag atc gac aag 398
His Gln Leu Gly Asn Asn Thr Leu Ser Ser His Leu Gln Ile Asp Lys
              95                      100                      105

atg acc gac aac aag aca gga gag gtg ctg atc tcc gag aat gtg gtg 446
Met Thr Asp Asn Lys Thr Gly Glu Val Leu Ile Ser Glu Asn Val Val
      110                      115                      120

gca tcc att caa cca gcg gag ggg agc ttc gag ggt gat ttg aag gta 494
Ala Ser Ile Gln Pro Ala Glu Gly Ser Phe Glu Gly Asp Leu Lys Val
      125                      130                      135                      140

ccc agg atg gag gag aag gag gcc ctg gta ccc atc cag aag gcc acg 542
Pro Arg Met Glu Glu Lys Glu Ala Leu Val Pro Ile Gln Lys Ala Thr
              145                      150                      155

gac agc ttc cac aca gaa ctc cat ccc cgg gtg gcc ttc tgg atc att 590
Asp Ser Phe His Thr Glu Leu His Pro Arg Val Ala Phe Trp Ile Ile
              160                      165                      170

aag ctg cca cgg cgg agg tcc cac cag gat gcc ctg gag ggc ggc cac 638
Lys Leu Pro Arg Arg Arg Ser His Gln Asp Ala Leu Glu Gly Gly His
              175                      180                      185

tgg ctc agc gag aag cga cac cgc ctg cag gcc atc cgg gat gga ctc 686
Trp Leu Ser Glu Lys Arg His Arg Leu Gln Ala Ile Arg Asp Gly Leu
      190                      195                      200

cgc aag ggg acc cac aag gac gtc cta gaa gag ggg acc gag agc tcc 734

```

- 18 -

Arg Lys Gly Thr His Lys Asp Val Leu Glu Glu Gly Thr Glu Ser Ser
 205 210 215 220
 tcc cac tcc agg ctg tcc ccc cga aag acc cac tta ctg tac atc ctc 782
 Ser His Ser Arg Leu Ser Pro Arg Lys Thr His Leu Leu Tyr Ile Leu
 225 230 235
 agg ccc tct cgg cag ctg taggggtggg gaccggggag cacctgcctg 830
 Arg Pro Ser Arg Gln Leu
 240
 tagcccccacat cagaccctgc cccaagcacc atatggaaat aaagttcttt cttacatcta 890
 aaaaaaaaaa aaaaaaaaaa aaaaaaattg gcggccgc 928

<210> 14
 <211> 242
 <212> PRT
 <213> Homo sapiens

<400> 14
 Met Gly Glu Ala Ser Pro Pro Ala Pro Ala Arg Arg His Leu Leu Val
 1 5 10 15
 Leu Leu Leu Leu Leu Ser Thr Leu Val Ile Pro Ser Ala Ala Ala Pro
 20 25 30
 Ile His Asp Ala Asp Ala Gln Glu Ser Ser Leu Gly Leu Thr Gly Leu
 35 40 45
 Gln Ser Leu Leu Gln Gly Phe Ser Arg Leu Phe Leu Lys Gly Asn Leu
 50 55 60
 Leu Arg Gly Ile Asp Ser Leu Phe Ser Ala Pro Met Asp Phe Arg Gly
 65 70 75 80
 Leu Pro Gly Asn Tyr His Lys Glu Glu Asn Gln Glu His Gln Leu Gly
 85 90 95
 Asn Asn Thr Leu Ser Ser His Leu Gln Ile Asp Lys Met Thr Asp Asn
 100 105 110
 Lys Thr Gly Glu Val Leu Ile Ser Glu Asn Val Val Ala Ser Ile Gln
 115 120 125
 Pro Ala Glu Gly Ser Phe Glu Gly Asp Leu Lys Val Pro Arg Met Glu
 130 135 140
 Glu Lys Glu Ala Leu Val Pro Ile Gln Lys Ala Thr Asp Ser Phe His
 145 150 155 160
 Thr Glu Leu His Pro Arg Val Ala Phe Trp Ile Ile Lys Leu Pro Arg
 165 170 175
 Arg Arg Ser His Gln Asp Ala Leu Glu Gly Gly His Trp Leu Ser Glu
 180 185 190
 Lys Arg His Arg Leu Gln Ala Ile Arg Asp Gly Leu Arg Lys Gly Thr

- 19 -

195	200	205	
His Lys Asp Val Leu Glu Glu Gly Thr Glu Ser Ser Ser His Ser Arg			
210	215	220	
Leu Ser Pro Arg Lys Thr His Leu Leu Tyr Ile Leu Arg Pro Ser Arg			
225	230	235	240
Gln Leu			
<210> 15			
<211> 726			
<212> DNA			
<213> Homo sapiens			
<220>			
<221> CDS			
<222> (1)..(726)			
<400> 15			
atg gga gaa gcc tcc cca cct gcc ccc gca agg cgg cat ctg ctg gtc	48		
Met Gly Glu Ala Ser Pro Pro Ala Pro Ala Arg Arg His Leu Leu Val			
1 5 10 15			
ctg ctg ctg ctc ctc tct acc ctg gtg atc ccc tcc gct gca gct cct	96		
Leu Leu Leu Leu Leu Ser Thr Leu Val Ile Pro Ser Ala Ala Ala Pro			
20 25 30			
atc cat gat gct gac gcc caa gag agc tcc ttg ggt ctc aca ggc ctc	144		
Ile His Asp Ala Asp Ala Gln Glu Ser Ser Leu Gly Leu Thr Gly Leu			
35 40 45			
cag agc cta ctc caa ggc ttc agc cga ctt ttc ctg aaa ggt aac ctg	192		
Gln Ser Leu Leu Gln Gly Phe Ser Arg Leu Phe Leu Lys Gly Asn Leu			
50 55 60			
ctt cgg ggc ata gac agc tta ttc tct gcc ccc atg gac ttc cgg ggc	240		
Leu Arg Gly Ile Asp Ser Leu Phe Ser Ala Pro Met Asp Phe Arg Gly			
65 70 75 80			
ctc cct ggg aac tac cac aaa gag gag aac cag gag cac cag ctg ggg	288		
Leu Pro Gly Asn Tyr His Lys Glu Glu Asn Gln Glu His Gln Leu Gly			
85 90 95			
aac aac acc ctc tcc agc cac ctc cag atc gac aag atg acc gac aac	336		
Asn Asn Thr Leu Ser Ser His Leu Gln Ile Asp Lys Met Thr Asp Asn			
100 105 110			
aag aca gga gag gtg ctg atc tcc gag aat gtg gtg gca tcc att caa	384		
Lys Thr Gly Glu Val Leu Ile Ser Glu Asn Val Val Ala Ser Ile Gln			
115 120 125			
cca gcg gag ggg agc ttc gag ggt gat ttg aag gta ccc agg atg gag	432		
Pro Ala Glu Gly Ser Phe Glu Gly Asp Leu Lys Val Pro Arg Met Glu			
130 135 140			
gag aag gag gcc ctg gta ccc atc cag aag gcc acg gac agc ttc cac	480		

- 20 -

Glu Lys Glu Ala Leu Val Pro Ile Gln Lys Ala Thr Asp Ser Phe His
 145 150 155 160
 aca gaa ctc cat ccc cgg gtg gcc ttc tgg atc att aag ctg cca cgg 528
 Thr Glu Leu His Pro Arg Val Ala Phe Trp Ile Ile Lys Leu Pro Arg
 165 170 175
 cgg agg tcc cac cag gat gcc ctg gag ggc ggc cac tgg ctc agc gag 576
 Arg Arg Ser His Gln Asp Ala Leu Glu Gly Gly His Trp Leu Ser Glu
 180 185 190
 aag cga cac cgc ctg cag gcc atc cgg gat gga ctc cgc aag ggg acc 624
 Lys Arg His Arg Leu Gln Ala Ile Arg Asp Gly Leu Arg Lys Gly Thr
 195 200 205
 cac aag gac gtc cta gaa gag ggg acc gag agc tcc tcc cac tcc agg 672
 His Lys Asp Val Leu Glu Glu Gly Thr Glu Ser Ser Ser His Ser Arg
 210 215 220
 ctg tcc ccc cga aag acc cac tta ctg tac atc ctc agg ccc tct cgg 720
 Leu Ser Pro Arg Lys Thr His Leu Leu Tyr Ile Leu Arg Pro Ser Arg
 225 230 235 240
 cag ctg 726
 Gln Leu

 <210> 16
 <211> 2380
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> CDS
 <222> (109)..(1155)

 <400> 26
 gtcgacccac gcgtccgctg tggcagccca gctaccggtc gtgaccagat ccagcttgca 60
 gctcagcttt gttcattcga attgggcggc ggccagcgcg gaacaaac atg cag cgg 117
 Met Gln Arg
 1
 ctc ggg ggt att ttg ctg tgt aca ctg ctg gcg gcg gcg gtc ccc act 165
 Leu Gly Gly Ile Leu Leu Cys Thr Leu Leu Ala Ala Ala Val Pro Thr
 5 10 15
 gct cct gct cct tcc ccg acg gtc act tgg act ccg gcg gag ccg ggc 213
 Ala Pro Ala Pro Ser Pro Thr Val Thr Trp Thr Pro Ala Glu Pro Gly
 20 25 30 35
 cca gct ctc aac tac cct cag gag gaa gct acg ctc aat gag atg ttt 261
 Pro Ala Leu Asn Tyr Pro Gln Glu Glu Ala Thr Leu Asn Glu Met Phe
 40 45 50
 cga gag gtg gag gag ctg atg gaa gac act cag cac aaa ctg cgc agt 309
 Arg Glu Val Glu Glu Leu Met Glu Asp Thr Gln His Lys Leu Arg Ser
 55 60 65

- 21 -

gcc gtg gag gag atg gag gcg gaa gaa gca gct gct aaa acg tcc tct	357
Ala Val Glu Glu Met Glu Ala Glu Glu Ala Ala Ala Lys Thr Ser Ser	
70 75 80	
gag gtg aac ctg gca agc tta cct ccc aac tat cac aat gag acc agc	405
Glu Val Asn Leu Ala Ser Leu Pro Pro Asn Tyr His Asn Glu Thr Ser	
85 90 95	
acg gag acc agg gtg gga aat aac aca gtc cat gtg cac cag gaa gtt	453
Thr Glu Thr Arg Val Gly Asn Asn Thr Val His Val His Gln Glu Val	
100 105 110 115	
cac aag ata acc aac aac cag agt gga cag gtg gtc ttt tct gag aca	501
His Lys Ile Thr Asn Asn Gln Ser Gly Gln Val Val Phe Ser Glu Thr	
120 125 130	
gtc att aca tct gta ggg gat gaa gaa ggc aag agg agc cat gaa tgt	549
Val Ile Thr Ser Val Gly Asp Glu Glu Gly Lys Arg Ser His Glu Cys	
135 140 145	
atc att gat gaa gac tgt ggg ccc acc agg tac tgc cag ttc tcc agc	597
Ile Ile Asp Glu Asp Cys Gly Pro Thr Arg Tyr Cys Gln Phe Ser Ser	
150 155 160	
ttc aag tac acc tgc cag cca tgc cgg gac cag cag atg cta tgc acc	645
Phe Lys Tyr Thr Cys Gln Pro Cys Arg Asp Gln Gln Met Leu Cys Thr	
165 170 175	
cga gac agt gag tgc tgt gga gac cag ctg tgt gcc tgg ggt cac tgc	693
Arg Asp Ser Glu Cys Cys Gly Asp Gln Leu Cys Ala Trp Gly His Cys	
180 185 190 195	
acc caa aag gcc acc aaa ggt ggc aat ggg acc atc tgt gac aac cag	741
Thr Gln Lys Ala Thr Lys Gly Gly Asn Gly Thr Ile Cys Asp Asn Gln	
200 205 210	
agg gat tgc cag cct ggc ctg tgt tgt gcc ttc caa aga ggc ctg ctg	789
Arg Asp Cys Gln Pro Gly Leu Cys Cys Ala Phe Gln Arg Gly Leu Leu	
215 220 225	
ttc ccc gtg tgc aca ccc ctg ccc gtg gag gga gag ctc tgc cat gac	837
Phe Pro Val Cys Thr Pro Leu Pro Val Glu Gly Glu Leu Cys His Asp	
230 235 240	
ccc acc agc cag ctg ctg gat ctc atc acc tgg gaa ctg gag cct gaa	885
Pro Thr Ser Gln Leu Leu Asp Leu Ile Thr Trp Glu Leu Glu Pro Glu	
245 250 255	
gga gct ttg gac cga tgc ccc tgc gcc agt ggc ctc cta tgc cag cca	933
Gly Ala Leu Asp Arg Cys Pro Cys Ala Ser Gly Leu Leu Cys Gln Pro	
260 265 270 275	
cac agc cac agt ctg gtg tac atg tgc aag cca gcc ttc gtg ggc agc	981
His Ser His Ser Leu Val Tyr Met Cys Lys Pro Ala Phe Val Gly Ser	
280 285 290	
cat gac cac agt gag gag agc cag ctg ccc agg gag gcc ccg gat gag	1029
His Asp His Ser Glu Glu Ser Gln Leu Pro Arg Glu Ala Pro Asp Glu	
295 300 305	

- 22 -

```

tac gaa gat gtt ggc ttc ata ggg gaa gtg cgc cag gag ctg gaa gac 1077
Tyr Glu Asp Val Gly Phe Ile Gly Glu Val Arg Gln Glu Leu Glu Asp
      310                      315                      320

ctg gag cgg agc cta gcc cag gag atg gca ttt gag ggg cct gcc cct 1125
Leu Glu Arg Ser Leu Ala Gln Glu Met Ala Phe Glu Gly Pro Ala Pro
      325                      330                      335

gtg gag tca cta ggc gga gag gag gag att taggcccaga cccagctgag 1175
Val Glu Ser Leu Gly Gly Glu Glu Glu Ile
      340                      345

tcactggtag atgtgcaata gaaatggcta atttattttc ccaggagtgt cccaagtgt 1235
ggaatggccg cagctccttc ccagtagctt ttctctctggc ttgacaaggt acagtgcagt 1295
acatttcttc cagccgcctt gcttctctga cttgggaaag acaggcatgg cgggtaaggg 1355
cagcggtgag tcgtccctcg ctgttgctag aaacgctgtc ttgttcttca tggatggaag 1415
atttgtttga agggagagga tgggaagggg tgaagtctgc tcatgatgga tttgggggat 1475
acagggagga ggatgcctgc cttgcagacg tggacttggc aaaatgtaac ctttgctttt 1535
gtcttgcgcc gctcccatgg gctgaggcag tggctacaca agagctatgc tgctctgtgg 1595
cctcccacat attcatccct gtgtttcagc tcctacctca ctgtcagcac agcccttcat 1655
agccacgccc cctcttgctc accacagcct aggaggggac cagaggggac ttctctcaga 1715
gcccacatgct ctctctctca accccatacc agcctctgtg ccagcgacag tccttccaaa 1775
tgaggggagt gaaatccttt ggtttaatta ttttctcctt caaggcacgc ctgccactaa 1835
ggtcaggctg acttgcattg ccctctaacg ttcgtagcag tgtggtggac actgtcttcc 1895
accgactgct tcaatacctc tgaaagccag tgctcggagt gcagttcgtg taaattaatt 1955
tgcaggaagt atacttggct aattgtaggg ctaggattgt gaatgaaatt tgcaaagtcg 2015
cttagcaaca atggaaagcc tttctcagtc acaccgagaa gtcacaacca agccaggttg 2075
tgtagagtac agctgtgaca tacagacaga agaaggctgg gctggatgtc aggctcaga 2135
tgacggtttc aggtgccagg aactattacc attctgtatc tatccagagt tattaaaatt 2195
gaaagttgca cacatttgta taagcatgcc tttctcctga gttttaaatt atatgtatac 2255
acaaacatgt ggccctcaaa gatcatgcac aaaccactac tctttgctaa ttcttggact 2315
tttctctttg attttcaata aatacaaatc cccttcatgc aaaaaaaaaa aaaaagggcg 2375
gccgc 2380

```

<210> 17
 <211> 349
 <212> PRT

- 23 -

<213> Homo sapiens

<400> 17

```

Met Gln Arg Leu Gly Gly Ile Leu Leu Cys Thr Leu Leu Ala Ala Ala
 1           5           10           15

Val Pro Thr Ala Pro Ala Pro Ser Pro Thr Val Thr Trp Thr Pro Ala
          20           25           30

Glu Pro Gly Pro Ala Leu Asn Tyr Pro Gln Glu Glu Ala Thr Leu Asn
          35           40           45

Glu Met Phe Arg Glu Val Glu Glu Leu Met Glu Asp Thr Gln His Lys
 50           55           60

Leu Arg Ser Ala Val Glu Glu Met Glu Ala Glu Glu Ala Ala Ala Lys
 65           70           75           80

Thr Ser Ser Glu Val Asn Leu Ala Ser Leu Pro Pro Asn Tyr His Asn
          85           90           95

Glu Thr Ser Thr Glu Thr Arg Val Gly Asn Asn Thr Val His Val His
          100          105          110

Gln Glu Val His Lys Ile Thr Asn Asn Gln Ser Gly Gln Val Val Phe
          115          120          125

Ser Glu Thr Val Ile Thr Ser Val Gly Asp Glu Glu Gly Lys Arg Ser
          130          135          140

His Glu Cys Ile Ile Asp Glu Asp Cys Gly Pro Thr Arg Tyr Cys Gln
          145          150          155          160

Phe Ser Ser Phe Lys Tyr Thr Cys Gln Pro Cys Arg Asp Gln Gln Met
          165          170          175

Leu Cys Thr Arg Asp Ser Glu Cys Cys Gly Asp Gln Leu Cys Ala Trp
          180          185          190

Gly His Cys Thr Gln Lys Ala Thr Lys Gly Gly Asn Gly Thr Ile Cys
          195          200          205

Asp Asn Gln Arg Asp Cys Gln Pro Gly Leu Cys Cys Ala Phe Gln Arg
          210          215          220

Gly Leu Leu Phe Pro Val Cys Thr Pro Leu Pro Val Glu Gly Glu Leu
          225          230          235          240

Cys His Asp Pro Thr Ser Gln Leu Leu Asp Leu Ile Thr Trp Glu Leu
          245          250          255

Glu Pro Glu Gly Ala Leu Asp Arg Cys Pro Cys Ala Ser Gly Leu Leu
          260          265          270

Cys Gln Pro His Ser His Ser Leu Val Tyr Met Cys Lys Pro Ala Phe
          275          280          285

Val Gly Ser His Asp His Ser Glu Glu Ser Gln Leu Pro Arg Glu Ala
          290          295          300

```

- 24 -

Pro Asp Glu Tyr Glu Asp Val Gly Phe Ile Gly Glu Val Arg Gln Glu
305 310 315 320

Leu Glu Asp Leu Glu Arg Ser Leu Ala Gln Glu Met Ala Phe Glu Gly
325 330 335

Pro Ala Pro Val Glu Ser Leu Gly Gly Glu Glu Glu Ile
340 345

<210> 18

<211> 1047

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(1047)

<400> 18

atg cag cgg ctc ggg ggt att ttg ctg tgt aca ctg ctg gcg gcg gcg 48
Met Gln Arg Leu Gly Gly Ile Leu Leu Cys Thr Leu Leu Ala Ala Ala
1 5 10 15

gtc ccc act gct cct gct cct tcc ccg acg gtc act tgg act ccg gcg 96
Val Pro Thr Ala Pro Ala Pro Ser Pro Thr Val Thr Trp Thr Pro Ala
20 25 30

gag ccg ggc cca gct ctc aac tac cct cag gag gaa gct acg ctc aat 144
Glu Pro Gly Pro Ala Leu Asn Tyr Pro Gln Glu Glu Ala Thr Leu Asn
35 40 45

gag atg ttt cga gag gtg gag gag ctg atg gaa gac act cag cac aaa 192
Glu Met Phe Arg Glu Val Glu Glu Leu Met Glu Asp Thr Gln His Lys
50 55 60

ctg cgc agt gcc gtg gag gag atg gag gcg gaa gaa gca gct gct aaa 240
Leu Arg Ser Ala Val Glu Glu Met Glu Ala Glu Glu Ala Ala Lys
65 70 75 80

acg tcc tct gag gtg aac ctg gca agc tta cct ccc aac tat cac aat 288
Thr Ser Ser Glu Val Asn Leu Ala Ser Leu Pro Pro Asn Tyr His Asn
85 90 95

gag acc agc acg gag acc agg gtg gga aat aac aca gtc cat gtg cac 336
Glu Thr Ser Thr Glu Thr Arg Val Gly Asn Asn Thr Val His Val His
100 105 110

cag gaa gtt cac aag ata acc aac aac cag agt gga cag gtg gtc ttt 384
Gln Glu Val His Lys Ile Thr Asn Asn Gln Ser Gly Gln Val Val Phe
115 120 125

tct gag aca gtc att aca tct gta ggg gat gaa gaa ggc aag agg agc 432
Ser Glu Thr Val Ile Thr Ser Val Gly Asp Glu Glu Gly Lys Arg Ser
130 135 140

cat gaa tgt atc att gat gaa gac tgt ggg ccc acc agg tac tgc cag 480
His Glu Cys Ile Ile Asp Glu Asp Cys Gly Pro Thr Arg Tyr Cys Gln

- 25 -

145	150	155	160	
ttc tcc agc ttc aag tac acc tgc cag cca tgc cgg gac cag cag atg				528
Phe Ser Ser Phe Lys Tyr Thr Cys Gln Pro Cys Arg Asp Gln Gln Met				
	165	170	175	
cta tgc acc cga gac agt gag tgc tgt gga gac cag ctg tgt gcc tgg				576
Leu Cys Thr Arg Asp Ser Glu Cys Cys Gly Asp Gln Leu Cys Ala Trp				
	180	185	190	
ggc cac tgc acc caa aag gcc acc aaa ggt ggc aat ggg acc atc tgt				624
Gly His Cys Thr Gln Lys Ala Thr Lys Gly Gly Asn Gly Thr Ile Cys				
	195	200	205	
gac aac cag agg gat tgc cag cct ggc ctg tgt tgt gcc ttc caa aga				672
Asp Asn Gln Arg Asp Cys Gln Pro Gly Leu Cys Cys Ala Phe Gln Arg				
	210	215	220	
ggc ctg ctg ttc ccc gtg tgc aca ccc ctg ccc gtg gag gga gag ctc				720
Gly Leu Leu Phe Pro Val Cys Thr Pro Leu Pro Val Glu Gly Glu Leu				
	225	230	235	240
tgc cat gac ccc acc agc cag ctg ctg gat ctc atc acc tgg gaa ctg				768
Cys His Asp Pro Thr Ser Gln Leu Leu Asp Leu Ile Thr Trp Glu Leu				
	245	250	255	
gag cct gaa gga gct ttg gac cga tgc ccc tgc gcc agt ggc ctc cta				816
Glu Pro Glu Gly Ala Leu Asp Arg Cys Pro Cys Ala Ser Gly Leu Leu				
	260	265	270	
tgc cag cca cac agc cac agt ctg gtg tac atg tgc aag cca gcc ttc				864
Cys Gln Pro His Ser His Ser Leu Val Tyr Met Cys Lys Pro Ala Phe				
	275	280	285	
gtg ggc agc cat gac cac agt gag gag agc cag ctg ccc agg gag gcc				912
Val Gly Ser His Asp His Ser Glu Glu Ser Gln Leu Pro Arg Glu Ala				
	290	295	300	
ccg gat gag tac gaa gat gtt ggc ttc ata ggg gaa gtg cgc cag gag				960
Pro Asp Glu Tyr Glu Asp Val Gly Phe Ile Gly Glu Val Arg Gln Glu				
	305	310	315	320
ctg gaa gac ctg gag cgg agc cta gcc cag gag atg gca ttt gag ggg				1008
Leu Glu Asp Leu Glu Arg Ser Leu Ala Gln Glu Met Ala Phe Glu Gly				
	325	330	335	
cct gcc cct gtg gag tca cta ggc gga gag gag gag att				1047
Pro Ala Pro Val Glu Ser Leu Gly Gly Glu Glu Glu Ile				
	340	345		

<210> 19

<211> 8

<212> PRT

<213> synthtic construct

<400> 19

Asp Tyr Lys Asp Asp Asp Asp Lys

1

5

```
<220>  
<221> CDS  
<222> (712)..(1500)
```

<400>	20																																		
gtc	gac	cc	ca	c	gcg	tcc	gg	cg	ccc	gcg	gcg	agc	gtag	cg	ca	agt	tcc	g	ctc	ct	agg	c	60												
atc	gct	gc	gc		tgg	cag	cg	at	t	cgt	gt	tct	t	ttg	t	gag	tca	g	ggg	gac	a	acg	ctt	cgg	ggg	ca	120								
act	gt	gag	t	g	cg	gt	gt	ggg	g	gac	ct	cg	at	t	tct	ctt	caga	t	tct	cg	agg	at	t	cg	gt	tcc	ggg	180							
gac	gt	tct	ct	c	gat	ccc	ct	ac	t	aaa	agc	gc	ct	g	cta	act	ttg	a	aaa	agg	ag	ca	ct	gt	gt	cct	g	240							
caa	agt	ttt	ga		cac	ata	aa	agg	a	atag	g	aaa	ag	a	agagg	ag	aga	a	aa	g	ca	act	g	agt	tga	agg	a	300							
ga	agg	ag	ct	g	at	gc	ggg	gc	ct	gat	ca	at		ta	ag	agg	aga	g	gt	taa	acc	gc	cg	ag	at	ccc	g	360							
gc	ggg	ac	caa		gg	agg	t	gc	g	gg	ca	aga	ag	a	ac	g	ga	ag	cg	g	tg	cg	at	cca	ca	ggg	ct	ggg	420						
ttt	t	ctt	g	ca	c	ctt	g	g	t	ca	cg	ct	c	ctt	g	cg	aga	aa	gc	g	c	ct	cg	cat	t	g	att	g	ctt	480					
cag	tt	att	t	gc	a	ga	act	t	ct	gt	g	gt	g	aga	ag	cg	gg	t	ct	cg	ctt	ggg	t	t	cc	g	cta	at	540						
tt	ct	gt	c	ct	g	t	agg	cg	t	g	aga	ct	g	ag	gt	t	cat	a	agg	gt	c	ct	g	g	gt	cccc	ga	ac	ca	gg	a	agg	gt	600	
tg	agg	ga	a	ca	ca	at	ct	g	caa	g	cccc	cg	cg	a	cca	ag	t	g	g	g	cccc	ct	g	tt	ggg	gt	c	ct	660						
cc	ct	c	c	ct	t	t		gc	att	ccc	ac	c	ct	c	gg	g	g	g	c	ttt	g	cg	t	ctt	c	ct	g	ggg	g	acc	c	ct	c	gc	717
																								Pro Arg	1										
cg	g	g	at	g	gc	g	t	at	g	cg	ag	a	g	at	t	cg	t	cc	g	g	t	g	g	ct					765						
Arg	Glu	Met	Ala	Ala	Leu	Met	Arg	Ser	Lys	Asp	Ser	Ser	Cys	Cys	Leu																				
																								5	10	15									
ct	c	t	a	ct	g	gc	g	t	g	ct	at	g	gt	g	g	ag	ag	t	c	a	c	ag	at	g	g	c	ag	t			813				
Leu	Leu	Leu	Ala	Ala	Val	Leu	Met	Val	Glu	Ser	Ser	Gln	Ile	Gly	Ser																				
																								20	25	30									
t	g	cg	g	gc	a	a	ct	a	c	t	cc	at	a	g	t	cc	t	ct	g	g	g	g	g	g	g	g	g	g	g	g	g	861			
Ser	Arg	Ala	Lys	Leu	Asn	Ser	Ile	Lys	Ser	Ser	Leu	Gly	Gly	Glu	Thr																				
																								35	40	45	50								
c	c	t	g	g	t	c	ag	g	c	c	a	a	t	c	g	a	t	c	t	g	g	c	at	t	a	c	a	a	g	g	a	g	909		
Pro	Gly	Gln	Ala	Ala	Asn	Arg	Ser	Ala	Gly	Met	Tyr	Gln	Gly	Leu	Ala																				
																								55	60	65									
t	t	c	g	g	g	g	g</																												


```

gttagagttt aacaaaatac tcctagaata acttggtata caataggttc taaaaataaa 1960
attgctaaac aagaaatgaa aacatggagc attgttaatt tacaacagaa aattaccttt 2020
tgatttgtaa cactacttct gctgttcaat caagagtctt ggtagataag aaaaaaatca 2080
gtcaatatat ccaaataatt gcaaaataat ggccagttgt ttaggaaggc ctttaggaag 2140
acaaataaat aacaaacaaa cagccacaaa tacttttttt tcaaaatatt agttttacct 2200
gtaattaata agaactgata caagacaaaa acagttcctt cagattctac ggaatgacag 2260
tatatctctc tttatcctat gtgattcctg ctctgaatgc attatatatt ccaaagtata 2320
cccataaatt gtgactagta aaatacttac acagagcaga attttcacag atggcaaaaa 2380
aatttaaaga tgtccaatat atgtgggaaa agagctaaca gagagatcat tatttcttaa 2440
agattggcca taacctgtat tttgatagaa ttagattggg aaatacatgt attcatacat 2500
actctgtggg aatagagact tgagctggat ctgtactgca ctggagtaag caagaaaatt 2560
gggaaaactt tttcgtttgt tcaggttttg gcaacacata gatcatatgt ctgaggcaca 2620
agttggctgt tcacttttga aaccagggga tgcacagtct aaatgaatat ctgcatggga 2680
tttgtatcat aatatttact atgcagatga attcagtggt aggtcctgtg tccgtactat 2740
cctcaaatta tttattttat agtgctgaga tcctcaaata atctcaattt caggagggtt 2800
cacaaaatgg actcctgaag tagacagagt agtgagggtt cattgccctc tataagcttc 2860
tgactagcca atggcatcat ccaattttct tcccaaacct ctgcagcatc tgctttattg 2920
ccaaagggtc agtttcggtt ttctgcagcc attgcggtta aaaaatataa gtaggataac 2980
ttgtaaaacc tgcatattgc taatctatag acaccacagt ttctaaattc tttgaaacca 3040
ctttactact ttttttaaac ttaactcagt tctaaatact ttgtctggag cacaaaacaa 3100
taaaaggtta tcttatagtt gtgactttaa actttttag accacaattc actttttagt 3160
tttcttttac ttaaatecca tctgcagtct caaatttaag ttctcccagt agagattgag 3220
tttgagcctg tatatctatt aaaaatttca acttcccaca tatatttact aagatgatta 3280
agacttacat tttctgcaca ggtctgcaa aacaaaaatt ataaactagt ccatccaaga 3340
accaaagttt gtataaacag gttgctataa gcttggtgaa atgaaaatgg aacatttcaa 3400
tcaaacattt cctatataac aattattata ttacaattt gggttctgca atatttttct 3460
tatgtccacc cttttaaaaa ttattatttg aagtaattta ttacaggaa atgttaatga 3520
gatgtatttt cttatagaga tatttcttac agaaagcttt gtagcagaat atatttgcag 3580
ctattgactt tgtaatttag gaaaaatgta taataagata aaatctatta aatttttctc 3640
ctctaaaaac tgaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaagggc ggccgc 3696

```

- 29 -

<210> 21

<211> 263

<212> PRT

<213> Homo sapiens

<400> 21

Pro	Arg	Arg	Glu	Met	Ala	Ala	Leu	Met	Arg	Ser	Lys	Asp	Ser	Ser	Cys
1				5					10					15	

Cys	Leu	Leu	Leu	Leu	Ala	Ala	Val	Leu	Met	Val	Glu	Ser	Ser	Gln	Ile
			20					25					30		

Gly	Ser	Ser	Arg	Ala	Lys	Leu	Asn	Ser	Ile	Lys	Ser	Ser	Leu	Gly	Gly
		35					40					45			

Glu	Thr	Pro	Gly	Gln	Ala	Ala	Asn	Arg	Ser	Ala	Gly	Met	Tyr	Gln	Gly
	50					55					60				

Leu	Ala	Phe	Gly	Gly	Ser	Lys	Lys	Gly	Lys	Asn	Leu	Gly	Gln	Ala	Tyr
65					70					75					80

Pro	Cys	Ser	Ser	Asp	Lys	Glu	Cys	Glu	Val	Gly	Arg	Tyr	Cys	His	Ser
				85					90					95	

Pro	His	Gln	Gly	Ser	Ser	Ala	Cys	Met	Val	Cys	Arg	Arg	Lys	Lys	Lys
			100					105					110		

Arg	Cys	His	Arg	Asp	Gly	Met	Cys	Cys	Pro	Ser	Thr	Arg	Cys	Asn	Asn
		115					120					125			

Gly	Ile	Cys	Ile	Pro	Val	Thr	Glu	Ser	Ile	Leu	Thr	Pro	His	Ile	Pro
	130					135					140				

Ala	Leu	Asp	Gly	Thr	Arg	His	Arg	Asp	Arg	Asn	His	Gly	His	Tyr	Ser
145					150					155					160

Asn	His	Asp	Leu	Gly	Trp	Gln	Asn	Leu	Gly	Arg	Pro	His	Thr	Lys	Met
			165						170					175	

Ser	His	Ile	Lys	Gly	His	Glu	Gly	Asp	Pro	Cys	Leu	Arg	Ser	Ser	Asp
			180					185					190		

Cys	Ile	Glu	Gly	Phe	Cys	Cys	Ala	Arg	His	Phe	Trp	Thr	Lys	Ile	Cys
		195					200					205			

Lys	Pro	Val	Leu	His	Gln	Gly	Glu	Val	Cys	Thr	Lys	Gln	Arg	Lys	Lys
	210					215					220				

Gly	Ser	His	Gly	Leu	Glu	Ile	Phe	Gln	Arg	Cys	Asp	Cys	Ala	Lys	Gly
225					230					235					240

Leu	Ser	Cys	Lys	Val	Trp	Lys	Asp	Ala	Thr	Tyr	Ser	Ser	Lys	Ala	Arg
				245					250					255	

Leu	His	Val	Cys	Gln	Lys	Ile
				260		

- 30 -

<210> 22
 <211> 789
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(789)

<400> 22
 cct cgc cgg gag atg gcc gcg ttg atg cgg agc aag gat tcg tcc tgc 48
 Pro Arg Arg Glu Met Ala Ala Leu Met Arg Ser Lys Asp Ser Ser Cys
 1 5 10 15
 tgc ctg ctc cta ctg gcc gcg gtg ctg atg gtg gag agc tca cag atc 96
 Cys Leu Leu Leu Leu Ala Ala Val Leu Met Val Glu Ser Ser Gln Ile
 20 25 30
 ggc agt tcg cgg gcc aaa ctc aac tcc atc aag tcc tct ctg ggc ggc 144
 Gly Ser Ser Arg Ala Lys Leu Asn Ser Ile Lys Ser Ser Leu Gly Gly
 35 40 45
 gag acg cct ggt cag gcc gcc aat cga tct gcg ggc atg tac caa gga 192
 Glu Thr Pro Gly Gln Ala Ala Asn Arg Ser Ala Gly Met Tyr Gln Gly
 50 55 60
 ctg gca ttc ggc ggc agt aag aag ggc aaa aac ctg ggg cag gcc tac 240
 Leu Ala Phe Gly Gly Ser Lys Lys Gly Lys Asn Leu Gly Gln Ala Tyr
 65 70 75 80
 cct tgt agc agt gat aag gag tgt gaa gtt ggg agg tat tgc cac agt 288
 Pro Cys Ser Ser Asp Lys Glu Cys Glu Val Gly Arg Tyr Cys His Ser
 85 90 95
 ccc cac caa gga tca tcg gcc tgc atg gtg tgt cgg aga aaa aag aag 336
 Pro His Gln Gly Ser Ser Ala Cys Met Val Cys Arg Arg Lys Lys Lys
 100 105 110
 cgc tgc cac cga gat ggc atg tgc tgc ccc agt acc cgc tgc aat aat 384
 Arg Cys His Arg Asp Gly Met Cys Cys Pro Ser Thr Arg Cys Asn Asn
 115 120 125
 ggc atc tgt atc cca gtt act gaa agc atc tta acc cct cac atc ccg 432
 Gly Ile Cys Ile Pro Val Thr Glu Ser Ile Leu Thr Pro His Ile Pro
 130 135 140
 gct ctg gat ggt act cgg cac aga gat cga aac cac ggt cat tac tca 480
 Ala Leu Asp Gly Thr Arg His Arg Asp Arg Asn His Gly His Tyr Ser
 145 150 155 160
 aac cat gac ttg gga tgg cag aat cta gga aga cca cac act aag atg 528
 Asn His Asp Leu Gly Trp Gln Asn Leu Gly Arg Pro His Thr Lys Met
 165 170 175
 tca cat ata aaa ggc cat gaa gga gac ccc tgc cta cga tca tca gac 576
 Ser His Ile Lys Gly His Glu Gly Asp Pro Cys Leu Arg Ser Ser Asp
 180 185 190

- 31 -

tgc att gaa ggg ttt tgc tgt gct cgt cat ttc tgg acc aaa atc tgc 624
 Cys Ile Glu Gly Phe Cys Cys Ala Arg His Phe Trp Thr Lys Ile Cys
 195 200 205

aaa cca gtg ctc cat cag ggg gaa gtc tgt acc aaa caa cgc aag aag 672
 Lys Pro Val Leu His Gln Gly Glu Val Cys Thr Lys Gln Arg Lys Lys
 210 215 220

ggc tct cat ggg ctg gaa att ttc cag cgt tgc gac tgt gcg aag ggc 720
 Gly Ser His Gly Leu Glu Ile Phe Gln Arg Cys Asp Cys Ala Lys Gly
 225 230 235 240

ctg tct tgc aaa gta tgg aaa gat gcc acc tac tcc tcc aaa gcc aga 768
 Leu Ser Cys Lys Val Trp Lys Asp Ala Thr Tyr Ser Ser Lys Ala Arg
 245 250 255

ctc cat gtg tgt cag aaa att 789
 Leu His Val Cys Gln Lys Ile
 260

<210> 23

<211> 54

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: consensus
sequence

<220>

<223> Xaa's at positions
2,3,5,6,8-12,14-26,28,29,31-36,38-42,45-48, and
50-53 may be any amino acid

<220>

<223> Xaa's at postions 22-26 may be absent

<400> 23

Cys Xaa Xaa Asp Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa Xaa
 1 5 10 15

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa Cys Xaa Xaa
 20 25 30

Xaa Xaa Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa Cys Cys Xaa Xaa Xaa Xaa
 35 40 45

Cys Xaa Xaa Xaa Xaa Cys
 50

<210> 24

<211> 123

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: consensus

- 32 -

sequence

<220>

<223> Xaa's at positions

2,5,8,9,11,14-23,25,27-30,32,33,35-53,60,62,63,65,
68,70,71,73-96,98,100,101,104, and 106-122 may be
any amino acid

<220>

<223> Xaa's at positions 22,23,51-58,89-96 and 116-122
may be absent

<400> 24

Cys Xaa Xaa Xaa Xaa Asp Cys Xaa Xaa Gly Xaa Cys Cys Xaa Xaa Xaa
1 5 10 15

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Pro Xaa Xaa Xaa Xaa Gly Xaa
20 25 30

Xaa Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
35 40 45

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Cys Xaa Xaa Pro
50 55 60

Xaa Xaa Xaa Xaa Gly Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
65 70 75 80

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
85 90 95

Cys Xaa Cys Xaa Xaa Gly Leu Xaa Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa
100 105 110

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys
115 120

<210> 25

<211> 87

<212> PRT

<213> Homo sapiens

<400> 25

Ile Asn Leu Glu Asn Gly Glu Leu Cys Met Asn Ser Ala Gln Cys Lys
1 5 10 15

Ser Asn Cys Cys Gln His Ser Ser Ala Leu Gly Leu Ala Arg Cys Thr
20 25 30

Ser Met Ala Ser Glu Asn Ser Glu Cys Ser Val Lys Thr Leu Tyr Gly
35 40 45

Ile Tyr Tyr Lys Cys Pro Cys Glu Arg Gly Leu Thr Cys Glu Gly Asp
50 55 60

Lys Thr Ile Val Gly Ser Ile Thr Asn Thr Asn Phe Gly Ile Cys His
65 70 75 80

- 33 -

Asp Ala Gly Arg Ser Lys Gln
85

<210> 26

<211> 835

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (57)..(746)

<400> 25

gaattcggca cgaggcagaa ggcgcgaatg aaggcaaagc ctcccaccca cctgca atg 59
Met
1

tgt cga ctg agg gtc ttg ctg ctg ctg ctc ccc ttg gcc ttc gtg tcc 107
Cys Arg Leu Arg Val Leu Leu Leu Leu Leu Pro Leu Ala Phe Val Ser
5 10 15

tcc tct gct ctc ccc atc cat gat gtc gac tct cag cag aac acc tcc 155
Ser Ser Ala Leu Pro Ile His Asp Val Asp Ser Gln Gln Asn Thr Ser
20 25 30

ggg ttc ctg ggc ctt cag agg ctt ctc caa agc ttt agt cga ctg ttc 203
Gly Phe Leu Gly Leu Gln Arg Leu Leu Gln Ser Phe Ser Arg Leu Phe
35 40 45

cta aaa aat gac ctg cta cga gac ctg gac aac ttc ttc tcc tcc ccc 251
Leu Lys Asn Asp Leu Leu Arg Asp Leu Asp Asn Phe Phe Ser Ser Pro
50 55 60 65

atg gac ttc cga gac ctt cct agg aac ttc cat cag gaa gag aac cag 299
Met Asp Phe Arg Asp Leu Pro Arg Asn Phe His Gln Glu Glu Asn Gln
70 75 80

gag cac aga atg ggc aac cat acc ctc tcc agc cac cta cag ata gac 347
Glu His Arg Met Gly Asn His Thr Leu Ser Ser His Leu Gln Ile Asp
85 90 95

aag gtg act gac aac cag aca ggg gag gtg cac atc tcg gag aaa gtc 395
Lys Val Thr Asp Asn Gln Thr Gly Glu Val His Ile Ser Glu Lys Val
100 105 110

gag gcc tcc att gag cca gaa cgg aac ccg gaa ggg gac tgg aag gtt 443
Glu Ala Ser Ile Glu Pro Glu Arg Asn Pro Glu Gly Asp Trp Lys Val
115 120 125

ccc aaa gta gaa gca aaa gag ccc ccg gtg cct gtg cag aag gtc acc 491
Pro Lys Val Glu Ala Lys Glu Pro Pro Val Pro Val Gln Lys Val Thr
130 135 140 145

gac agc ttg cac cca gag ccc cgg cag gtg gct ttc tgg atc atg aag 539
Asp Ser Leu His Pro Glu Pro Arg Gln Val Ala Phe Trp Ile Met Lys
150 155 160

atg cca agg cgg agg acc cag ccc gat gtc cag gat gga ggc cgc tgg 587

- 34 -

Met Pro Arg Arg Arg Thr Gln Pro Asp Val Gln Asp Gly Gly Arg Trp
165 170 175

ctc ata gaa aag cga cat cgc atg cag gcc atc cgg gat ggg ctc cgt 635
Leu Ile Glu Lys Arg His Arg Met Gln Ala Ile Arg Asp Gly Leu Arg
180 185 190

gga ggc gcc cgt gag gac agc ctg gag gat ggg gtc cat atc ccc caa 683
Gly Gly Ala Arg Glu Asp Ser Leu Glu Asp Gly Val His Ile Pro Gln
195 200 205

cac gcc aag ctg cct gtc aga aag aca cac ttt ctc tac atc ctc agg 731
His Ala Lys Leu Pro Val Arg Lys Thr His Phe Leu Tyr Ile Leu Arg
210 215 220 225

cca tcc caa cag ctg taagtgggga ccagatgtcc cacaccctac cccaacacca 786
Pro Ser Gln Gln Leu
230

tatggaaata aaggttttct tacatctaaa aaaaaaaaaa aaaaaaaaaa 835

<210> 27
<211> 230
<212> PRT
<213> Mus musculus

<400> 27
Met Cys Arg Leu Arg Val Leu Leu Leu Leu Leu Pro Leu Ala Phe Val
1 5 10 15

Ser Ser Ser Ala Leu Pro Ile His Asp Val Asp Ser Gln Gln Asn Thr
20 25 30

Ser Gly Phe Leu Gly Leu Gln Arg Leu Leu Gln Ser Phe Ser Arg Leu
35 40 45

Phe Leu Lys Asn Asp Leu Leu Arg Asp Leu Asp Asn Phe Phe Ser Ser
50 55 60

Pro Met Asp Phe Arg Asp Leu Pro Arg Asn Phe His Gln Glu Glu Asn
65 70 75 80

Gln Glu His Arg Met Gly Asn His Thr Leu Ser Ser His Leu Gln Ile
85 90 95

Asp Lys Val Thr Asp Asn Gln Thr Gly Glu Val His Ile Ser Glu Lys
100 105 110

Val Glu Ala Ser Ile Glu Pro Glu Arg Asn Pro Glu Gly Asp Trp Lys
115 120 125

Val Pro Lys Val Glu Ala Lys Glu Pro Pro Val Pro Val Gln Lys Val
130 135 140

Thr Asp Ser Leu His Pro Glu Pro Arg Gln Val Ala Phe Trp Ile Met
145 150 155 160

Lys Met Pro Arg Arg Arg Thr Gln Pro Asp Val Gln Asp Gly Gly Arg

- 35 -

	165		170		175										
Trp	Leu	Ile	Glu	Lys	Arg	His	Arg	Met	Gln	Ala	Ile	Arg	Asp	Gly	Leu
	180							185					190		
Arg	Gly	Gly	Ala	Arg	Glu	Asp	Ser	Leu	Glu	Asp	Gly	Val	His	Ile	Pro
	195						200					205			
Gln	His	Ala	Lys	Leu	Pro	Val	Arg	Lys	Thr	His	Phe	Leu	Tyr	Ile	Leu
	210					215					220				
Arg	Pro	Ser	Gln	Gln	Leu										
225					230										

<210> 28
 <211> 690
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (1)..(690)

<400> 28

atg	tgt	cga	ctg	agg	gtc	ttg	ctg	ctg	ctg	ctc	ccc	ttg	gcc	ttc	gtg	48
Met	Cys	Arg	Leu	Arg	Val	Leu	Leu	Leu	Leu	Leu	Pro	Leu	Ala	Phe	Val	
1				5					10					15		
tcc	tcc	tct	gct	ctc	ccc	atc	cat	gat	gtc	gac	tct	cag	cag	aac	acc	96
Ser	Ser	Ser	Ala	Leu	Pro	Ile	His	Asp	Val	Asp	Ser	Gln	Gln	Asn	Thr	
			20					25					30			
tcc	ggg	ttc	ctg	ggc	ctt	cag	agg	ctt	ctc	caa	agc	ttt	agt	cga	ctg	144
Ser	Gly	Phe	Leu	Gly	Leu	Gln	Arg	Leu	Leu	Gln	Ser	Phe	Ser	Arg	Leu	
		35					40					45				
ttc	cta	aaa	aat	gac	ctg	cta	cga	gac	ctg	gac	aac	ttc	ttc	tcc	tcc	192
Phe	Leu	Lys	Asn	Asp	Leu	Leu	Arg	Asp	Leu	Asp	Asn	Phe	Phe	Ser	Ser	
	50					55					60					
ccc	atg	gac	ttc	cga	gac	ctt	cct	agg	aac	ttc	cat	cag	gaa	gag	aac	240
Pro	Met	Asp	Phe	Arg	Asp	Leu	Pro	Arg	Asn	Phe	His	Gln	Glu	Glu	Asn	
	65				70				75						80	
cag	gag	cac	aga	atg	ggc	aac	cat	acc	ctc	tcc	agc	cac	cta	cag	ata	288
Gln	Glu	His	Arg	Met	Gly	Asn	His	Thr	Leu	Ser	Ser	His	Leu	Gln	Ile	
				85				90					95			
gac	aag	gtg	act	gac	aac	cag	aca	ggg	gag	gtg	cac	atc	tcg	gag	aaa	336
Asp	Lys	Val	Thr	Asp	Asn	Gln	Thr	Gly	Glu	Val	His	Ile	Ser	Glu	Lys	
			100					105					110			
gtc	gag	gcc	tcc	att	gag	cca	gaa	cgg	aac	ccg	gaa	ggg	gac	tgg	aag	384
Val	Glu	Ala	Ser	Ile	Glu	Pro	Glu	Arg	Asn	Pro	Glu	Gly	Asp	Trp	Lys	
		115					120					125				
gtt	ccc	aaa	gta	gaa	gca	aaa	gag	ccc	ccg	gtg	cct	gtg	cag	aag	gtc	432
Val	Pro	Lys	Val	Glu	Ala	Lys	Glu	Pro	Pro	Val	Pro	Val	Gln	Lys	Val	

- 36 -

130	135	140	
acc gac agc ttg cac cca gag ccc cgg cag gtg gct ttc tgg atc atg			480
Thr Asp Ser Leu His Pro Glu Pro Arg Gln Val Ala Phe Trp Ile Met			
145	150	155	160
aag atg cca agg cgg agg acc cag ccc gat gtc cag gat gga ggc cgc			528
Lys Met Pro Arg Arg Arg Thr Gln Pro Asp Val Gln Asp Gly Gly Arg			
	165	170	175
tgg ctc ata gaa aag cga cat cgc atg cag gcc atc cgg gat ggg ctc			576
Trp Leu Ile Glu Lys Arg His Arg Met Gln Ala Ile Arg Asp Gly Leu			
	180	185	190
cgt gga ggc gcc cgt gag gac agc ctg gag gat ggg gtc cat atc ccc			624
Arg Gly Gly Ala Arg Glu Asp Ser Leu Glu Asp Gly Val His Ile Pro			
	195	200	205
caa cac gcc aag ctg cct gtc aga aag aca cac ttt ctc tac atc ctc			672
Gln His Ala Lys Leu Pro Val Arg Lys Thr His Phe Leu Tyr Ile Leu			
	210	215	220
agg cca tcc caa cag ctg			690
Arg Pro Ser Gln Gln Leu			
225	230		

<210> 29

<211> 51

<212> PRT

<213> Artificial Sequence

<220>

<223> Xaa at positions

3-5, 7, 9-15, 18, 20-22, 24-27, 29, 31, 33, 34, 36-39, 42, 44,
45, and 47-50 may be any amino acid

<220>

<223> Description of Artificial Sequence: consensus sequence

<400> 29

Leu	Pro	Xaa	Xaa	Xaa	His	Xaa	Glu	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Gly
1					5				10						15

Asn	Xaa	Thr	Xaa	Xaa	Xaa	His	Xaa	Xaa	Xaa	Xaa	Lys	Xaa	Thr	Xaa	Asn
			20					25					30		

Xaa	Xaa	Gly	Xaa	Xaa	Xaa	Xaa	Ser	Glu	Xaa	Val	Xaa	Xaa	Ser	Xaa	Xaa
		35					40					45			

Xaa	Xaa	Glu
		50

<210> 30

<211> 20

<212> DNA

<213> Artificial Sequence

- 37 -

<220>

<223> Description of Artificial Sequence: primer

<400> 30

cagtgagtgc tgtggagacc

20

<210> 31

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 31

tcttcagtca ggctcctctc

20

<210> 32

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 32

acctgcaatg tgtcgactga g

21

<210> 33

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 33

cacttacagc tgttgggatg

20

<210> 34

<211> 10

<212> PRT

<213> Homo sapiens

<220>

<223> Xaa at position 1 may be any amino acid

<400> 34

Xaa Val Leu Asp Phe Asn Asn Ile Arg Ser
1 5 10

<210> 35

<211> 10

- 38 -

<212> PRT

<213> Homo sapiens

<400> 35

Ser Gln Gly Arg Lys Gly Gln Glu Gly Ser
 1 5 10

<210> 36

<211> 272

<212> PRT

<213> Mus musculus

<400> 36

Met Met Val Val Cys Ala Pro Ala Ala Val Arg Phe Leu Ala Val Phe
 1 5 10 15

Thr Met Met Ala Leu Cys Ser Leu Pro Leu Leu Gly Ala Ser Ala Thr
 20 25 30

Leu Asn Ser Val Leu Ile Asn Ser Asn Ala Ile Lys Asn Leu Pro Pro
 35 40 45

Pro Leu Gly Gly Ala Gly Gly Gln Pro Gly Ser Ala Val Ser Val Ala
 50 55 60

Pro Gly Val Leu Tyr Glu Gly Gly Asn Lys Tyr Gln Thr Leu Asp Asn
 65 70 75 80

Tyr Gln Pro Tyr Pro Cys Ala Glu Asp Glu Glu Cys Gly Ser Asp Glu
 85 90 95

Tyr Cys Ser Ser Pro Ser Arg Gly Ala Ala Gly Val Gly Gly Val Gln
 100 105 110

Ile Cys Leu Ala Cys Arg Lys Arg Arg Lys Arg Cys Met Thr His Ala
 115 120 125

Met Cys Cys Pro Gly Asn Tyr Cys Lys Asn Gly Ile Cys Met Pro Ser
 130 135 140

Asp His Ser His Phe Pro Arg Gly Glu Ile Glu Glu Ser Ile Ile Glu
 145 150 155 160

Asn Leu Gly Asn Asp His Asn Ala Ala Ala Gly Asp Gly Tyr Pro Arg
 165 170 175

Arg Thr Thr Leu Thr Ser Lys Ile Tyr His Thr Lys Gly Gln Glu Gly
 180 185 190

Ser Val Cys Leu Arg Ser Ser Asp Cys Ala Ala Gly Leu Cys Cys Ala
 195 200 205

Arg His Phe Trp Ser Lys Ile Cys Lys Pro Val Leu Lys Glu Gly Gln
 210 215 220

Val Cys Thr Lys His Lys Arg Lys Gly Ser His Gly Leu Glu Ile Phe
 225 230 235 240

His His Gln Ala Ser Asn Ser Ser Arg Leu His Thr Cys Gln Arg His
260 265 270

```
<210> 37
<211> 259
<212> PRT
<213> Xenopus laevis
```

```
<400> 37
Met Gly Ser Asn Met Phe Pro Val Pro Leu Ile Val Phe Trp Gly Phe
  1             5             10             15
```

Ile Leu Asp Gly Ala Leu Gly Phe Val Met Met Thr Asn Ser Asn Ser
20 25 30

Ile Lys Asn Val Pro Ala Ala Pro Ala Gly Gln Pro Ile Gly Tyr Tyr
35 40 45

Pro Val Ser Val Ser Pro Asp Ser Leu Tyr Asp Ile Ala Asn Lys Tyr
50 55 60

Gln Pro Leu Asp Ala Tyr Pro Leu Tyr Ser Cys Thr Glu Asp Asp Asp
65 70 75 80

Cys Ala Leu Asp Glu Phe Cys His Ser Ser Arg Asn Gly Asn Ser Leu
85 90 95

Val Cys Leu Ala Cys Arg Lys Arg Arg Lys Arg Cys Leu Arg Asp Ala
100 105 110

Met Cys Cys Thr Gly Asn Tyr Cys Ser Asn Gly Ile Cys Val Pro Val
115 120 125

Glu Gln Asp Gln Glu Arg Phe Gln His Gln Gly Tyr Leu Glu Glu Thr
130 135 140

Ile Leu Glu Asn Tyr Asn Asn Ala Asp His Ala Thr Met Asp Thr His
145 150 155 160

Ser Lys Leu Thr Thr Ser Pro Ser Gly Met Gln Pro Phe Lys Gly Arg
165 170 175

Asp Gly Asp Val Cys Leu Arg Ser Thr Asp Cys Ala Pro Gly Leu Cys
180 185 190

Cys Ala Arg His Phe Trp Ser Lys Ile Cys Lys Pro Val Leu Asp Glu
195 200 205

Gly Gln Val Cys Thr Lys His Arg Arg Lys Gly Ser His Gly Leu Glu
210 215 220

Ile Phe Gln Arg Cys His Cys Gly Ala Gly Leu Ser Cys Arg Leu Gln
225 230 235 240

Lys Gly Glu Phe Thr Thr Val Pro Lys Thr Ser Arg Leu His Thr Cys

245	250	255
Gln Arg His		
 <210> 38		
<211> 350		
<212> PRT		
<213> Gallus gallus		
 <400> 38		
Met Arg Arg Gly Glu Gly Pro Ala Pro Arg Arg Arg Trp Leu Leu Leu		
1 5 10 15		
Leu Ala Val Leu Ala Ala Leu Cys Cys Ala Ala Ala Gly Ser Gly Gly		
20 25 30		
Arg Arg Arg Ala Ala Ser Leu Gly Glu Met Leu Arg Glu Val Glu Ala		
35 40 45		
Leu Met Glu Asp Thr Gln His Lys Leu Arg Asn Ala Val Gln Glu Met		
50 55 60		
Glu Ala Glu Glu Glu Gly Ala Lys Lys Leu Ser Glu Val Asn Phe Glu		
65 70 75 80		
Asn Leu Pro Pro Thr Tyr His Asn Glu Ser Asn Thr Glu Thr Arg Ile		
85 90 95		
Gly Asn Lys Thr Val Gln Thr His Gln Glu Ile Asp Lys Val Thr Asp		
100 105 110		
Asn Arg Thr Gly Ser Thr Ile Phe Ser Glu Thr Ile Ile Thr Ser Ile		
115 120 125		
Lys Gly Gly Glu Asn Lys Arg Asn His Glu Cys Ile Ile Asp Glu Asp		
130 135 140		
Cys Glu Thr Gly Lys Tyr Cys Gln Phe Ser Thr Phe Glu Tyr Lys Cys		
145 150 155 160		
Gln Pro Cys Lys Thr Gln His Thr His Cys Ser Arg Asp Val Glu Cys		
165 170 175		
Cys Gly Asp Gln Leu Cys Val Trp Gly Glu Cys Arg Lys Ala Thr Ser		
180 185 190		
Arg Gly Glu Asn Gly Thr Ile Cys Glu Asn Gln His Asp Cys Asn Pro		
195 200 205		
Gly Thr Cys Cys Ala Phe Gln Lys Glu Leu Leu Phe Pro Val Cys Thr		
210 215 220		
Pro Leu Pro Glu Glu Gly Glu Pro Cys His Asp Pro Ser Asn Arg Leu		
225 230 235 240		
Leu Asn Leu Ile Thr Trp Glu Leu Glu Pro Asp Gly Val Leu Glu Arg		
245 250 255		

Cys	Pro	Cys	Ala	Ser	Gly	Leu	Ile	Cys	Gln	Pro	Gln	Ser	Ser	His	Ser
			260					265					270		
Thr	Thr	Ser	Val	Cys	Glu	Leu	Ser	Ser	Asn	Glu	Thr	Arg	Lys	Asn	Glu
		275					280					285			
Lys	Glu	Asp	Pro	Leu	Asn	Met	Asp	Glu	Met	Pro	Phe	Ile	Ser	Leu	Ile
	290					295					300				
Pro	Arg	Asp	Ile	Leu	Ser	Asp	Tyr	Glu	Glu	Ser	Ser	Val	Ile	Gln	Glu
305					310					315					320
Val	Arg	Lys	Glu	Leu	Glu	Ser	Leu	Glu	Asp	Gln	Ala	Gly	Val	Lys	Ser
				325					330					335	
Glu	His	Asp	Pro	Ala	His	Asp	Leu	Phe	Leu	Gly	Asp	Glu	Ile		
			340					345					350		

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/05452

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07K14/47 C12N15/12 C07K16/18 G01N33/50 G01N33/53
C12Q1/68 A61K38/17

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 98 46755 A (MCCARTHY SEAN A ;MILLENNIUM BIOTHERAPEUTICS INC (US)) 22 October 1998 (1998-10-22) Note: 100.0% nt seq identity of SEQ ID NO:1 (3) with SEQ ID NO:1 (3) in 2479 bp (1050 bp) overlap, 100.0% aa seq identity of SEQ ID NO:2 with SEQ ID NO:2 in 350 aa overlap. the whole document example 1 page 85, line 31 -page 86, line 7 page 89-95 claims 1-22</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	1-22

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

° Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

5 July 2000

Date of mailing of the international search report

18.10.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

van de Kamp, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/05452

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 98 27932 A (HUMAN GENOME SCIENCES INC ;RUBEN STEVEN M (US); SOPPET DANIEL R (U) 2 July 1998 (1998-07-02) Note: 99.7% nt seq identity of SEQ ID NO:1 with SEQ ID NO:1 (3) in 2445 bp (1050 bp) overlap, 99.1% aa seq identity of SEQ ID NO:2 with SEQ ID NO:2 in 350 aa overlap. the whole document page 2, line 22 -page 3, line 12 page 4, line 8-12 examples 1-4 page 45-49 claims 1-4,8-20 figures 1,2</p> <p>---</p>	1-22
A	<p>GLINKA ET AL: "Dickkopf-1 is a member of a new family of secreted proteins and functions in head induction" NATURE, vol. 391, no. 6665, 22 January 1998 (1998-01-22), pages 357-362, XP002140766 the whole document page 357, right-hand column, line 4-13; figure 1B</p> <p>---</p>	1-22
A	<p>FINCH P W ET AL.: "Purification and molecular cloning of a secreted, Frizzled-related antagonist of Wnt action" PROC. NATL. ACAD. SCI. USA, vol. 94, no. 13, June 1997 (1997-06), pages 6770-6775, XP002140787 abstract; figure 2</p> <p>---</p>	1-22
P,X	<p>KRUPNIK V E ET AL.: "Functional and structural diversity of the human Dickkopf gene family" GENE, vol. 238, no. 2, 1 October 1999 (1999-10-01), pages 301-313, XP002140768 Note: 99.9% (99.7%) nt seq identity of hdkk-3 with SEQ ID NO:1 (3) in 2479 bp (1050 bp) overlap, 99.1% aa seq identity of hDkk-3 with SEQ ID NO:2 in 350 aa overlap. the whole document page 304, left-hand column, line 21-34; figure 1</p> <p>---</p> <p>-/--</p>	1-22

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/05452

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 99 22000 A (DEUTSCHES KREBSFORSCH ;GLINKA ANDREI (DE); NIEHRS CHRISTOF (DE)) 6 May 1999 (1999-05-06) Note: 67.6% nt seq identity of SEQ ID NO:7 with SEQ ID NO:3 in 873 bp overlap, 65.4% aa seq identity of translated SEQ ID NO:7 with SEQ ID NO:2 in 254 aa overlap. the whole document claim 4A; figure 2 ---	1-22
E	WO 00 18194 A (ARNOLD & RICHTER KG ;BRAUCKMANN WILFRIED (DE); STEGMAIER KLAUS DIE) 30 March 2000 (2000-03-30) Note: 99.7% nt seq identity of SEQ ID NO:2 with SEQ ID NO:1 (3) in 1053 bp (1050 bp) overlap, 99.1% aa seq identity of SEQ ID NO:9 with SEQ ID NO:2 in 350 aa overlap. the whole document figures 2,9; examples 2,10,11 claims 1-20 -----	1-22

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 00/05452

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-22 all partially

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1-22 (all partially)

An isolated nucleic acid molecule: a) comprising a nucleotide (nt) sequence which is at least 60% homologous to a nt sequence of SEQ ID NO:1 or 3, or of the DNA insert of the plasmid with ATCC Accession No. 98452, or a complement thereof; or b) comprising at least 1000 nts from said nt sequences; or c) encoding a polypeptide comprising an amino acid (aa) sequence at least about 60% homologous to the aa sequence of SEQ ID NO:2, or to the aa sequence encoded by the insert of said plasmid; or d) encoding a fragment of at least 15 contiguous aa residues of said aa sequence; or e) encoding a naturally occurring allelic variant of said polypeptide, wherein the nucleic acid molecule hybridizes to a nucleic acid molecule comprising SEQ ID NO:1 or 3 or the DNA insert of said plasmid, under stringent conditions. An isolated nucleic acid molecule: f) comprising the nt sequence of SEQ ID NO:1 or 3, or of the DNA insert of the plasmid with ATCC Accession No. 98452; or g) encoding a polypeptide comprising the aa sequence of SEQ ID NO:2, or encoded by the DNA insert of said plasmid. Vectors, host cells. Isolated polypeptides analogous to said isolated nucleic acid molecules (a,c,d,e,g). Fusion proteins. Antibodies. Methods for producing polypeptides. Methods for detecting the presence of said polypeptides. Methods for detecting the presence of said nucleic acid molecules. Methods for identifying compounds which bind to or modulate said polypeptides. Methods for modulating the activity of said polypeptides.

2. Claims: 1-22 (all partially)

As invention 1, but concerning SEQ ID NO:4-6, and the plasmid with ATCC Accession No. ____.

3. Claims: 1-22 (all partially)

As invention 1, but concerning SEQ ID NO:7-9, and the plasmid with ATCC Accession No. 98633.

4. Claims: 1-22 (all partially)

As invention 1, but concerning SEQ ID NO:13-15, and the plasmid with ATCC Accession No. ____.

5. Claims: 1-22 (all partially)

As invention 1, but concerning SEQ ID NO:20-22, and the plasmid with ATCC Accession No. ____.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/05452

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9846755 A	22-10-1998	AU 7137398 A EP 0975755 A	11-11-1998 02-02-2000
WO 9827932 A	02-07-1998	AU 5613498 A EP 0954575 A	17-07-1998 10-11-1999
WO 9922000 A	06-05-1999	DE 19747418 C EP 1027440 A	15-07-1999 16-08-2000
WO 0018194 A	30-03-2000	DE 19844293 A	30-03-2000